

Representations of compact Lie groups and the osculating spaces of their orbits

CLAUDIO GORODSKI* AND GUDLAUGUR THORBERGSSON

INSTITUTO DE MATEMÁTICA E ESTATÍSTICA
UNIVERSIDADE DE SÃO PAULO
RUA DO MATÃO, 1010
SÃO PAULO, SP 05508-900
BRAZIL

E-MAIL: gorodski@ime.usp.br

MATHEMATISCHES INSTITUT
UNIVERSITÄT ZU KÖLN
WEYERTAL 86-90
50931 KÖLN
GERMANY

E-MAIL: gthorber@mi.uni-koeln.de

**Alexander von Humboldt Research Fellow* at the University of Cologne during the completion of this work.

Contents

1	Introduction	4
2	A review of basic definitions and results	6
3	A collection of geometric propositions	8
3.1	Variationally complete representations	8
3.2	Taut representations	12
3.3	A reduction principle	15
4	A necessary condition for a representation to be of class $\mathcal{O}^2$	16
5	The candidates to a position in class $\mathcal{O}^2$	22
5.1	The case where ρ is of quaternionic type	24
5.2	The case where ρ is of complex type	25
5.3	The case where ρ is of real type	30
5.3.1	The case where G has four simple factors	34
5.3.2	The case where G has three simple factors	34
5.3.3	The case where G has two simple factors and ρ has two factors of real type	35
5.3.4	The case where G has two simple factors and ρ has two factors of quaternionic type	35
5.3.5	The case where G is a simple group	36
6	The proof of the converse to a Theorem of Bott and Samelson	50
7	Taut representations	63
8	Other applications	77
A	Description of the maximal subsets of strongly orthogonal roots of the complex simple Lie algebras	79
B	Table for Dadok's invariant $k(\lambda)$	81
C	Table of candidates to class $\mathcal{O}^2$	83

Abstract

Several classes of irreducible orthogonal representations of compact Lie groups that are of importance in Differential Geometry have the property that the second osculating space of all of their nontrivial orbits coincide with the representation space. We say that representations with this property are of class $\mathcal{O}^2$. Our approach in the present paper will be to find restrictions on the class $\mathcal{O}^2$ and then apply them to classify variationally complete and taut representations. The known classifications of cohomogeneity one and two orthogonal representations and more generally of polar representations will also follow easily.

⁰2000 *Mathematics Subject Classification*: primary, 57S15; secondary, 53C30, 53C40, 53C42.

1 Introduction

Several classes of irreducible orthogonal representations of compact Lie groups that are of importance in Differential Geometry have the property that the second osculating space of all of their nontrivial orbits coincide with the representation space. We say that representations with this property are of class $\mathcal{O}^2$. Our approach in the present paper will be to find restrictions on the class $\mathcal{O}^2$ and then apply them to classify variationally complete and taut representations. The known classifications of cohomogeneity one and two orthogonal representations and more generally of polar representations will also follow easily. We refer to Section 2 for the definitions of concepts used in this introduction.

Variationally complete actions were introduced by Bott in [4]. Bott and Samelson proved in [5] that isotropy representations of symmetric spaces are variationally complete. We will prove the following converse of their theorem in Section 6.

1.1 Theorem *A variationally complete representation of a compact connected Lie group is orbit equivalent to the isotropy representation of a symmetric space.*

We will in fact prove more than we have stated here and give a complete classification of variationally complete representations (up to image equivalence; see the beginning of Section 5 for this concept). As a corollary we get a new proof of the classification of polar representations due to Dadok ([12]) since these are variationally complete (see [9]) and isotropy representations of symmetric spaces are easily seen to be polar. It follows that an orthogonal representation is variationally complete if and only if it is polar, and it is polar if and only if it is orbit equivalent to the isotropy representation of a symmetric space.

Another important result of Bott and Samelson in [5] is that the distance functions to orbits of variationally complete representations are perfect Morse functions. We say that submanifolds with this property are *taut* and call representations *taut* if all of their orbits are taut. In Section 7 we will prove the following classification theorem which provides new examples of taut submanifolds and taut representations.

1.2 Theorem *A taut irreducible representation ρ of a compact connected Lie group G is either orbit equivalent to the isotropy representation of a symmetric space or it is one of the following orthogonal representations ($n \geq 2$):*

G	ρ
$\mathbf{SO}(2) \times \mathbf{Spin}(9)$	$(standard) \otimes_{\mathbf{R}} (spin)$
$\mathbf{U}(2) \times \mathbf{Sp}(n)$	$(standard) \otimes_{\mathbf{C}} (standard)$
$\mathbf{SU}(2) \times \mathbf{Sp}(n)$	$(standard)^3 \otimes_{\mathbf{H}} (standard)$

As a consequence of this theorem, the classes of polar, variationally complete and taut irreducible representations of compact *simple* groups coincide. Notice that the three exceptional representations in Theorem 1.2 are precisely the irreducible representations of cohomogeneity three that are not polar (see [46]). Hence, in the case of a general compact Lie group, the taut irreducible representations are precisely those that are either polar or of cohomogeneity three. A further related result is the classification of homogeneous taut submanifolds with flat normal bundle due to Olmos ([32]). He shows that such a submanifold is a principal

orbit of the isotropy representation of a symmetric space. It follows that the principal orbits of the three nonpolar representations in Theorem 1.2 do not have flat normal bundle which is also easy to see directly.

Bott and Samelson proved the tautness of variationally complete representations as an application of their K -cycles which are concrete representatives of a basis for the homology of the orbits. (In [5] the group acting is denoted by K which explains the terminology ‘ K -cycle’). The three nonpolar representations in Theorem 1.2 admit generalizations of the K -cycles of Bott and Samelson. We will not verify this claim in the present paper since it requires methods quite different from those employed here and would make the paper too long; we refer instead to the forthcoming paper [20]. Instead we will prove their tautness with shorter, although maybe less illuminating arguments based on a theorem of Floyd ([17]). This approach is adapted from the paper [14] of Duistermaat.

Kuiper observed in [27] that the second osculating space of a taut submanifold in a Euclidean space V coincides with V if the submanifold is not contained in a proper affine subspace. In fact, he proved this more generally for tight submanifolds, but this is unimportant for us since an orbit is tight if and only if it is taut. Since the classes of representations we are dealing with are all taut, it follows from this observation of Kuiper that they belong to class $\mathcal{O}^2$ if they are irreducible. The class $\mathcal{O}^2$ is much more tractable than the other classes of representations we are dealing with since it involves an infinitesimal condition. The technique of Dadok ([12]), notably his invariant $k(\lambda)$, turns out to be an extremely powerful tool to reduce the class $\mathcal{O}^2$ in size so that the remaining cases are accessible to the geometric methods developed in Section 3.

We have already pointed out that Dadok’s classification of polar representations follows as a consequence of Theorem 1.1. Some other classification results can also easily be proved with our methods as shown in Section 8. These include the classification of cohomogeneity one representations as well as the classification of cohomogeneity two representations due to Hsiang-Lawson ([26]). The cohomogeneity two representations are polar and therefore included in Dadok’s classification, but the point here is that it is very easy to see directly that they belong to class $\mathcal{O}^2$ without referring to tautness.

One could also use our methods to classify symmetric spaces and their isotropy representations. This would not lead to a simpler proof than those existing in the literature. Insisting on the details needed to classify symmetric spaces and their isotropy representations would make the paper much longer and more involved than it is. We will therefore use this classification in the paper¹ and assume knowledge of the lists in [45], Tables 8.11.2 and 8.11.5. (See also [24], but notice that the isotropy representations are not listed in this reference). Otherwise, our guiding principle in writing the paper has been to make it as self-contained as possible.

Previous to this paper taut representations were studied in [19] and [11]. It is decided in [19], with some exceptions, which representations of $\mathbf{SU}(n)$ and $\mathbf{U}(n)$ can be taut. In [11] it is proved among other things that a compact group admitting an almost faithful taut representation can have at most four simple factors. Eschenburg and Heintze use the theory of isoparametric submanifolds in [16] to reprove Dadok’s characterization up to orbit equivalence of polar representations under the assumption of cohomogeneity at least three. In [15]

¹Except for the classification of cohomogeneity one representations.

they use the same theory to decide which irreducible representations are orbit equivalent to isotropy representations of symmetric spaces. This follows immediately from our work in Section 6. Bergmann in [1] decides which reducible representations are polar. Her result is equivalent to the last step in our classification of variationally complete representations. Di Scala and Olmos give in [13] a direct proof of the fact that a variationally complete representation is polar.

The first author wishes to thank the *Alexander von Humboldt Foundation* for its generous support and constant assistance during the completion of this work.

2 A review of basic definitions and results

Let G be a compact Lie group acting on a Riemannian manifold M by isometries. A geodesic γ in M is called *G -transversal* if it is orthogonal to the G -orbit through $\gamma(t)$ for every t . One can show that a geodesic γ is G -transversal if there is a point t_0 such that $\dot{\gamma}(t_0)$ is orthogonal to $G\gamma(t_0)$. A Jacobi field along a geodesic in M is called *G -transversal* if it is the variational vector field of a variation through G -transversal geodesics. The action of G on M is called *variationally complete* if every G -transversal Jacobi field J in M that is tangent to the G -orbits at two different parameter values is the restriction of a Killing field on M induced by the G -action (see [4] and [5]). It is proved in [5] on p. 974 that instead of requiring tangency at two different points in the definition of variational completeness it is equivalent to require tangency at one point and vanishing at another point.

Let ρ be a variationally complete reducible representation and $\tilde{\rho}$ a summand of ρ . Then it is easy to see that $\tilde{\rho}$ is also variationally complete, see Proposition 3.4. Notice though that it is not true that the direct sum of two variationally complete representations of a compact Lie group G is variationally complete as can be seen by taking the direct sum of two copies of $\mathbf{SO}(2)$ acting on $\mathbf{R}^2$.

Let N be a properly embedded submanifold of a Euclidean space V and let x be some point in V . Then we define the *distance function* $L_x : N \rightarrow \mathbf{R}$ from x to N by setting $L_x(p) = \|p - x\|^2$. It follows that L_x is a non-negative proper function since N is properly embedded. Hence it is possible to apply Morse theory to L_x . We say that L_x is *perfect with respect to a field F* if it is a Morse function and the Morse inequalities for L_x with respect to F are equalities. Furthermore we say that N is *F -taut* or simply *taut* if L_x is perfect with respect to F whenever L_x is a Morse function, see [8]. The concept of tautness can be extended to submanifolds of complete Riemannian manifolds, see [43], but we will not need that here. We will say that an orthogonal representation $\rho : G \rightarrow \mathbf{O}(V)$ of a compact Lie group G is *F -taut* or simply *taut* if the the orbits of G are F -taut submanifolds of V .

If ρ is a reducible taut representation and $\tilde{\rho}$ one of its factors, then $\tilde{\rho}$ is clearly taut. Notice that it is not true that direct sums of taut representations of a compact Lie group G are also taut. An example of this is the direct sum of n copies of $\mathbf{SU}(n)$ acting on $\mathbf{C}^n$. This representation is not taut since one can compute that $\mathbf{SU}(n)$ is not taut in the matrix space $M(n; \mathbf{C})$.

The following theorem mentioned in the introduction was proved by Bott and Samelson more generally for variationally complete actions on complete Riemannian manifolds. In fact it is an immediate corollary of Theorem I in [5]. Notice that Bott and Samelson do not use

the concept of a taut submanifold which was introduced later.

2.1 Theorem (Bott-Samelson) *A variationally complete representation of a compact connected Lie group on an Euclidean space is $\mathbf{Z}_2$ -taut.*

Bott proved in [4], Propositions 7.1 and 11.6, that the action of a compact Lie group G with a bi-invariant metric on itself by conjugations and the adjoint representation of G on its Lie algebra $\mathfrak{g}$ are variationally complete. This was generalized by Bott and Samelson in [5], Theorem II as follows. Let (L, G) be a symmetric pair and $\mathfrak{l} = \mathfrak{g} \oplus \mathfrak{p}$ the corresponding Cartan decomposition. Then the action of $L \times L$ on L , the action of G on L/G and the action of $\text{Ad}_L(G)$ restricted to $\mathfrak{p}$ are variationally complete. The action of $\text{Ad}_L(G)$ on $\mathfrak{p}$ is equivalent to the isotropy representation of G on the tangent space $T_p(L/G)$ where p denotes the coset G . Therefore we have the following theorem mentioned in the introduction.

2.2 Theorem (Bott-Samelson) *The isotropy representation of a symmetric space is variationally complete.*

Conlon considered in [9] actions of a Lie group G on a complete Riemannian manifold M with the property that there is a connected submanifold Σ of M that meets all orbits of G in such a way that the intersections between Σ and the orbits of G are all orthogonal. Such a submanifold is called a *section* and an action admitting a section is now usually called *polar* if Σ is properly embedded. Notice that Conlon does not assume in [9] that Σ is properly embedded, but it is usually required in the recent literature on the subject. It is easy to see that a section Σ is totally geodesic in M . An action admitting a section that is flat in the induced metric is called *hyperpolar*. There is clearly no difference between polar and hyperpolar representations since totally geodesic submanifolds of a Euclidean space are affine subspaces. Moreover, the question whether Σ should be required to be properly embedded or not becomes redundant. Conlon proved the following theorem in [9].

2.3 Theorem (Conlon) *A hyperpolar action of a compact Lie group on a complete Riemannian manifold is variationally complete.*

Polar representations were classified by Dadok in [12]. We recall that two representations $\rho_1 : G_1 \rightarrow O(V_1)$ and $\rho_2 : G_2 \rightarrow O(V_2)$ are said to be *orbit equivalent* if there is an isometry $A : V_1 \rightarrow V_2$ under which the orbits of G_1 and G_2 correspond. As a consequence of his classification he obtained the following result.

2.4 Theorem (Dadok) *A polar representation of a compact connected Lie group is orbit equivalent to the isotropy representation of a symmetric space.*

We can summarize this discussion as follows. Let G be a compact connected Lie group. Denote by $\mathcal{I}$ the representations of G that are isotropy representations of symmetric spaces, by $\mathcal{P}$ those that are polar, by $\mathcal{V}$ those that are variationally complete, and by $\mathcal{T}$ those that are taut. Then

$$\mathcal{I} \subset \mathcal{P} \subset \mathcal{V} \subset \mathcal{T}.$$

The starting point in the present paper will be to consider a class of representations that is a priori larger than $\mathcal{T}$, but easier to deal with, see [11]. We recall that the *second osculating*

space $O_p^2(N)$ at a point p of a submanifold N in a Euclidean space V is the vector space spanned by the first and second derivatives at p of the inclusion of N into V . It is easy to see that

$$O_p^2(N) = T_p N \oplus \langle \{ \alpha(X, Y) \mid X, Y \in T_p N \} \rangle.$$

where α denotes the second fundamental form of N and $\langle S \rangle$ stands for the linear hull of the set S . The following is a corollary of the discussion of Kuiper in [27].

2.5 Theorem (Kuiper) *Let N be a taut submanifold of a Euclidean space. Then the affine hull of N coincides with $p + O_p^2(N)$ for every point p in N .*

We let $\mathcal{O}^2$ denote the class of representations of a compact connected Lie group G such that the representation space coincides with $O_p^2(Gp)$ for all nonzero p . The representations in $\mathcal{O}^2$ are clearly irreducible. If a taut representation is irreducible, then Theorem 2.5 implies that it belongs to class $\mathcal{O}^2$. Thus we have the inclusions

$$\mathcal{I}_i \subset \mathcal{P}_i \subset \mathcal{V}_i \subset \mathcal{T}_i \subset \mathcal{O}^2,$$

where $\mathcal{I}_i$, $\mathcal{P}_i$, $\mathcal{V}_i$, and $\mathcal{T}_i$ are the subclasses of $\mathcal{I}$, $\mathcal{P}$, $\mathcal{V}$, and $\mathcal{T}$ consisting of irreducible representations.

3 A collection of geometric propositions

In this section we collect technical results of geometric nature that will be needed to prove our main theorems.

3.1 Variationally complete representations

We first study focal points of orbits of variationally complete representations with the aim to prove that a representation that is orbit equivalent to a variationally complete one is also variationally complete. We then show how reducible variationally complete representations relate to irreducible ones. Finally we prove an important criterion which can be used to decide in some cases whether a variationally complete representation is orbit equivalent to the isotropy representation of a symmetric space.

Let q be a focal point of an orbit Gp of an orthogonal representation of a compact connected Lie group G . It is clear that the orbit Gq consists of focal points of Gp . We therefore call Gq a *focal orbit of Gp* . It is also clear that the singular orbits of a representation are always focal orbits of the principal orbits. Slightly more generally, Gq is a focal orbit of Gp if the segment $\overline{qp}$ is perpendicular to Gp in p and there is a one-parameter group in G_q that does not fix p . We will use the following lemma to prove Proposition 3.3.

3.1 Lemma *A representation $\rho : G \rightarrow \mathbf{O}(V)$ is variationally complete if and only if every orbit Gp has the following property: if q is a focal point of Gp relative to p with multiplicity $k > 0$, then $k = \dim G_q p$.*

3.2 Remark (Compare [42].) Let $\rho : G \rightarrow \mathbf{O}(V)$ be a variationally complete representation of a compact connected Lie group G . Then Lemma 3.1 implies that the focal orbits of any orbit of G are singular orbits. Furthermore, the multiplicity of the focal points of a principal orbit Gp in a singular orbit Gq is equal to the difference between the dimension of Gp and the dimension of Gq . In particular, it follows that any principal orbit is a *totally focal submanifold*; see [42] for a definition of this concept. Terng claims in [42] that this fact can be used to show that variationally complete representations are polar. However, as Terng herself pointed out in [43], p. 197, there is a gap in that argument.

Proof of Lemma 3.1. We first assume that ρ is variationally complete. Let q be a focal point of Gp along the line segment $\overline{pq}$ with multiplicity k . Let $\mathcal{J}$ be the space of Jacobi fields along $\overline{pq}$ that are the variational vector fields of variations γ_s of $\overline{pq}$ through line segments such that all γ_s meet the orbit Gp orthogonally for $t = 0$ and the Jacobi field J of the variation of γ vanishes for $t = 1$. The multiplicity k of the focal point q is equal to the dimension of the space $\mathcal{J}$. The action of G_q on $\overline{pq}$ induces variations of $\overline{pq}$ whose variational vector fields are contained in $\mathcal{J}$. Let us denote by $\hat{\mathcal{J}}$ the subspace of $\mathcal{J}$ that consists of Jacobi fields induced by the action of G_q on $\overline{pq}$. It follows that $\dim \hat{\mathcal{J}} = \dim G_q v \leq k$ where v is the tangent vector of $\overline{pq}$ at q . We now prove that $\dim G_q v = k$. Let γ_s be a variation of $\overline{pq}$ through line segments such that all γ_s meet the orbit Gp orthogonally for $t = 0$ and the Jacobi field J of the variation along γ vanishes for $t = 1$. The segments γ_s are orthogonal to the orbit Gp . Hence they are G -transversal and J is a G -transversal Jacobi field. It is clear that J is tangent to the orbits through $p = \gamma(0)$ and $q = \gamma(1)$. By variational completeness, there is a Killing field X on M induced by the G -action such that $J = X \circ \gamma$. Hence $J \in \hat{\mathcal{J}}$ and it follows that $\hat{\mathcal{J}} = \mathcal{J}$. Thus $\dim G_q v = k$.

To prove the converse, we have to show that $\dim G_q v = k$ for all focal points q implies that ρ is variationally complete. This is equivalent to show that a Jacobi field J that vanishes in a point q and is tangent to an orbit Gp in p is induced by the group action. We let the symbols $\mathcal{J}$ and $\hat{\mathcal{J}}$ have the same meaning as in the first part of the proof. We have to show that $\hat{\mathcal{J}} = \mathcal{J}$. This is clear since $\dim \mathcal{J} = k$ and $\dim \hat{\mathcal{J}} = \dim G_q v = k$. $\square$

The following proposition that we will use to classify variationally complete representations is now easy.

3.3 Proposition *Let ρ be a representation that is orbit equivalent to a variationally complete representation. Then ρ is itself variationally complete.*

Proof. We can assume that we have representations ρ and $\hat{\rho}$ of the groups G and $\hat{G}$ with the same representation space and the same orbits and that $\hat{\rho}$ is variationally complete. Clearly the focal orbits of the two representations coincide since their orbits are the same. Let v be a normal vector in q of the orbit $Gq = \hat{G}q$. It follows from the slice theorem and the fact that orbits of ρ and $\hat{\rho}$ are the same that $G_q v = \hat{G}_q v$. Since $\hat{\rho}$ is variationally complete, Lemma 3.1 now implies that ρ is also variationally complete. $\square$

The next proposition will be important in the proof of Theorem 1.1 in the introduction. It is analogous to a result of Dadok on polar representations, see Theorem 4 in [12].

3.4 Proposition *Let G be a compact connected Lie group. Let $\rho : G \rightarrow \mathbf{O}(V)$ be a non-trivial reducible variationally complete representation and let $\rho = \rho_1 \oplus \rho_2$ be a nontrivial decomposition. Let V_1 and V_2 be the representation spaces of ρ_1 and ρ_2 respectively. Then we have the following:*

- (a) *The representations ρ_1 and ρ_2 are variationally complete.*
- (b) *The restriction of ρ_2 to the connected component of the isotropy subgroup of any point in V_1 is orbit equivalent to ρ_2 .*
- (c) *The representations ρ_1 and ρ_2 are not equivalent as representations.*
- (d) *The representation ρ is orbit equivalent to the representation $\hat{\rho} : G \times G \rightarrow \mathbf{O}(V)$ defined by $\hat{\rho}(g_1, g_2) = \rho_1(g_1) + \rho_2(g_2)$.*

Proof. The claim in part (a) is clear, since the orbits of ρ_1 and ρ_2 are also orbits of ρ .

To prove part (b), note that it is true in general, without assuming ρ is variationally complete, that any point $p_1 \in V_1$ is a focal point of any orbit $Gp_2 \subset V_2$ with multiplicity equal to $\dim Gp_2$. Now if ρ is variationally complete then the claim follows from Lemma 3.1 and the fact that the G -orbits are connected.

Next we prove (c). Assume $A : V_1 \rightarrow V_2$ is an isomorphism such that $\rho_2 = A\rho_1A^{-1}$. Notice that $G_p = G_{Ap}$ for all $p \in V_1$. Part (b) implies that G_p is transitive on the orbit $G(Ap)$, which is impossible if we choose $p \in V_1$ such that $\dim Gp > 0$. Therefore such an isomorphism A does not exist.

Finally we prove part (d). Let $v = v_1 + v_2$ be a point in V and v_1 and v_2 its components in V_1 and V_2 respectively. Let H_1 (resp. H_2) be the isotropy group of v_1 (resp. v_2). Notice that v_2 is a focal point of Gv_1 with multiplicity $\dim Gv_1$. It follows from part (b) that $H_2v_1 = Gv_1$ and similarly that $H_1v_2 = Gv_2$. We next prove the factorization $G = H_1H_2$. Let $g \in G$. Then there is an element $h_1 \in H_1$ such that $h_1v_2 = gv_2$. Then $h_1^{-1}g \in H_2$. Set $h_2 = h_1^{-1}g$. Then $g = h_1h_2$. It follows that

$$\mathfrak{g} = \mathfrak{h}_1 + \mathfrak{h}_2$$

where $\mathfrak{g}$, $\mathfrak{h}_1$ and $\mathfrak{h}_2$ are the Lie algebras of G , H_1 and H_2 respectively. It is clear that the orbit of v under $\rho(G)$ is contained in its orbit under $\hat{\rho}(G \times G)$. The tangent space of $\hat{\rho}(G \times G)v$ at v is

$$(\mathfrak{g} \times \mathfrak{g})v = \mathfrak{g}v_1 + \mathfrak{g}v_2 = \mathfrak{h}_2v_1 + \mathfrak{h}_1v_2.$$

Let $X_1 \in \mathfrak{h}_1$ and $X_2 \in \mathfrak{h}_2$. Then $X_1v_1 = 0$ and $X_2v_2 = 0$ and hence

$$X_1v_2 + X_2v_1 = (X_1 + X_2)v \in \mathfrak{g}v.$$

Hence the dimension of the tangent spaces of the orbits of v under ρ and $\hat{\rho}$ coincide. Since G is connected the orbits coincide. Since v was chosen arbitrarily, all orbits ρ and $\hat{\rho}$ coincide. $\square$

3.5 Remark Part (c) in Proposition 3.4 generalizes Proposition 2.10 in [23].

The next lemma is a preparation for Proposition 3.7. We follow the terminology of Bredon in [7], pp. 180-181, and distinguish between three types of orbits, namely principal (regular), exceptional and singular. Recall that orbits are called *exceptional* if they have maximal dimension without being principal, and *singular* if their dimension is not maximal. We also call points *regular* when they belong to principal orbits.

3.6 Lemma *Let G be a closed subgroup of the compact connected Lie group K and let $\rho : K \rightarrow \mathbf{O}(V)$ be a representation. If ρ has an orbit of maximal dimension in common with $\rho|G$, then ρ and $\rho|G$ are orbit equivalent.*

Proof. We denote the maximal dimension of an orbit of ρ by n . Let us denote by $\mathcal{O}_G$ the union of the G -orbits of maximal dimension and by $\mathcal{O}_K$ the corresponding set relative to K . It is well-known that both $\mathcal{O}_G$ and $\mathcal{O}_K$ contain an open and dense subset of V . Hence the set $\mathcal{O}_G \cap \mathcal{O}_K$ also contains an open and dense set. The dimensions of the orbits of G and K through points in $\mathcal{O}_G \cap \mathcal{O}_K$ are all equal to n . It follows that the orbits of G and K through points in $\mathcal{O}_G \cap \mathcal{O}_K$ coincide since K is connected. Let $p \in V \setminus \mathcal{O}_G \cap \mathcal{O}_K$. We would like to show that $Gp = Kp$. Let $k(p)$ be some other point in Kp . Let (p_n) be a sequence in $\mathcal{O}_G \cap \mathcal{O}_K$ that converges to p . Then $(k(p_n))$ converges to $k(p)$. There is for every n an element g_n in G such that $g_n(p_n) = k(p_n)$. Now $(g_n(p_n))$ converges to $k(p)$. By going to a subsequence if necessary we can assume that g_n converges to an element g in G , so that the continuity of $G \times V \rightarrow V$ implies that $(g_n(p_n))$ converges to $g(p)$. Hence $g(p) = k(p)$ and we have seen that all orbits of G and K coincide. $\square$

3.7 Proposition *Let G be a closed subgroup of the compact connected Lie group K and let $\rho : K \rightarrow \mathbf{O}(V)$ be the isotropy representation of an irreducible symmetric space. If the restriction $\rho|G$ is variationally complete and $\rho|G$ has a nontrivial orbit in common with ρ , then ρ and $\rho|G$ are orbit equivalent.*

Proof. Let $p \in V$ be a point such that the orbits of G and K through p coincide and have the largest possible dimension with this property. Assume that Kp is not an orbit of maximal dimension for the action of K . Let $\mathfrak{a}$ be a Cartan subspace in V with respect to the symmetric space representation of K such that $p \in \mathfrak{a}$. Let T be a tubular neighborhood of the common orbit $Kp = Gp$. Let $q \in \mathfrak{a}$ be a point in $Gp = Kp$ different from p and such that the segment $\overline{pq}$ from p to q passes through a point r in T with $\dim K_p r \geq 1$. (Here it is important that the symmetric space is irreducible.) It follows that $\dim Kp < \dim Kr$ by the slice theorem. Hence $\dim Gr < \dim Kr$ since the dimension of $Gp = Kp$ is maximal among the common orbits of G and K . It follows that $\dim G_p r < \dim K_p r$.

Let X be an element of the Lie algebra $\mathfrak{k}_p$ of K_p such that $\dot{\gamma}_{X,r}(0)$ is not in $T_r(G_p r)$ where $\gamma_{X,r}(s) = \exp(sX)r$. Now let us consider the variation $(\overline{pq})_s = \exp(sX)(\overline{pq})$ of $\overline{pq}$. Let J be the corresponding Jacobi field along $\overline{pq}$. This variation is K -transversal and hence also G -transversal. The value of J over p and q are tangent vectors of the common orbit $Gp = Kp$. By the variational completeness of the action of G , the Jacobi field J is the restriction of a Killing field on V to $\overline{pq}$ that is induced by the action of G on V . Hence all values of J are tangent vectors to orbits of G . The value of J over r is $\dot{\gamma}_{X,r}(0)$ which is not a tangent vector of the orbit of G through r . This is a contradiction. Hence G and K have an orbit in

common which is of maximal dimension among the K -orbits. It now follows from Lemma 3.6 that all orbits of G and K coincide. $\square$

3.2 Taut representations

We will first prove that the slice representation of a taut representation is also taut and end the section with an important proposition about taut reducible representations. It also follows immediately from our methods that a taut representation does not have exceptional orbits.

3.8 Proposition *Let G be a compact connected Lie group with a taut representation $\rho : G \rightarrow \mathbf{O}(V)$. Let $p \in V$ and let $\nu_p : G_p \rightarrow \mathbf{O}(N_p(Gp))$ be the slice representation of ρ at p . Then ν_p is taut.*

Proof. Let $N^\epsilon(Gp)$ denote the bundle of normal vectors of length less than ϵ over the orbit Gp . Let $\epsilon > 0$ be so small that $T = \exp(N^\epsilon(Gp))$ is a tubular neighborhood around the orbit Gp . Let $v \in N_p(Gp)$. We want to show that $G_p v$ is taut in the normal space $N_p(Gp)$. Let $\alpha > 0$ be a number that is so small that $w = \alpha v$ has length less than ϵ . It is clear that $G_p v$ is taut in $N_p(Gp)$ if and only if $G_p w$ is taut in $N_p(Gp)$. It is also clear that $G_p w$ is taut in $N_p(Gp)$ if and only if $\exp_p(G_p w) = G_p \exp_p(w)$ is taut in $\exp_p(N_p(Gp))$ since $\exp_p$ is an isometry. We set $q = \exp_p(w)$. A submanifold in an affine subspace A of V is taut in A if and only if it is taut in V . Hence $G_p v$ is taut in $N_p(Gp)$ if and only if $G_p q$ is taut in V .

Now let $L_p : Gq \rightarrow \mathbf{R}$ be the distance function. The segment $\overline{pq}$ is orthogonal to Gp and hence also to Gq . It follows that q is a critical point of L_p . Since q is in the tubular neighborhood T and the length of the segment $\overline{pq}$ is smaller than ϵ it follows that L_p takes on its minimum in q . Let C be the subset of Gq on which L_p takes on its minimum value. Clearly $C = Gq \cap \exp_p(N_p^\epsilon(Gp))$ since a geodesic segment between Gp and Gq that is orthogonal to Gp is orthogonal to Gq and vice versa. By the slice theorem $G_p q = Gq \cap \exp_p(N_p^\epsilon(Gp))$. Hence $G_p q = C$. The set of critical points of a distance function to a taut submanifold is a union over taut submanifolds by a theorem of Ozawa, see [34]. It follows that $G_p q$ is taut and hence that the slice representation ν_p is taut. $\square$

3.9 Remark We are assuming in Proposition 3.8 that G is connected. Hence its orbits are connected and it follows from their tautness that the set of points where a distance function takes on its minimal value is connected. The proof of Proposition 3.8 now implies that the orbits of G_p are connected even if G_p is not connected.

It is well-known that the isotropy representation of a symmetric space does not have an exceptional orbit. The methods used in the proof of Proposition 3.8 immediately give that this is also the case for taut representations, as we show in the next proposition.

3.10 Proposition *A taut representation $\rho : G \rightarrow \mathbf{O}(V)$ of a compact connected Lie group G does not have exceptional orbits.*

Proof. Assume the orbit through p in V is exceptional. Then the slice representation of G_p at p has a disconnected orbit. Arguing exactly as in the proof of Proposition 3.8, we see that the distance function from p to a principal orbit through some regular point close to

p assumes its minimum value on a disconnected set, contradicting tautness. Thus ρ cannot have an exceptional orbit. $\square$

We close this subsection with a discussion of taut reducible representations.

3.11 Proposition *Let ρ_1 and ρ_2 be representations of a compact connected Lie group G with representation spaces V_1 and V_2 respectively. Assume that $\rho_1 \oplus \rho_2$ is F -taut. Then the restriction of ρ_2 to the isotropy group G_{v_1} is taut for every $v_1 \in V_1$.*

Furthermore, we have $p(G(v_1, v_2); F) = p(Gv_1; F) p(G_{v_1}v_2; F)$ where $p(M; F)$ denotes the Poincaré polynomial of M with respect to the field F . In particular, $G_{v_1}v_2$ is connected and

$$b_1(G(v_1, v_2); F) = b_1(Gv_1; F) + b_1(G_{v_1}v_2; F)$$

where $b_1(M; F)$ denotes the first Betti number of M with respect to F .

Proof. We can work with height functions instead of distance functions in this proof since a submanifold contained in a round sphere in a Euclidean space is taut if and only if all height functions are perfect Morse functions, see [35]. Furthermore, in this situation the set of critical points of a distance function will also occur as the set of critical points of a height function, and vice versa. Fix $(v_1, v_2) \in V_1 \oplus V_2$. Let $a \in V_1$ be such that the height function $h_a : Gv_1 \rightarrow \mathbf{R}$ defined by $h_a(v) = \langle a, v \rangle$ is a Morse function. We define the height function $h_{(a,0)} : G(v_1, v_2) \rightarrow \mathbf{R}$ similarly. The point (u_1, u_2) is a critical point of $h_{(a,0)}$ with index i if and only if u_1 is a critical point of h_a with index i . Hence the critical set C on the critical level $h_{(a,0)}(u_1, u_2) = h_a(u_1)$ is

$$C = \{(w_1, w_2) \in G(v_1, v_2) \mid w_1 = u_1\} = \{(u_1, w_2) \mid w_2 \in G_{u_1}v_2\}.$$

Ozawa proves in [34] that the set of critical points of a distance function on a taut submanifold is a union over nondegenerate critical submanifolds that are again taut submanifolds. It follows that C is taut. The projection of C into V_2 , which coincides with $G_{u_1}v_2$, is then also taut. We can choose a such that h_a is a Morse function of which v_1 is a critical point. It follows that $G_{v_1}v_2$ is taut for every $v_1 \in V_1$ and $v_2 \in V_2$ and hence that the restriction of ρ_2 to G_{v_1} is taut.

Now fix again a point (v_1, v_2) . If $C \subset G(v_1, v_2)$ is an arbitrary critical submanifold of $h_{(a,0)}$, then it is diffeomorphic to $G_{v_1}v_2$. To see this let $\pi_1 : G(v_1, v_2) \rightarrow Gv_1$ be the projection onto the first factor. Then one easily sees that

$$\pi_1^{-1}(gv_1) = G_{gv_1}gv_2 = gG_{v_1}g^{-1}(gv_2) = gG_{v_1}v_2.$$

Hence we have that π_1 is a G -equivariant fibration. The critical submanifolds of $h_{(a,0)}$ are fibers of π_1 and hence diffeomorphic to each other.

Since the orbit $G(v_1, v_2)$ is taut we have by [34] that the Morse-Bott inequalities are equalities. Let $C_1, \dots, C_k$ be the critical manifolds of $h_{(a,0)}$. Let $i(C_j)$ be the index of C_j for $j = 1, \dots, k$. Then

$$p(G(v_1, v_2)) = \sum_{j=1}^k p(C_j) t^{i(C_j)} = p(G_{v_1}v_2) \sum_{j=1}^k t^{i(C_j)}.$$

since the critical submanifolds are all diffeomorphic to $G_{v_1}v_2$. We have

$$p(Gv_1) = \sum_{j=1}^k t^{i(C_j)}$$

since h_a is a perfect Morse functions with critical points $p_1 = \pi_1(C), \dots, p_k = \pi_1(C_k)$ and the index of p_j is equal to $i(C_j)$. It follows that

$$p(G(v_1, v_2)) = p(Gv_1) p(G_{v_1}v_2).$$

Multiplying out the Poincaré polynomials gives $b_0(G_{v_1}v_2) = 1$ and

$$b_1(G(v_1, v_2)) = b_1(Gv_1) + b_1(G_{v_1}v_2).$$

This finishes the proof of the proposition. $\square$

3.12 Remark The property that $\rho_2|_{G_{v_1}}$ is taut in Proposition 3.11 is a weakening of the property that $\rho_2|_{G_{v_1}}$ is orbit equivalent to ρ_2 for a variationally complete reducible representation $\rho_1 \oplus \rho_2$ (cf. Proposition 3.4, part (b)).

We give examples of how Proposition 3.11 can be used.

3.13 Examples (i) Let $G = \mathbf{SO}(n)$ and let ρ_1 be the $\mathbf{SO}(n)$ -conjugation on the space V_1 of real traceless symmetric $n \times n$ matrices. Then ρ_1 is taut since it is the isotropy representation of the symmetric space $\mathbf{SL}(n, \mathbf{R})/\mathbf{SO}(n)$. Let ρ_2 be any other nontrivial representation of $\mathbf{SO}(n)$ with representation space V_2 . Then $\rho_1 \oplus \rho_2$ cannot be taut if $n \geq 3$. To see this, let $v_1 \in V_1$ be a regular point. Then G_{v_1} is the discrete group consisting of all diagonal matrices with determinant one and entries ± 1 on the diagonal. The kernel of ρ_2 is contained in the center of $\mathbf{SO}(n)$. Since $n \geq 3$, we see that G_{v_1} cannot be contained in the center of ρ_2 . Hence there is an element $v_2 \in V_2$ that is not fixed by G_{v_1} . It follows that $G_{v_1}v_2$ is disconnected. Now Proposition 3.11 implies that $\rho_1 \oplus \rho_2$ is not taut.

(ii) Now let G be a compact connected simple Lie group of rank at least two and let ρ_1 denote the adjoint representation of G . We assume that G is simply connected. Let ρ_2 be any other nontrivial representation of G . Then $\rho_1 \oplus \rho_2$ is not taut. To see this let T be a maximal torus in G . We denote the representation spaces of ρ_1 and ρ_2 by V_1 and V_2 respectively. There is a regular element $v_1 \in V_1$ with $G_{v_1} = T$. The restriction of ρ_2 to T has a discrete kernel that is contained in the center of G . If $v_2 \in V_2$ is a T -regular point then the isotropy subgroup T_{v_2} coincides with the kernel of $\rho_2|_T$. Hence $G_{v_1}v_2$ is diffeomorphic to T and it follows that $b_1(G_{v_1}v_2; F)$ is equal to the rank of G . In particular $b_1(G_{v_1}v_2; F) \geq 2$. Now notice that the isotropy group of (v_1, v_2) is also T_{v_2} . Hence $\pi_1(G(v_1, v_2)) = T_{v_2}$ which implies $H_1(G(v_1, v_2); \mathbf{Z}) = T_{v_2}$ since T_{v_2} is Abelian. If $G \neq \mathbf{Spin}(4k)$ then the center of G is a cyclic group and it follows that $b_1(G(v_1, v_2); F) \leq 1$. If $G = \mathbf{Spin}(4k)$, then $k \geq 2$ and we get $b_1(G_{v_1}v_2; F) = 2k \geq 4$; since the center of $\mathbf{Spin}(4k)$ is $\mathbf{Z}_2 \times \mathbf{Z}_2$, we have $b_1(G(v_1, v_2); F) \leq 2$. In either case, $b_1(G_{v_1}v_2; F) > b_1(G(v_1, v_2); F)$ which implies by Proposition 3.11 that $\rho_1 \oplus \rho_2$ is not taut.

3.3 A reduction principle

We now discuss a reduction principle which will be used later to prove that certain representations are or are not taut. Let $\rho : G \rightarrow \mathbf{O}(V)$ be an orthogonal linear action of a compact Lie group G which is not assumed to be connected. Denote by H a fixed principal isotropy subgroup of the G -action on V and let V^H be the subspace of V that is left point-wise fixed by the action of H . Let N be the normalizer of H in G . Then the group N/H acts on V^H with trivial principal isotropy subgroup. Moreover, the following result is known ([21, 28, 29, 37, 38, 39]):

3.14 Theorem (Luna-Richardson) *The inclusion $V^H \rightarrow V$ induces a stratification pre-serving homeomorphism between orbit spaces*

$$(3.15) \quad V^H/N \rightarrow V/G.$$

The injectivity of the map (3.15) means that $Np = Gp \cap V^H$ for $p \in V^H$. In particular, the H -fixed point set of a G -orbit is a smooth manifold. Observe also that for a regular point $p \in V^H$ the normal space to the principal orbit $M = Gp$ at p is contained in V^H , because the slice representation at p is trivial.

3.16 Lemma *Let $p \in V^H$ be a regular point and consider a normal vector ξ to $M = Gp$ at p . Then the Weingarten operator of M at ξ restricts to the Weingarten operator of $M^H = M \cap V^H$ at ξ , in symbols, $A_\xi^M|_{T_p M^H} = A_\xi^{M^H}$.*

Proof. Let $v \in T_p M^H$. Consider an extension $\tilde{\xi}(t)$ of ξ to a normal vector field along a curve $\gamma(t)$ in M^H with $\dot{\gamma}(0) = v$. Then $A_\xi^M v = -\tilde{\xi}'(0)^T$, where “ T ” denotes the orthogonal projection onto $T_p M$. Now $\tilde{\xi}(t) \in V^H$, so the derivative $\tilde{\xi}'(0) \in V^H$. The normal component of $\tilde{\xi}'(0)$ is already in V^H , so its component in $T_p M$ is also in V^H . Hence $A_\xi^{M^H} v = -\tilde{\xi}'(0)^T$ which proves the claim. $\square$

3.17 Lemma *Let $p, q \in V^H$ and suppose that q is a regular point for G . Consider $M = Gp$ and let $L_q : M \rightarrow \mathbf{R}$ be the distance function. Then the critical set of L_q is contained in $M^H = M \cap V^H$, namely, $\text{Crit}(L_q) = \text{Crit}(L_q|_{M^H})$.*

Proof. Let $x \in M$ be a critical point of $L_q : M \rightarrow \mathbf{R}$. Then $x = gp$ for some $g \in G$ and $v = q - x$ is a normal vector to M at x . Consider first the case where p is a regular point for G . Then the isotropy subgroup at x is principal, so $G_x = gHg^{-1}$ acts trivially on the normal space. In particular $gHg^{-1}v = v$, which implies that $gHg^{-1}q = q$. Now $gHg^{-1} \subset G_q = H$ and therefore $g \in N$. Thus $x = gp \in V^H$ and x is a critical point for the restriction $L_q : M^H \rightarrow \mathbf{R}$. Consider now the case where p is not a regular point for G . Then there is $\epsilon > 0$ sufficiently small such that $y = x + \epsilon v$ is a regular point for G . Now y is a critical point for $L_q : Gy \rightarrow \mathbf{R}$, since a line segment meeting one orbit orthogonally is orthogonal to all orbits it meets. By the previous case $y \in V^H$. Hence $x \in V^H$ and again x is a critical point for the restriction $L_q : M^H \rightarrow \mathbf{R}$.

On the other hand, let $x \in M^H$ be a critical point of $L_q : M^H \rightarrow \mathbf{R}$. Consider first the case where p is a regular point for G . Then x is also a regular point for G so that the normal spaces $N_x(M^H)$ and $N_x(M)$ coincide. It follows that x is a critical point of

$L_q : M \rightarrow \mathbf{R}$. Consider now the case where p is not a regular point for G . Then there is $\epsilon > 0$ sufficiently small such that $y = x + \epsilon v \in V^H$ is a regular point for G , where $v = q - x$. Since $Gy \cap V^H = Ny$, we get that y is a critical point for $L_q : Ny \rightarrow \mathbf{R}$. By the previous case y is a critical point for $L_q : Gy \rightarrow \mathbf{R}$. Hence again x is a critical point of $L_q : M \rightarrow \mathbf{R}$. $\square$

For the next lemma and proposition we found inspiration in [14].

3.18 Lemma *Suppose there is a subgroup $L \subset H$ which is a finitely iterated $\mathbf{Z}_2$ -extension of the identity and such that the fixed point sets $V^L = V^H$. Let $M = Gp$ be a principal orbit. Then the sum of the Betti numbers (with respect to $\mathbf{Z}_2$ coefficients) of M^H is less than or equal the sum of the Betti numbers of M , in symbols $\beta(M^H; \mathbf{Z}_2) \leq \beta(M; \mathbf{Z}_2)$.*

Proof. We can write $L = \{h_1, \dots, h_n\} \subset H$ such that h_{i+1} normalizes H_i and $h_{i+1}^2 \in H_i$ for $i : 0, \dots, n-1$, where H_0 is the identity group and H_i is the subgroup of H generated by $h_1, \dots, h_i$. Now each transformation h_{i+1} has order 2 in the fixed point set $M^{H_i} = M \cap V^{H_i}$, so by a theorem of Floyd ([17])

$$\beta(M^L; \mathbf{Z}_2) \leq \beta(M^{H_{n-1}}; \mathbf{Z}_2) \leq \dots \leq \beta(M^{H_0}; \mathbf{Z}_2).$$

But $M^L = M^H$ and $M^{H_0} = M$ and this completes the proof. $\square$

3.19 Proposition *Suppose there is a subgroup $L \subset H$ which is a finitely iterated $\mathbf{Z}_2$ -extension of the identity and such that the fixed point sets $V^L = V^H$. Suppose also that the reduced representation $\bar{\rho} : (N/H)^0 \rightarrow \mathbf{O}(V^H)$ is $\mathbf{Z}_2$ -taut where $(N/H)^0$ denotes the connected component of the identity of N/H . Then $\rho : G \rightarrow \mathbf{O}(V)$ is $\mathbf{Z}_2$ -taut.*

Proof. Let $M = Gp$ be an arbitrary orbit. We shall prove that M is $\mathbf{Z}_2$ -taut. Since V^H intersects all orbits, we may assume $p \in V^H$. Suppose $L_q : M \rightarrow \mathbf{R}$ is a Morse distance function. We must show that L_q is $\mathbf{Z}_2$ -perfect, and in fact it is enough to do that for q in a dense subset of V , so we may take q to be a regular point. Since G acts on V by isometries, we may moreover assume that $q \in V^H$. If $\bar{\rho}$ is $\mathbf{Z}_2$ -taut, then each connected component of M^H is $\mathbf{Z}_2$ -taut, hence M^H itself is $\mathbf{Z}_2$ -taut. Note that $L_q|_{M^H}$ is a Morse function. Hence $L_q|_{M^H}$ is $\mathbf{Z}_2$ -perfect. Now Lemmas 3.17 and 3.18 and the Morse inequalities imply that

$$\beta(M^H; \mathbf{Z}_2) \leq \beta(M; \mathbf{Z}_2) \leq \#\text{Crit}(L_q) = \#\text{Crit}(L_q|_{M^H}),$$

where $\#A$ denotes the cardinality of the finite set A . It follows that L_q is $\mathbf{Z}_2$ -perfect. $\square$

4 A necessary condition for a representation to be of class $\mathcal{O}^2$

Let π be a complex representation of the compact connected Lie group G on a finite-dimensional vector space. We say that π is of *real type* if it comes from a representation of G on a real vector space by extension of scalars, and we say that π is of *quaternionic type*

if it comes from a representation of G on a quaternionic vector space by restriction of scalars. If π is neither of real type nor of quaternionic type, we say that π is of *complex type*².

Now it is known that the finite-dimensional real irreducible representations ρ of G fall into one of the following disjoint classes:

- (a) the complexification ρ^c is irreducible and $\rho^c = \pi$ is a complex representation of real type;
- (b) the complexification ρ^c is reducible and $\rho^c = \pi \oplus \pi$ where π is a complex irreducible representation of quaternionic type;
- (c) the complexification ρ^c is reducible and $\rho^c = \pi \oplus \pi^*$ where π is a complex irreducible representation of complex type and π^* is not equivalent to π (where π^* denotes the dual representation of π).

The relation between ρ and π is that ρ is a real form of π in the first case ($\rho^c = \pi$), but ρ is π viewed as a real representation in the other two cases ($\rho = \pi^r$). We shall call ρ of *real*, *quaternionic* or *complex type* according to whether the associated π is of real, quaternionic or complex type. Note also that π is self-dual precisely in the first two cases.

Suppose now that G is semisimple, let $\mathfrak{g}$ denote its Lie algebra and $\mathfrak{g}^c$ its complexification. Write Δ for the root system of $\mathfrak{g}^c$ with respect to a chosen Cartan subalgebra, Δ^+ for the positive root system with respect to an ordering of the roots, and $\mathcal{S} = \{\alpha_1, \dots, \alpha_r\}$ for the corresponding simple root system. Let $\lambda_1, \dots, \lambda_r$ be the fundamental highest weights defined by the relations $2(\lambda_i, \alpha_j)/(\alpha_j, \alpha_j) = \delta_{ij}$ where $(\cdot, \cdot)$ is the Cartan-Killing form. The Theorem of the Highest Weight of É. Cartan states that the complex irreducible representations of $\mathfrak{g}^c$ are parametrized by their highest weights, and these are exactly the linear combinations $\lambda = \sum_{i=1}^r m_i \lambda_i$ for $m_i \in \{0, 1, 2, \dots\}$. The following proposition is a useful criterium of Dadok to decide for which λ the corresponding representation π_λ is of real, quaternionic or complex type. Recall that the roots $\alpha, \beta \in \Delta$ are called *strongly orthogonal* if $\alpha \pm \beta$ is not a root.

4.1 Proposition ([12]) *There is a maximal subset $\mathcal{B} = \{\beta_1, \dots, \beta_s\} \subset \Delta^+$ of strongly orthogonal roots such that:*

- (a) *we have $s_0 = s_{\beta_1} \cdots s_{\beta_s}$ is the Weyl group element that maps the positive Weyl chamber into its negative;*
- (b) *the representation π_λ is of complex type if and only if λ does not belong to the real span of $\mathcal{B}$;*
- (c) *the representation π_λ is of real type (resp. quaternionic type) if and only if λ belongs to the real span of $\mathcal{B}$ and*

$$k(\lambda) = \sum_{i=1}^s \frac{(\lambda, \beta_i) - (s_0 \lambda, \beta_i)}{(\beta_i, \beta_i)}$$

²It is useful to remark that π is of real type (resp., quaternionic type) if and only if there exists an invariant real (resp., quaternionic) structure on the representation space, i. e. a $\mathbf{C}$ -conjugate linear, G -invariant endomorphism ϵ of the representation space such that $\epsilon^2 = 1$ (resp., $\epsilon^2 = -1$).

is an even (resp. odd) integer.

4.2 Remark For a simple Lie algebra the set $\mathcal{B}$ can be constructed as follows, as is explained in [12] and will be assumed throughout our paper (cf. Appendix A). Let β_1 be the highest root. The root system $\{\alpha \in \Delta : (\alpha, \beta_1) = 0\}$ is either irreducible or equals $\{\pm\zeta_1\} \cup \Delta_1$, with Δ_1 being irreducible and $\zeta_1 \in \Delta^+$. In the former case set β_2 equal to the highest root of Δ_1 (with the inherited order from Δ), and proceed by induction. In the latter case set $\beta_2 = \zeta_1$ and β_3 equal to the highest root of Δ_1 , and proceed by induction.

We continue to assume that G is semisimple. Let H_α , $\alpha \in \Delta$, be the coroots of $\mathfrak{g}^c$. It is possible to choose root vectors X_α for $\mathfrak{g}^c$, $\alpha \in \Delta$, such that the compact real form $\mathfrak{g}$ is spanned by

$$(4.3) \quad iH_\alpha, \quad X_\alpha - X_{-\alpha}, \quad i(X_\alpha + X_{-\alpha}), \quad \text{where } \alpha \in \Delta^+.$$

Now if π is a complex representation of G , we have that the adjoint map $\pi(X_\alpha)^* = \pi(X_{-\alpha})$ with respect to any G -invariant Hermitian product on the representation space. Moreover, if π is of real (resp. quaternionic) type and ϵ is an invariant real (resp. quaternionic) structure on the representation space, then $\pi(X_\alpha)\epsilon = \epsilon\pi(X_{-\alpha})$.

By compactness of G , any real representation is equivalent to an orthogonal one. The following proposition, which is stated as a remark in [12], p. 128, and the ensuing lemma, which is a refinement of a result in [11], are precisely the ingredients we need to establish a necessary condition for an orthogonal representation of G to be of class $\mathcal{O}^2$.

4.4 Proposition ([12]) *Let $\mathcal{U}^k(\mathfrak{g}^c)$ be the k th level in the natural filtration of the universal enveloping algebra of $\mathfrak{g}^c$. Fix π_λ , the irreducible representation of $\mathfrak{g}^c$ with highest weight λ , and fix v_λ , a highest weight vector. Let*

$$n_i = \frac{(\lambda, \beta_i) - (s_0\lambda, \beta_i)}{(\beta_i, \beta_i)},$$

for $i = 1, \dots, s$. Then the element s_0v_λ is in $\mathcal{U}^{k(\lambda)}(\mathfrak{g}^c)v_\lambda$, but not in $\mathcal{U}^l(\mathfrak{g}^c)v_\lambda$ for any $l < k(\lambda)$, where s_0v_λ is given as

$$s_0v_\lambda = \pi(X_{-\beta_1})^{n_1} \cdots \pi(X_{-\beta_s})^{n_s} v_\lambda.$$

There is no proof of the above proposition in [12]. In [10], p. 270, there is an attempt to prove the proposition which in our view contains serious gaps.

Proof. It is enough to consider the case where $\mathfrak{g}^c$ is a complex simple Lie algebra. We first claim that the element s_0v_λ is not zero. In fact, consider the complex subalgebra $\mathfrak{k} \subset \mathfrak{g}^c$ generated by the root spaces of $\mathfrak{g}^c$ corresponding to $\pm\beta_1, \dots, \pm\beta_s$. Then $\mathfrak{k}$ is isomorphic to the direct product of s copies of $\mathfrak{sl}(2, \mathbf{C})$. We restrict π_λ to $\mathfrak{k}$ and let U_λ be the unique irreducible $\mathfrak{k}$ -module generated by v_λ . Since $\lambda - s_0\lambda = \sum_{i=1}^s n_i \beta_i$, it is now clear that U_λ is the representation space of $\mathfrak{k} : \underset{n_1}{\circ} \otimes \cdots \otimes \underset{n_s}{\circ}$ and that $s_0v_\lambda \in U_\lambda$. Therefore, our claim is reduced to the case of the $(n+1)$ -dimensional complex irreducible representation of $\mathfrak{sl}(2, \mathbf{C})$, namely $\circ$, which is immediate to verify.

It is obvious that $s_0 v_\lambda \in \mathcal{U}^{k(\lambda)}(\mathfrak{g}^c) v_\lambda$. We next prove that $s_0 v_\lambda \notin \mathcal{U}^l(\mathfrak{g}^c) v_\lambda$ for $l < k(\lambda)$ by contradiction. In fact, enumerate the positive roots $\Delta^+ = \{\alpha_1, \dots, \alpha_t\}$ and suppose $s_0 v_\lambda \in \mathcal{U}^l(\mathfrak{g}^c) v_\lambda$ for $l < k(\lambda)$. It follows from the Poincaré-Birkhoff-Witt theorem that we can write

$$s_0 v_\lambda = \sum_{p_1, \dots, p_t \geq 0} c_{p_1 \dots p_t} X_{-\alpha_1}^{p_1} \cdots X_{-\alpha_t}^{p_t} v_\lambda,$$

for some complex constants $c_{p_1 \dots p_t}$. Now, using that $X_\alpha^* = X_{-\alpha}$,

$$\begin{aligned} 0 &\neq (s_0 v_\lambda, s_0 v_\lambda) \\ &= \sum_{p_1, \dots, p_t \geq 0} c_{p_1 \dots p_t} (X_{-\alpha_1}^{p_1} \cdots X_{-\alpha_t}^{p_t} v_\lambda, X_{-\beta_1}^{n_1} \cdots X_{-\beta_s}^{n_s} v_\lambda) \\ &= \sum_{p_1, \dots, p_t \geq 0} c_{p_1 \dots p_t} (X_{\beta_s}^{n_s} \cdots X_{\beta_1}^{n_1} X_{-\alpha_1}^{p_1} \cdots X_{-\alpha_t}^{p_t} v_\lambda, v_\lambda). \end{aligned}$$

Recall that the highest weight space is one-dimensional; hence we can write

$$X_{\beta_s}^{n_s} \cdots X_{\beta_1}^{n_1} X_{-\alpha_1}^{p_1} \cdots X_{-\alpha_t}^{p_t} v_\lambda = c v_\lambda,$$

for some nonnegative integers $p_1, \dots, p_t$ such that $k(\lambda) = n_1 + \cdots + n_s > p_1 + \cdots + p_t = l$ and for some nonzero complex constant c . Note that $n_1 \beta_1 + \cdots + n_s \beta_s = p_1 \alpha_1 + \cdots + p_t \alpha_t$. The contradiction we are aiming at now follows from the following claim.

CLAIM 1 *If $N > M$ then*

$$X_{\delta_N} \cdots X_{\delta_1} X_{-\gamma_1} \cdots X_{-\gamma_M} v_\lambda = 0,$$

where $\delta_1, \dots, \delta_N \in \mathcal{B}$, and $\gamma_1, \dots, \gamma_M \in \Delta^+$.

We proceed by induction on the integer N . The cases $N = 1$ and $N = 2$ are trivial (use that v_λ is a highest weight vector for $N = 1$ and that the sum of two roots in $\mathcal{B}$ is not a root for $N = 2$). Assume the assertion is true for some $N - 1 \geq 2$ and let us prove it for N . Since $X_{\beta_1}, \dots, X_{\beta_s}$ pairwise commute, we may assume that $\delta_1 = \beta_i$ where $i \leq j$ for every $j = 1, \dots, s$ which satisfies $\beta_j = \delta_k$ for some $k = 1, \dots, N$.

CLAIM 2 *For each $j = 1, \dots, M$, if $\delta_1 - \gamma_j$ is a root then it is a positive root.*

Indeed we have $\delta_1 = \beta_i$ where i is as above. If $i = 1$, then we are done because β_1 is the highest root. If not, $0 = (\beta_1, \delta_1 + \cdots + \delta_N) = (\beta_1, \gamma_1 + \cdots + \gamma_M)$. Now $(\beta_1, \gamma_1) = -(\beta_1, \gamma_2 + \cdots + \gamma_M)$, and since β_1 is the highest root we have $(\beta_1, \gamma_1) \geq 0, \dots, (\beta_1, \gamma_M) \geq 0$. It follows that $(\beta_1, \gamma_1) = 0$ and $(\beta_1, \gamma_2 + \cdots + \gamma_M) = 0$. An easy induction argument shows that $(\beta_1, \gamma_j) = 0$ for $j = 1, \dots, M$. Consider now the root system $\{\alpha \in \Delta : (\alpha, \beta_1) = 0\}$. The first case occurs when it is irreducible. Then β_2 is by definition its highest root so that $\beta_2 \geq \gamma_j$ for $j = 1, \dots, M$. If $i = 2$ we are done. If not, we see that $(\beta_2, \gamma_j) = 0$ for $j = 1, \dots, M$ and we proceed by induction. The second case occurs when the above root system is of the form $\{\pm \zeta_1\} \cup \Delta_1$, with Δ_1 being irreducible and $\zeta_1 \in \Delta^+$. Here $\beta_2 = \zeta_1$ and β_3 is the highest root of Δ_1 . If $\gamma_j \in \Delta_1$ for $j = 1, \dots, M$ then $(\beta_2, \gamma_j) = 0$ for $j = 1, \dots, M$ so that $i \geq 3$ and we proceed by induction as above. On the other hand, if $\zeta_1 = \gamma_{j_0}$ for some j_0 , then $i = 2$. In this case, for each $j = 1, \dots, M$ either $\beta_2 = \gamma_j$ or $-\gamma_j \in \Delta_1$. So $\beta_2 - \gamma_j$ is never a root. This completes the proof of Claim 2.

CLAIM 3 *We have*

$$X_{\delta_N} \cdots X_{\delta_2} X_{-\gamma_1} \cdots X_{-\gamma_{j-1}} H_\zeta X_{-\gamma_{j+1}} \cdots X_{-\gamma_M} v_\lambda = 0,$$

where H_ζ is the coroot vector corresponding to $\zeta \in \Delta^+$ and $j = 1, \dots, M$.

In order to prove Claim 3, proceed by induction on j . The initial case is $j = M$, which follows from the induction hypothesis on N , since $H_\zeta v_\lambda = (\zeta, \lambda) v_\lambda$. Next write

$$(4.5) \quad \begin{aligned} & X_{\delta_N} \cdots X_{\delta_2} X_{-\gamma_1} \cdots X_{-\gamma_{j-1}} H_\zeta X_{-\gamma_{j+1}} \cdots X_{-\gamma_M} v_\lambda = \\ & X_{\delta_N} \cdots X_{\delta_2} X_{-\gamma_1} \cdots X_{-\gamma_{j-1}} X_{-\gamma_{j+1}} H_\zeta \cdots X_{-\gamma_M} v_\lambda \\ & + X_{\delta_N} \cdots X_{\delta_2} X_{-\gamma_1} \cdots X_{-\gamma_{j-1}} [H_\zeta, X_{-\gamma_{j+1}}] \cdots X_{-\gamma_M} v_\lambda. \end{aligned}$$

The first summand on the right hand side of (4.5) is zero by the induction hypothesis on j , and the second summand is zero because $[H_\zeta, X_{-\gamma_{j+1}}] = -(\zeta, \gamma_{j+1}) X_{-\gamma_{j+1}}$ so that we can use the induction hypothesis on N . This proves Claim 3.

CLAIM 4 *We have*

$$X_{\delta_N} \cdots X_{\delta_2} X_{-\gamma_1} \cdots X_{-\gamma_{j-1}} X_\zeta X_{-\gamma_{j+1}} \cdots X_{-\gamma_M} v_\lambda = 0,$$

where X_ζ is the root vector corresponding to $\zeta \in \Delta^+$ and $j = 1, \dots, M$.

In order to prove Claim 4, proceed by induction on j . The initial case is $j = M$, which is trivial because v_λ is a highest weight vector. Next write

$$(4.6) \quad \begin{aligned} & X_{\delta_N} \cdots X_{\delta_2} X_{-\gamma_1} \cdots X_{-\gamma_{j-1}} X_\zeta X_{-\gamma_{j+1}} \cdots X_{-\gamma_M} v_\lambda = \\ & X_{\delta_N} \cdots X_{\delta_2} X_{-\gamma_1} \cdots X_{-\gamma_{j-1}} X_{-\gamma_{j+1}} X_\zeta \cdots X_{-\gamma_M} v_\lambda \\ & + X_{\delta_N} \cdots X_{\delta_2} X_{-\gamma_1} \cdots X_{-\gamma_{j-1}} [X_\zeta, X_{-\gamma_{j+1}}] \cdots X_{-\gamma_M} v_\lambda. \end{aligned}$$

The first summand on the right hand side of (4.6) is zero by the induction hypothesis on j . The second summand is also zero for the following reasons. If $\zeta = \gamma_{j+1}$ we use Claim 3. If not, then either $\zeta - \gamma_{j+1}$ is not a root and then $[X_\zeta, X_{-\gamma_{j+1}}] = 0$, or else $\zeta - \gamma_{j+1}$ is a root. In the latter case, if it is a positive root we can use the induction hypothesis on j , and if it is a negative root then we can use the induction hypothesis on N . This completes the proof of Claim 4. We finally turn to the proof of Claim 1. We can write

$$(4.7) \quad \begin{aligned} & X_{\delta_N} \cdots X_{\delta_1} X_{-\gamma_1} \cdots X_{-\gamma_M} v_\lambda = \\ & \sum_{j=1}^M X_{\delta_N} \cdots X_{\delta_2} X_{-\gamma_1} \cdots X_{-\gamma_{j-1}} [X_{\delta_1}, X_{-\gamma_j}] X_{-\gamma_{j+1}} \cdots X_{-\gamma_M} v_\lambda \\ & + X_{\delta_N} \cdots X_{\delta_2} X_{-\gamma_1} \cdots X_{-\gamma_M} X_{\delta_1} v_\lambda. \end{aligned}$$

The second summand in the right hand side of (4.7) is zero because v_λ is a highest weight vector. The first summand is also zero because if $\delta_1 = \gamma_j$ we can apply Claim 3. Otherwise, either $\delta_1 - \gamma_j$ is not a root and then $[X_{\delta_1}, X_{-\gamma_j}] = 0$ or else $\delta_1 - \gamma_j$ is a root and then it is a positive root by Claim 2 so that we can apply Claim 4. This completes the proof of Claim 1 and the proof of the proposition. $\square$

4.8 Lemma *Let π_λ be the complex irreducible representation of G with highest weight λ and representation space V_λ . Let μ be a weight of π_λ and fix a weight vector v_μ . Let ρ denote a real form of π_λ in case π_λ is of real type, or the realification of π_λ in case it is either of quaternionic or of complex type. Suppose that ρ is of class $\mathcal{O}^2$.*

(a) *If π_λ is of real type, then*

$$\mathcal{U}^2(\mathfrak{g}^c)v_\mu + \mathcal{U}^2(\mathfrak{g}^c)\epsilon(v_\mu) = V_\lambda,$$

where ϵ is the real structure on V_λ defined by ρ . In particular, by taking $\mu = \lambda$, we find that

$$\mathcal{U}^2(\mathfrak{g}^c)v_\lambda + \mathcal{U}^2(\mathfrak{g}^c)\epsilon(v_\lambda) = V_\lambda. \quad (C_2)$$

(b) *If π_λ is of complex or quaternionic type, then $\mathcal{U}^2(\mathfrak{g}^c)v_\mu = V_\lambda$.*

Proof. (a) Consider the real vector $p = v_\mu + \epsilon v_\mu$. The second osculating space $\mathcal{O}_p^2(Gp)$ is spanned over $\mathbf{R}$ by Xp , XYp , where $X, Y \in \mathfrak{g}$. Taking linear combinations with complex coefficients of these vectors, we can write that the complexification

$$\mathcal{O}_p^2(Gp)^c \subset \mathcal{U}^2(\mathfrak{g}^c)v_\mu + \mathcal{U}^2(\mathfrak{g}^c)\epsilon(v_\mu) \subset V_\lambda.$$

But ρ of class $(\mathcal{O}^2)$ implies that $\mathcal{O}_p^2(Gp)^c = V_\lambda$.

(b) Let $p = v_\mu$. We have that $\mathcal{O}_p^2(Gp) \subset \mathcal{U}^2(\mathfrak{g}^c)v_\mu \subset V_\lambda$ as real vector spaces and the proof is similar as in (a). $\square$

We now state the main result of this section. Notice that the same result is claimed in [10], p. 271, with an attempt of a proof which in our opinion is not satisfactory.

4.9 Proposition *Let ρ be a real (orthogonal) irreducible representation of a compact connected semisimple Lie group G with complexified Lie algebra $\mathfrak{g}^c$, and let π_λ be the associated complex irreducible representation. Suppose that ρ is of class $\mathcal{O}^2$.*

(a) *If ρ is of quaternionic type, then $k(\lambda) = 1$.*

(b) *If ρ is of complex type, then $k(\lambda) = 1, 2$.*

(c) *If ρ is of real type, then $k(\lambda) = 2, 4$.*

Proof. First consider ρ to be of quaternionic or complex type, i. e. $\rho^c = \pi_\lambda \oplus \pi_\lambda^*$. Then ρ of class $\mathcal{O}^2$ forces $\mathcal{U}^2(\mathfrak{g}^c)v_\lambda = V_\lambda$ (Lemma 4.8) and then $s_0 v_\lambda \in \mathcal{U}^2(\mathfrak{g}^c)v_\lambda$ implies that $k(\lambda) \leq 2$ (Proposition 4.4). From this follow (a) and (b).

Now take ρ to be of real type. Let v_λ be a highest weight vector of $\rho^c = \pi_\lambda$. Since π_λ is self-dual,

$$\text{lowest weight of } \pi_\lambda = -\text{highest weight of } \pi_\lambda^* = -\lambda = s_0 \lambda.$$

Also, there exists a $\mathbf{C}$ -conjugate linear, G -invariant involution ϵ of V_λ and $\epsilon(v_\lambda)$ is a lowest weight vector. We shall assume $k(\lambda) \geq 6$ and derive a contradiction. Since $k(\lambda)$ is even, it is possible to write

$$\frac{k(\lambda)}{2} = n_1 + \dots + n_{i_0} + m$$

for some integers $0 \leq i_0 < s$, $0 \leq m < n_{i_0+1}$, where n_i is as in Proposition 4.4. Set

$$\mu = \lambda - n_1\beta_1 - \dots - n_{i_0}\beta_{i_0} - m\beta_{i_0+1}.$$

Note that μ is a weight: a μ -weight vector is

$$u = X_{-\beta_1}^{n_1} \dots X_{-\beta_{i_0}}^{n_{i_0}} X_{-\beta_{i_0+1}}^m v_\lambda.$$

Use Lemma 4.8 to decompose $u = u_1 + u_2$ where $u_1 \in \mathcal{U}^2(\mathfrak{g}^c)v_\lambda$ and $u_2 \in \mathcal{U}^2(\mathfrak{g}^c)\epsilon(v_\lambda)$. First, assume that $u_2 \neq 0$. Since $\frac{k(\lambda)}{2} \geq 3$, we have

$$(4.10) \quad u_2 = u - u_1 \in \mathcal{U}^{\frac{k(\lambda)}{2}}(\mathfrak{g}^c)v_\lambda + \mathcal{U}^2(\mathfrak{g}^c)v_\lambda \subset \mathcal{U}^{\frac{k(\lambda)}{2}}(\mathfrak{g}^c)v_\lambda.$$

Moreover, it is clear that u_2 may be assumed to be a μ -weight vector (because a component of u_2 in a different weight space has to cancel with the corresponding component of u_1 in the same weight space) and that

$$u_2 = \sum_{\gamma, \delta \in \Delta^+} c_{\gamma, \delta} X_{-\gamma} X_{-\delta} \epsilon(v_\lambda)$$

for some complex constants $c_{\gamma, \delta}$ (we do not need to consider terms of first order in the sum because $\lambda - \mu$ cannot be a root). Therefore,

$$0 \neq (u_2, u_2) = (u_2, \sum_{\gamma, \delta \in \Delta^+} c_{\gamma, \delta} X_{-\gamma} X_{-\delta} \epsilon(v_\lambda)) = (\sum_{\gamma, \delta \in \Delta^+} \bar{c}_{\gamma, \delta} X_\gamma X_\delta u_2, \epsilon(v_\lambda)).$$

This shows that

$$\sum_{\gamma, \delta \in \Delta^+} \bar{c}_{\gamma, \delta} X_\gamma X_\delta u_2$$

is a nonzero multiple of the lowest weight vector $\epsilon(v_\lambda)$. This, combined with (4.10), gives that $\epsilon(v_\lambda)$ is in $\mathcal{U}^{\frac{k(\lambda)}{2}+2}(\mathfrak{g}^c)v_\lambda$. But $\frac{k(\lambda)}{2} + 2 < k(\lambda)$, contradicting Proposition 4.4.

In case $u_2 = 0$ we have that $u = u_1 \in \mathcal{U}^2(\mathfrak{g}^c)v_\lambda$. Since

$$X_{-\beta_{i_0+1}}^{n_{i_0+1}-m} X_{-\beta_{i_0+2}}^{n_{i_0+2}} \dots X_{-\beta_s}^{n_s} u = s_0 v_\lambda$$

is a nonzero multiple of the lowest weight vector $\epsilon(v_\lambda)$ and $n_{i_0+1} - m + n_{i_0+2} + \dots + n_s = \frac{k(\lambda)}{2}$, again $\epsilon(v_\lambda) \in \mathcal{U}^{\frac{k(\lambda)}{2}+2}(\mathfrak{g}^c)v_\lambda$, contradicting the same proposition. $\square$

5 The candidates to a position in class $\mathcal{O}^2$

In this section we want to elaborate a list of possibilities for real irreducible representations of class $\mathcal{O}^2$ which we shall use later in Sections 6 and 7 to classify variationally complete and taut representations. Our point of view, as is usually the case in other papers on the subject, is to classify orthogonal representations up to the following equivalence relation: we call two

representations $\rho : G \rightarrow \mathbf{O}(V)$, $\rho' : G' \rightarrow \mathbf{O}(V')$ *image equivalent* if there exists an isometry $\psi : V \rightarrow V'$ such that $\mathbf{O}(\psi)(\rho(G)) = \rho'(G')$, where $\mathbf{O}(\psi) : \mathbf{O}(V) \rightarrow \mathbf{O}(V')$ is the induced conjugation map³. In particular, the image equivalence class of ρ always contains its dual ρ^* (in fact any representation obtained from ρ by an automorphism of the Dynkin diagram), as well as the pull back representation $\tilde{\rho} = \rho \circ p$ of a covering group $p : \tilde{G} \rightarrow G$.

Let G be a compact connected Lie group. We have that G is finitely covered by $T^n \times G_s$, where T^n is an n -dimensional torus and G_s is a compact connected semisimple Lie group (which may also be assumed to be simply-connected, whenever convenient), and any representation of G pulls back to a representation of $T^n \times G_s$. In view of image equivalence, in order to study representations of G , we may assume that $G = T^n \times G_s$ and restrict to almost faithful representations. Next we observe that any complex irreducible representation of T^n is a character (hence of complex type). Therefore a complex irreducible representation of G can be of real type or of quaternionic type only if it is trivial on T^n , and in case it is of complex type, then it has an $(n - 1)$ -dimensional kernel on T^n . Now we have come to our working hypothesis (for a circle group S^1 , let x^n denote the n th power representation of that circle which is a complex representation of complex type):

Let ρ be a real irreducible representation of a compact connected Lie group G on a finite-dimensional real vector space V .

- If ρ is of real type, we assume that $G = G_s$ is a compact connected semisimple Lie group, and ρ is a real form of a complex irreducible representation $\pi = \pi_\lambda$ of G of real type on the complex vector space V_λ for some highest weight λ .
- If ρ is of quaternionic type, we assume that $G = G_s$, where G_s is as above, and ρ is the realification of a complex irreducible representation $\pi = \pi_\lambda$ of G of quaternionic type on the complex vector space V_λ for some highest weight λ .
- If ρ is of complex type, we assume that $G = G_s$ or $G = S^1 \times G_s$, where G_s is as above, and ρ is the realification of a complex irreducible representation π of G of complex type. According to the form of G , we write $\pi = \pi_\lambda$ or $\pi = x \otimes \pi_\lambda$ as a representation on the complex vector space $V_\lambda = \mathbf{C} \otimes_{\mathbf{C}} V_\lambda$ for some highest weight λ .


Isotropy representations of symmetric spaces and cohomogeneity one representations of compact connected Lie groups are of course two classes of representations that belong to class $\mathcal{O}^2$. In order to keep the exposition as simple as possible, we assume knowledge of the classification of symmetric spaces. So in the course of this section we will be disregarding all the isotropy representations of symmetric spaces that we encounter, according to tables 8.11.2 and 8.11.5 in [45]. Moreover, we classify in Section 8 the cohomogeneity one representations of compact connected Lie groups. (As a side remark, note that our proof of Theorem 8.2 is completely independent from the classification of the symmetric spaces.) Therefore, in addition to our previous hypotheses we assume throughout this section:

³This is finer than the notion of orbit equivalence which is explained in Section 2, but of course not as fine as the usual notion of equivalence for representations.

The real irreducible representation ρ is of class $\mathcal{O}^2$, cohomogeneity greater than one and it is not the isotropy representation of a symmetric space.

The results of our investigation in this section (namely, Propositions 5.1, 5.2, 5.17, 5.18, 5.19, 5.20 and 5.21) finally imply: the possibilities for ρ are those collected in the table of Proposition 6.3 and in Tables C.1, C.2, C.3 and C.4 in Appendix C.

We need some amount of notation (cf. [45], p. 237). Let $\mathbf{A}_n = \mathbf{SU}(n+1)$, $\mathbf{B}_n = \mathbf{Spin}(2n+1)$, $\mathbf{C}_n = \mathbf{Sp}(n)$, $\mathbf{D}_n = \mathbf{Spin}(2n)$, $\mathbf{G}_2$, $\mathbf{F}_4$, $\mathbf{E}_6$, $\mathbf{E}_7$ and $\mathbf{E}_8$ refer to É. Cartan's classification types of simple Lie groups and Lie algebras. For the complex semisimple Lie algebra $\mathfrak{g}^c$, we use the following notation for its complex irreducible representation π_λ : if the integer $2(\lambda, \alpha_i)/(\alpha_i, \alpha_i) \neq 0$, then we write it next to the vertex of the Dynkin diagram of $\mathfrak{g}^c$ which corresponds to α_i (recall that $\mathcal{S} = \{\alpha_1, \dots, \alpha_r\}$ is the simple root system). For example,



denote respectively the vector representations of $\mathbf{A}_n$, $\mathbf{B}_n$, $\mathbf{C}_n$, $\mathbf{D}_n$, and the adjoint representations are given by:

$\mathbf{A}_1$		$\mathbf{G}_2$	
$\mathbf{A}_n, n \geq 2$		$\mathbf{F}_4$	
$\mathbf{B}_n, n \geq 3$		$\mathbf{E}_6$	
$\mathbf{C}_n, n \geq 2$		$\mathbf{E}_7$	
$\mathbf{D}_n, n \geq 4$		$\mathbf{E}_8$	

Note that we are using the dot convention: if there are two lengths of roots, then the short roots are black in the Dynkin diagram. And a real irreducible representation is denoted by the diagram of its complexification.

We add that we shall refer repeatedly to Table B.1 in Appendix B, which contains the values of the invariant $k(\lambda)$ for the fundamental representations of the complex simple Lie algebras.

5.1 The case where ρ is of quaternionic type

Here there is nothing:

5.1 Proposition *The representation ρ cannot be of quaternionic type.*

Proof. If ρ is of class $\mathcal{O}^2$ and quaternionic type, Proposition 4.9 says that $k(\lambda) = 1$. A glance at Table B.1 now shows that $G = \mathbf{Sp}(n)$ and π_λ is the vector representation. But then ρ is of cohomogeneity one. $\square$

5.2 The case where ρ is of complex type

Here the result is:

5.2 Proposition *Let ρ be of complex type. Then ρ is one of the following:*

G	ρ	Conditions
$\mathbf{SU}(n) \times \mathbf{SU}(m)$	$(\overset{1}{\circ} \cdots \cdots \overset{1}{\circ} \otimes \overset{1}{\circ} \cdots \cdots \overset{1}{\circ})$ $\oplus (\overset{1}{\circ} \cdots \cdots \overset{1}{\circ} \otimes \overset{1}{\circ} \cdots \cdots \overset{1}{\circ})$	$n \neq m$
$\mathbf{SU}(n) \times \mathbf{Sp}(m)$	$(\overset{1}{\circ} \cdots \cdots \overset{1}{\circ} \oplus \overset{1}{\circ} \cdots \cdots \overset{1}{\circ})$ $\otimes \bullet \cdots \cdots \bullet \equiv \bullet$	$n \geq 3, m \geq 2$
$S^1 \times \mathbf{SU}(n) \times \mathbf{Sp}(m)$	$(x \otimes \overset{1}{\circ} \cdots \cdots \overset{1}{\circ} \oplus x^{-1} \otimes \overset{1}{\circ} \cdots \cdots \overset{1}{\circ})$ $\otimes \bullet \cdots \cdots \bullet \equiv \bullet$	$m \geq 2$
$\mathbf{SU}(n)$	$\overset{1}{\circ} \cdots \cdots \overset{1}{\circ} \oplus \overset{1}{\circ} \cdots \cdots \overset{1}{\circ}$	n odd
$\mathbf{Spin}(10)$	$\overset{1}{\circ} \cdots \cdots \overset{1}{\circ} \oplus \overset{1}{\circ} \cdots \cdots \overset{1}{\circ}$	—
$\mathbf{SO}(2) \times \mathbf{Spin}(7)$	$(x \oplus x^{-1}) \otimes \overset{1}{\circ} \cdots \cdots \bullet \equiv \bullet$	—
$\mathbf{SO}(2) \times \mathbf{Spin}(9)$	$(x \oplus x^{-1}) \otimes \overset{1}{\circ} \cdots \cdots \bullet \equiv \bullet$	—
$\mathbf{SO}(2) \times \mathbf{G}_2$	$(x \oplus x^{-1}) \otimes \bullet \equiv \bullet$	—

The proof of Proposition 5.2 will be given after the proof of the following two lemmas.

5.3 Lemma *Let π_λ be a complex irreducible representation of a compact connected semisimple Lie group G_s and suppose that the realification of the tensor product representation $\pi = x \otimes \pi_\lambda$ of $G = S^1 \times G_s$ is of class $\mathcal{O}^2$. We have:*

- (a) *if π_λ is of real type and G_s is simple, then $G_s = \mathbf{Spin}(7)$ or $G_s = \mathbf{Spin}(9)$ and π_λ is the respective spin representation, or $G_s = \mathbf{G}_2$ and π_λ is the 7-dimensional representation, or $G_s = \mathbf{SO}(m)$ for $m \neq 2, 4$ and π_λ is the vector representation;*
- (b) *if π_λ is of real type and G_s is not simple, then $G_s = \mathbf{Sp}(1) \times \mathbf{Sp}(m)$ and π_λ is the tensor product of the vector representations of each of the factors;*

(c) if π_λ is of quaternionic type or complex type, then $k(\lambda) = 1, 2$.

Proof. (a) and (b) Let V_λ be the representation space of π_λ and V be the real subspace where a real form acts. Note that the representation space for π is still V_λ , but π is of complex type. Let $p = 1 \otimes a \in \mathbf{C} \otimes_{\mathbf{R}} V = V^c = V_\lambda$, n the dimension of V and q the codimension of $G_s a$ in V . Since the realification of π is of class $\mathcal{O}^2$, we have that $\mathcal{O}_p^2(Gp) = V^{cr} \cong V \oplus iV$. But

$$\mathcal{O}_p^2(Gp) \subset 1 \otimes [\mathbf{R}a + \mathcal{O}_a^2(G_s a)] + i \otimes [\mathbf{R}a + T_a(G_s a)],$$

so by comparing real and imaginary parts we get that $V = \mathbf{R}a + T_a(G_s a)$. Therefore $q = 1$ and G acts with cohomogeneity one on V . Now (a) follows from Lemmas 5.11 and 5.14 below, and (b) follows from Lemmas 5.11 and 5.15 below.

(c) Consider the point $p = v_\lambda \in V_\lambda$. We have that $\mathcal{O}_p^2(Gp) \subset \mathcal{U}^2(\mathfrak{g}^c)v_\lambda \subset V_\lambda$ as real vector spaces. Since the realification of π is of class $\mathcal{O}^2$, we have that $\mathcal{U}^2(\mathfrak{g}^c)v_\lambda = V_\lambda$. But $\mathcal{U}^2(\mathfrak{g}_s^c)v_\lambda = \mathcal{U}^2(\mathfrak{g}^c)v_\lambda$, since the circle subgroup S^1 preserves the weight spaces of π_λ . Therefore $\mathcal{U}^2(\mathfrak{g}_s^c)v_\lambda = V_\lambda$ and we can apply Proposition 4.4 as in the first paragraph of the proof of Proposition 4.9. $\square$

Part (a) of the following lemma also follows from [41].

5.4 Lemma (a) Let π_λ be one of the following complex irreducible representations of complex type of the compact connected semisimple Lie group G_s :

G_s	π_λ	Conditions
$\mathbf{SU}(n) \times \mathbf{SU}(m)$		$n \neq m$
$\mathbf{SU}(n)$		$n = 2p + 1$
$\mathbf{Spin}(10)$		—

Consider the tensor product representation $\pi = x \otimes \pi_\lambda$ of $S^1 \times G_s$. Then the realifications of π_λ and π are orbit equivalent.

(b) The realifications of the following complex irreducible representations π_λ of complex type of the compact connected semisimple Lie group G_s are not of class $\mathcal{O}^2$:

G_s	π_λ	Conditions
$\mathbf{SU}(n)$		$n \geq 3$
$\mathbf{SU}(n)$		$n = 2p \geq 6$
$\mathbf{SU}(n) \times \mathbf{SU}(n)$		$n \geq 3$
$\mathbf{E}_6$		—

Proof. PRELIMINARY CONSIDERATIONS: Each one of these representations π_λ has the property that the realification of $\pi = x \otimes \pi_\lambda$ is the isotropy representation of a compact irreducible Hermitian symmetric space $X = L_0/K_0$, where K_0 is locally isomorphic to $S^1 \times G_s$. We refer to [24], Chapter VIII, §7, for results about Hermitian symmetric spaces that we will be using in the following. Let $\mathfrak{l}_0 = \mathfrak{k}_0 + \mathfrak{p}_0$ be the decomposition of the Lie algebra $\mathfrak{l}_0$ of L_0 into the ± 1 -eigenspaces of the symmetry. Let $\mathfrak{c}_0$ be the Lie algebra of S^1 , and let $\mathfrak{t}_0$ be some Cartan subalgebra of the Lie algebra $\mathfrak{g}_s$ of G_s . Then $\mathfrak{h}_0 = \mathfrak{c}_0 + \mathfrak{t}_0$ is a Cartan subalgebra of $\mathfrak{k}_0$ and of $\mathfrak{l}_0$. Let $\mathfrak{l}$ be the complexification of $\mathfrak{l}_0$ and let $\mathfrak{c}, \mathfrak{t}, \mathfrak{h}, \mathfrak{k}, \mathfrak{p}$ be the complex subspaces of $\mathfrak{l}$ spanned by $\mathfrak{c}_0, \mathfrak{t}_0, \mathfrak{h}_0, \mathfrak{k}_0, \mathfrak{p}_0$. Let Δ denote the root system of $(\mathfrak{l}, \mathfrak{h})$ and consider the root space decomposition $\mathfrak{l} = \mathfrak{h} + \sum_{\alpha \in \Delta} \mathfrak{l}_\alpha$. For each $\alpha \in \Delta$, we have that either $\mathfrak{l}_\alpha \subset \mathfrak{k}$ or $\mathfrak{l}_\alpha \subset \mathfrak{p}$, in which cases the root α is called respectively *compact* or *noncompact*. A root is compact if and only if it vanishes on $\mathfrak{c}$. Let Δ_c and Δ_n denote respectively the subsets of Δ of compact and noncompact roots. Then we have decompositions $\mathfrak{k} = \mathfrak{h} + \sum_{\alpha \in \Delta_c} \mathfrak{l}_\alpha$, $\mathfrak{p} = \sum_{\gamma \in \Delta_n} \mathfrak{l}_\gamma$. Each root is real-valued on $i\mathfrak{h}_0$. We introduce a lexicographic ordering on the dual of $i\mathfrak{h}_0$ that takes $i\mathfrak{c}_0$ before $i\mathfrak{t}_0$. Let Δ^+, Δ_c^+ and Δ_n^+ be the set of positive roots in Δ, Δ_c and Δ_n , and define

$$\mathfrak{p}^+ = \sum_{\gamma \in \Delta_n^+} \mathfrak{l}_\gamma, \quad \mathfrak{p}^- = \sum_{-\gamma \in \Delta_n^+} \mathfrak{l}_\gamma.$$

Then $\mathfrak{p} = \mathfrak{p}^+ + \mathfrak{p}^-$ is an $\text{ad}_{\mathfrak{k}}$ -invariant decomposition. The $\text{ad}_{\mathfrak{k}}$ -invariant complex structure J_0 on $\mathfrak{p}_0$ is given by ad_{H_0} , where $H_0 \in \mathfrak{c}_0$ is determined by $\gamma(H_0) = i$ for $\gamma \in \Delta_n^+$. The $\mathbf{C}$ -linear extension J of J_0 to $\mathfrak{p}$ has $\mathfrak{p}^\pm$ as $\pm i$ -eigenspaces. Let $\Gamma = \{\gamma_1, \dots, \gamma_s\} \subset \Delta_n^+$ be a maximal subset of strongly orthogonal roots. Then $\mathfrak{a}_0 = \sum_{j=1}^s \mathbf{R}(X_{\gamma_j} - X_{-\gamma_j})$ is a maximal Abelian subspace of $\mathfrak{p}_0$. We view π^r as the adjoint action of K_0 on $\mathfrak{p}_0$, and then π^r is polar and $\mathfrak{a}_0$ is a section for π^r . We also view π as the the adjoint action of K_0 on $\mathfrak{p}^+$, and then the weight system of π is Δ_n^+ . Now the root system of $(\mathfrak{g}_s^c, \mathfrak{t})$ is

$$\Delta_0 = \{\alpha|_{\mathfrak{t}} : \alpha \in \Delta_c\} \quad (\text{also } \Delta_0^+ = \{\alpha|_{\mathfrak{t}} : \alpha \in \Delta_c^+\}),$$

and the weight system of π_λ is

$$\Phi = \{\gamma|_{\mathfrak{t}} : \gamma \in \Delta_n^+\}.$$

It is also useful to remark that any $\text{ad}_{\mathfrak{k}}$ -invariant inner product $\langle \cdot, \cdot \rangle$ on $\mathfrak{p}_0$ is given by a negative multiple of the restriction of the Killing form of $\mathfrak{l}_0$; fix one and extend it to a Hermitian product $\ll \cdot, \cdot \gg$ on $\mathfrak{p}$. Then $\ll \cdot, \cdot \gg$ is $\text{ad}_{\mathfrak{k}}$ -invariant and J is a skew-Hermitian operator on $\mathfrak{p}$ with respect to $\ll \cdot, \cdot \gg$.

PROOF OF (a): Since $\mathfrak{a}_0$ is a section for π^r , it is clearly enough to check that given

$$p = \sum_{j=1}^s a_j (X_{\gamma_j} - X_{-\gamma_j}) \in \mathfrak{a}_0,$$

where $a_j \in \mathbf{R}$, there exists $H \in i\mathfrak{t}_0$ such that

$$(5.5) \quad i \text{ad}_H p = J_0 p.$$

Write $H = \sum_{\alpha \in \Delta_0^+} c_\alpha H_\alpha$, where H_α are the coroots. Then (5.5) is equivalent to the following system of linear equations:

$$\sum_{\alpha \in \Delta_0^+} c_\alpha(\alpha, \gamma_j) = 1, \quad j : 1, \dots, s,$$

which we can solve in each one of the three cases with the help of the tables in Appendix A.

Let $G_s = \mathbf{SU}(n) \times \mathbf{SU}(m)$, $n < m$, $\pi_\lambda : \overset{1}{\circ} \cdots \overset{1}{\circ} \otimes \overset{1}{\circ} \cdots \overset{1}{\circ}$. Then L_0 is $\mathbf{SU}(n+m)$,

$$\Delta_0^+ = \{\theta_i - \theta_j : 1 \leq i < j \leq n \text{ or } n+1 \leq i < j \leq n+m\},$$

$$\Phi = \{\theta_i - \theta_j : 1 \leq i \leq n \text{ and } n+1 \leq j \leq n+m\}$$

and $\Gamma = \{\theta_i - \theta_{n+m+1-i} : 1 \leq i \leq n\}$. Here we can take H to be a suitable multiple of $H_{\theta_m - \theta_{m+1}} + H_{\theta_m - \theta_{m+2}} + \dots + H_{\theta_m - \theta_{m+n}}$.

Let $G_s = \mathbf{SU}(n)$, n odd, $\pi_\lambda : \circ \cdots \circ$. Then L_0 is $\mathbf{SO}(2n)$, $\Delta_0^+ = \{\theta_i - \theta_j : 1 \leq i < j \leq n\}$, $\Phi = \{\theta_i + \theta_j : 1 \leq i < j \leq n\}$ and $\Gamma = \{\theta_{2i-1} + \theta_{2i} : 1 \leq i \leq \frac{n-1}{2}\}$. Here we can take H to be a suitable multiple of $H_{\theta_1 - \theta_n} + H_{\theta_3 - \theta_n} + \dots + H_{\theta_{n-2} - \theta_n}$.

Let $G_s = \mathbf{Spin}(10)$, $\pi_\lambda : \circ \cdots \circ \begin{array}{c} \circ \\ \diagup \quad \diagdown \\ \circ \end{array}$. Then L_0 is E_6 , $\Delta_0^+ = \{\pm\theta_i + \theta_j : 1 \leq i < j \leq 5\}$,

$\Phi = \{\frac{1}{2}(\theta_8 - \theta_7 - \theta_6 + \sum_{i=1}^5 \epsilon_i \theta_i) : \prod_{i=1}^5 \epsilon_i = +1\}$ and $\Gamma = \{\frac{1}{2}(\theta_8 - \theta_7 - \theta_6 + \theta_5 + \theta_4 + \theta_3 + \theta_2 + \theta_1), \frac{1}{2}(\theta_8 - \theta_7 - \theta_6 + \theta_5 - \theta_4 - \theta_3 - \theta_2 - \theta_1)\}$. Here we can take H to be a suitable multiple of $H_{\theta_4 + \theta_5} - H_{\theta_4 - \theta_5}$. This completes the proof of (a).

PROOF OF (b): We will verify directly that the second osculating space at the point

$$p = \sum_{j=1}^s X_{\gamma_j} - X_{-\gamma_j} \in \mathfrak{a}_0 \subset \mathfrak{p}_0$$

is not of maximal dimension. More precisely, we shall see that

$$(5.6) \quad \langle \text{ad}_X p, J_0 p \rangle = \langle \text{ad}_Y \text{ad}_X p, J_0 p \rangle = 0,$$

for all $X, Y \in \mathfrak{g}_s$. Notice that $J_0 p = \sum_{j=1}^s i(X_{\gamma_j} + X_{-\gamma_j}) \in \mathfrak{p}_0$.

It suffices to prove the identities (5.6) for the basis elements (4.3) of $\mathfrak{g}_s$. We have that $\langle \text{ad}_X p, J_0 p \rangle = \Re \ll \text{ad}_X p, Jp \gg$ and $\langle \text{ad}_Y \text{ad}_X p, J_0 p \rangle = -\Re \ll \text{ad}_X p, \text{ad}_Y Jp \gg$. The only cases where the real parts of the Hermitian products are not obviously zero are the following $(\alpha, \beta \in \Delta_0^+)$:

$$\begin{aligned} \ll \text{ad}_{H_\alpha} p, Jp \gg &= \sum_{j,k=1}^s \ll \text{ad}_{H_\alpha} (X_{\gamma_j} - X_{-\gamma_j}), X_{\gamma_k} + X_{-\gamma_k} \gg \\ &= \sum_{j,k=1}^s \gamma_j(H_\alpha) \ll X_{\gamma_j} + X_{-\gamma_j}, X_{\gamma_k} + X_{-\gamma_k} \gg \\ (5.7) \quad &= 2 \sum_{j=1}^s \gamma_j(H_\alpha) = 2 \left(\sum_{j=1}^s \gamma_j, \alpha \right), \end{aligned}$$

assuming the root vectors to have length one. We will show below that the last expression in (5.7) vanishes for the representations we are interested.

$$\begin{aligned}
\ll \text{ad}_{i(X_\alpha + X_{-\alpha})} p, Jp \gg &= \ll \text{ad}_{X_\alpha + X_{-\alpha}} p, \sum_{k=1}^s (X_{\gamma_k} + X_{-\gamma_k}) \gg \\
&= \sum_{j,k=1}^s \ll \text{ad}_{X_\alpha + X_{-\alpha}} (X_{\gamma_j} - X_{-\gamma_j}), X_{\gamma_k} + X_{-\gamma_k} \gg \\
(5.8) \qquad &= 0,
\end{aligned}$$

because $\text{ad}_{X_\alpha} X_{\gamma_j}$ is either a $(\gamma_j + \alpha)$ -root vector or 0; $\pm\gamma_j \pm \gamma_k \notin \Delta$; and root spaces corresponding to distinct roots are orthogonal.

$$\begin{aligned}
\ll \text{ad}_{iH_\alpha} p, \text{ad}_{X_\beta - X_{-\beta}} Jp \gg &= \sum_{j,k=1}^s \gamma_j(H_\alpha) \ll (X_{\gamma_j} + X_{-\gamma_j}), \text{ad}_{X_\beta - X_{-\beta}} (X_{\gamma_k} + X_{-\gamma_k}) \gg \\
&= 0,
\end{aligned}$$

for the same reason why (5.8) is zero.

$$\begin{aligned}
\Re \ll \text{ad}_{X_\alpha - X_{-\alpha}} Jp, \text{ad}_{i(X_\beta + X_{-\beta})} p \gg &= \Re \ll \text{ad}_{X_\alpha} Jp, i \text{ad}_{X_\beta} p \gg \\
&\quad - \Re \ll \text{ad}_{X_{-\alpha}} Jp, i \text{ad}_{X_{-\beta}} p \gg \\
&\quad + \Re \ll \text{ad}_{X_\alpha} Jp, i \text{ad}_{X_{-\beta}} p \gg \\
&\quad - \Re \ll \text{ad}_{X_{-\alpha}} Jp, i \text{ad}_{X_\beta} p \gg \\
(5.9) \qquad &= \Re \ll Jp, i \text{ad}_{[X_{-\beta}, X_\alpha]} p \gg \\
&\quad + \Re \ll Jp, i \text{ad}_{[X_\beta, X_\alpha]} p \gg,
\end{aligned}$$

because, for instance,

$$\begin{aligned}
\ll \text{ad}_{X_\alpha} Jp, i \text{ad}_{X_\beta} p \gg &= \ll \text{ad}_{X_{-\beta}} \text{ad}_{X_\alpha} Jp, ip \gg \\
&= \ll J \text{ad}_{X_{-\beta}} \text{ad}_{X_\alpha} p, ip \gg \\
&= -\ll \text{ad}_{X_{-\beta}} \text{ad}_{X_\alpha} p, Jip \gg \\
&= -\ll \text{ad}_{X_{-\beta}} \text{ad}_{X_\alpha} p, iJp \gg \\
&= \ll i \text{ad}_{X_{-\beta}} \text{ad}_{X_\alpha} p, Jp \gg \\
&= \overline{\ll Jp, i \text{ad}_{X_{-\beta}} \text{ad}_{X_\alpha} p \gg}.
\end{aligned}$$

Now (5.9) shows that everything amounts to checking that (5.7) vanishes. Next we do that in each one of the four cases. Let $\gamma = \sum_{j=1}^s \gamma_j$. Then the vanishing of (5.7) is equivalent to

$$(5.10) \qquad (\gamma, \Delta_0^+) = 0.$$

Let $G_s = \mathbf{SU}(n)$, $\pi_\lambda : \overset{2}{\circ} - \dots - \circ$. Then L_0 is $\mathbf{Sp}(n)$, $\Delta_0^+ = \{\theta_i - \theta_j : 1 \leq i < j \leq n\}$, $\Phi = \{\theta_i + \theta_j : 1 \leq i < j \leq n\} \cup \{2\theta_i : 1 \leq i \leq n\}$, $\Gamma = \{2\theta_i : 1 \leq i \leq n\}$ and $\gamma = 2\theta_1 + \dots + 2\theta_n$.

Let $G_s = \mathbf{SU}(n)$, n even, $\pi_\lambda : \overset{1}{\circ} \text{---} \overset{1}{\circ} \text{---} \dots \text{---} \overset{1}{\circ}$. Then L_0 is $\mathbf{SO}(2n)$, $\Delta_0^+ = \{\theta_i - \theta_j : 1 \leq i < j \leq n\}$, $\Phi = \{\theta_i + \theta_j : 1 \leq i < j \leq n\}$, $\Gamma = \{\theta_{2i-1} + \theta_{2i} : 1 \leq i \leq \frac{n}{2}\}$ and $\gamma = \theta_1 + \theta_2 + \dots + \theta_n$.

Let $G_s = \mathbf{SU}(n) \times \mathbf{SU}(n)$, $\pi_\lambda : \overset{1}{\circ} \text{---} \dots \text{---} \overset{1}{\circ} \otimes \overset{1}{\circ} \text{---} \dots \text{---} \overset{1}{\circ}$. Then L_0 is $\mathbf{SU}(2n)$, $\Delta_0^+ = \{\theta_i - \theta_j : 1 \leq i < j \leq n \text{ or } n+1 \leq i < j \leq 2n\}$, $\Phi = \{\theta_i - \theta_j : 1 \leq i \leq n \text{ and } n+1 \leq j \leq 2n\}$, $\Gamma = \{\theta_i - \theta_{2n+1-i} : 1 \leq i \leq n\}$ and $\gamma = \theta_1 + \theta_2 + \dots + \theta_n - \theta_{n+1} - \theta_{n+2} - \dots - \theta_{2n}$.

Let $G_s = \mathbf{E}_6$, $\pi_\lambda : \overset{1}{\circ} \text{---} \overset{1}{\circ} \text{---} \overset{1}{\circ} \text{---} \overset{1}{\circ} \text{---} \overset{1}{\circ}$. Then $L_0 = \mathbf{E}_7$,

$$\Delta_0^+ = \{\pm\theta_i + \theta_j : 1 \leq i < j \leq 5\} \cup \left\{ \frac{1}{2}(\theta_8 - \theta_7 - \theta_6 + \sum_{i=1}^5 \epsilon_i \theta_i) : \prod_{i=1}^5 \epsilon_i = +1 \right\},$$

$$\Phi = \{\theta_8 - \theta_7\} \cup \{\pm\theta_i + \theta_6 : 1 \leq i \leq 5\} \cup \left\{ \frac{1}{2}(\theta_8 - \theta_7 + \theta_6 + \sum_{i=1}^5 \epsilon_i \theta_i) : \prod_{i=1}^5 \epsilon_i = -1 \right\},$$

$\Gamma = \{\theta_8 - \theta_7, \theta_6 + \theta_5, \theta_6 - \theta_5\}$ and $\gamma = \theta_8 - \theta_7 + 2\theta_6$.


It is immediate to see that (5.10) holds in each case, and this completes the proof of (b). $\square$

Proof of Proposition 5.2. We know from Proposition 4.9 and Lemma 5.3 that $k(\lambda) = 1, 2$. In particular, G_s can have at most two simple factors.

Suppose $k(\lambda) = 1$. This implies G_s is simple. In the case $G = G_s$, we have π is of complex type, so $G = \mathbf{SU}(n)$ and π is the vector representation. In the case $G = S^1 \times G_s$, we have π_λ is of quaternionic or of complex type, so $G_s = \mathbf{Sp}(n)$ or $G_s = \mathbf{SU}(n)$ and π_λ is the respective vector representation. This only gives representations with cohomogeneity one.

Next suppose $k(\lambda) = 2$. The first case is $G_s = G_1 \times G_2$, where G_i is a simple group. Here π_λ decomposes as an outer tensor product $\pi_{\mu_1} \otimes \pi_{\mu_2}$, where π_{μ_i} is the representation of G_i of highest weight μ_i with $k(\mu_i) = 1$. This forces each G_i to be either $\mathbf{Sp}(n)$ or $\mathbf{SU}(n)$ for some n , and π_{μ_i} to be the corresponding vector representation. Using Lemma 5.3, item (b), and Lemma 5.4, item (b), we get the first three representations in the table.

The second case is G_s is a simple group. For $G = G_s$, an inspection of Table B.1 reveals four possibilities for π_λ , namely:



Now Lemma 5.4, item (b), restricts these to the next two representations in the table.

For $G = S^1 \times G_s$, if π_λ is of complex type, we only get representations associated to symmetric spaces, and if π_λ is of real type, Lemma 5.3, item (a), gives the last three representations in our table. $\square$

5.3 The case where ρ is of real type

The real case is much more involved than the previous two. We know from Proposition 4.9 that $k(\lambda) = 2, 4$. In particular, G can have at most four simple factors.

As a first step towards the classification in the case ρ is of real type, we introduce the following variations of condition (C_2) of Lemma 4.8 for a representation π_λ with representation space V_λ and highest weight vector v_λ :

$$\begin{aligned}\mathcal{U}^2(\mathfrak{g}^c)v_\lambda + \mathcal{U}^1(\mathfrak{g}^c)\epsilon(v_\lambda) &= V_\lambda & (C_{1\frac{1}{2}}) \\ \mathcal{U}^1(\mathfrak{g}^c)v_\lambda + \mathcal{U}^2(\mathfrak{g}^c)\epsilon(v_\lambda) &= V_\lambda & (C'_{1\frac{1}{2}}) \\ \mathcal{U}^1(\mathfrak{g}^c)v_\lambda + \mathcal{U}^1(\mathfrak{g}^c)\epsilon(v_\lambda) &= V_\lambda & (C_1) \\ \mathcal{U}^1(\mathfrak{g}^c)v_\lambda + \mathbf{C}\epsilon(v_\lambda) &= V_\lambda & (C_{\frac{1}{2}}) \\ \mathbf{C}v_\lambda + \mathcal{U}^1(\mathfrak{g}^c)\epsilon(v_\lambda) &= V_\lambda & (C'_{\frac{1}{2}})\end{aligned}$$

It follows from the identity $\epsilon X_\alpha = X_{-\alpha}\epsilon$ that conditions $(C_{1\frac{1}{2}})$ and $(C'_{1\frac{1}{2}})$ are equivalent, and that conditions $(C_{\frac{1}{2}})$ and $(C'_{\frac{1}{2}})$ are equivalent.

5.11 Lemma *The following conditions are equivalent for a complex irreducible representation π_λ of real type of a compact connected semisimple Lie group G , with invariant real form V :*

- (a) *condition (C_1) ;*
- (b) *G is transitive on the unit sphere of V ;*
- (c) *G acts with cohomogeneity 1 on V ;*
- (d) *$\{Xp : X \in \mathfrak{g}\} = (\mathbf{R}p)^\perp$ for all $p \in V$;*
- (e) *$\{Xp : X \in \mathfrak{g}\} = (\mathbf{R}p)^\perp$ for some $p \in V$.*

Proof. (b), (c), (d), and (e) are clearly equivalent. We prove that (a) is equivalent to (e). In fact, let $p = v_\lambda + \epsilon(v_\lambda)$. Clearly $\{Xp : X \in \mathfrak{g}\} \subset (\mathbf{R}p)^\perp$. Using the basis (4.3), we find that $\{Xp : X \in \mathfrak{g}\}^c$ is spanned over $\mathbf{C}$ by

$$v_\lambda - \epsilon(v_\lambda), \quad X_{-\alpha}v_\lambda, \quad X_\alpha\epsilon(v_\lambda), \quad \alpha \in \Delta^+.$$

But condition (C_1) holds precisely when this is also a system of generators for $(\mathbf{R}p)^\perp$ over $\mathbf{C}$. $\square$

5.12 Lemma *The following conditions are equivalent for a complex representation π_λ of quaternionic or complex type of a compact connected semisimple Lie group G on V_λ , with highest weight v_λ :*

- (a) $\mathcal{U}^1(\mathfrak{g}^c)v_\lambda = V_\lambda$;
- (b) *G is transitive on the unit sphere of V_λ^r ;*
- (c) *G acts with cohomogeneity 1 on V_λ^r ;*
- (d) *$\{Xp : X \in \mathfrak{g}\} = (\mathbf{R}p)^\perp$ for all $p \in V_\lambda^r$;*







(e) $\{Xp : X \in \mathfrak{g}\} = (\mathbf{R}p)^\perp$ for some $p \in V_\lambda^r$.

Proof. The proof is analogous to the proof of Lemma 5.11. $\square$

5.13 Lemma *The self-dual complex irreducible representations of compact connected simple Lie groups satisfying condition $(C_{\frac{1}{2}})$ are precisely⁴: the vector representation of $\mathbf{SO}(m)$ for $m \neq 2, 4$, the vector representation of $\mathbf{Sp}(m)$, the 7-dimensional representation of $\mathbf{G}_2$, and the spin representation of $\mathbf{Spin}(7)$.*

We will prove Lemma 5.13 together with the next lemma.

5.14 Lemma *The self-dual complex irreducible representations of compact connected simple Lie groups satisfying condition (C_1) are, besides those listed in Lemma 5.13, precisely the following:*

G	π_λ	$k(\lambda)$
$\mathbf{SU}(2)$		3
$\mathbf{SU}(6)$		3
$\mathbf{Spin}(9)$		2
$\mathbf{Spin}(11)$		3
$\mathbf{Spin}(12)$		3
$\mathbf{Sp}(3)$		3
$\mathbf{E}_7$		3

Proof of Lemmas 5.13 and 5.14. It is obvious that condition $(C_{\frac{1}{2}})$ implies condition (C_1) . So we consider (C_1) . It implies the following:

- (a) $k(\lambda) \leq 3$.
- (b) The dimension of π_λ is bounded by $\dim \mathfrak{g}^c - \text{rank } \mathfrak{g}^c + 2$.
- (c) If 0 is a weight, then its multiplicity is one. In particular, if π_λ is the adjoint representation, then the rank is at most one.

In fact, (a) follows from the fact that $k(\lambda) \geq 4$ contradicts Proposition 4.4. For (b) notice that the dimension of the left hand side of (C_1) is bounded by $2(\text{number of positive roots}) + 2$. Finally, if 0 is a weight, then the 0-weight space is contained in $\mathcal{U}^1(\mathfrak{g}^c)v_\lambda$ (because $\epsilon X_\alpha = X_{-\alpha}\epsilon$), and from this we see that λ is a root and the 0-weight space is in fact spanned by $X_{-\lambda}v_\lambda$, which implies (c).

⁴The half-spin representations of $\mathbf{Spin}(8)$ are image equivalent to the vector representation.

We will use the above observations to complete the proof of the lemmas.

Let π_λ be of real type. Then $k(\lambda) = 2$. The adjoint representations of $\mathbf{A}_n$, $\mathbf{B}_n$, $\mathbf{C}_n$, $\mathbf{D}_n$ with $n \geq 2$, and of $\mathbf{G}_2$, $\mathbf{F}_4$, $\mathbf{E}_6$, $\mathbf{E}_7$, $\mathbf{E}_8$ are immediately excluded. We run through all the remaining cases of representations of real type with $k(\lambda) = 2$.

Let G be of $\mathbf{A}_n$ -type. Then π_λ is $\circ$ or $\overset{1}{\circ} - \overset{1}{\circ} - \circ$ which are cited in the lemmas.

Let G be of $\mathbf{B}_n$ -type, $n \geq 2$. Then π_λ is $\overset{1}{\circ} - \cdots - \overset{1}{\circ} = \bullet$ or $\circ - \overset{1}{\circ} = \bullet$ or $\circ - \circ - \overset{1}{\circ} = \bullet$. All of them are cited in the lemmas.

Let G be of $\mathbf{C}_n$ -type, $n \geq 3$. Then π_λ is $\bullet - \bullet - \cdots - \bullet = \circ$, which cannot occur because the multiplicity of the 0-weight is $n - 1$.

Let G be of $\mathbf{D}_n$ -type, $n \geq 4$. Then π_λ is $\overset{1}{\circ} - \cdots - \overset{1}{\circ} \begin{array}{l} \circ \\ \circ \end{array}$, which is cited in the lemmas.

Let G be of exceptional type. Then π_λ is $\bullet = \circ$ or $\circ - \bullet = \bullet$. Only the former is cited in the lemmas; the latter has 0 as a weight of multiplicity 2.

Now let π_λ be of quaternionic type. Then $k(\lambda) = 1$ or $k(\lambda) = 3$. The case $k(\lambda) = 1$ gives the vector representation of $\mathbf{Sp}(m)$, which is cited in the lemmas. In the case $k(\lambda) = 3$, the only possibilities not cited in the lemmas are the following four representations, which we next eliminate:

- $G = \mathbf{B}_6$, π_λ : $\circ - \circ - \circ - \circ - \overset{1}{\circ} = \bullet$: here $\lambda = \frac{1}{2}(\theta_1 + \theta_2 + \theta_3 + \theta_4 + \theta_5 + \theta_6)$, $\mu = \frac{1}{2}(\theta_1 + \theta_2 + \theta_3 - \theta_4 - \theta_5 - \theta_6)$ is a weight and $\lambda \pm \mu$ are not roots which violates condition (C_1) .

In the remaining cases we make use of the bound on the dimension of the representation (see e. g. Table 5 in the Reference Chapter of [33]):



- $G = \mathbf{C}_n$, π_λ : $\overset{3}{\bullet} - \bullet - \cdots - \bullet = \circ$: $\dim \pi_\lambda = \frac{2}{3}n(n+1)(2n+1) > 2n^2+2 = \dim \mathbf{C}_n - n+2$, for $n \geq 2$;
- $G = \mathbf{C}_n$, π_λ : $\overset{1}{\bullet} - \overset{1}{\bullet} - \cdots - \bullet = \circ$: $\dim \pi_\lambda = \frac{8}{3}n(n-1)(n+1) > 2n^2+2 = \dim \mathbf{C}_n - n+2$, for $n \geq 2$;
- $G = \mathbf{C}_n$, π_λ : $\bullet - \bullet - \overset{1}{\bullet} - \cdots - \bullet = \circ$: $\dim \pi_\lambda = \frac{2}{3}n(n-2)(2n+1) > 2n^2+2 = \dim \mathbf{C}_n - n+2$, for $n \geq 4$;

On the other hand, the representations listed in the statements of the Lemmas 5.13 and 5.14 do satisfy (C_1) , as can be easily checked using the fact that each of them has all weights with multiplicities one. This concludes the classification of self-dual representations that satisfy (C_1) . As for condition $(C_{\frac{1}{2}})$, we observe that $(C_{\frac{1}{2}})$ implies that $k(\lambda) \leq 2$. This eliminates all representations in the table of Lemma 5.14 except $\circ - \circ - \overset{1}{\circ} = \bullet$; that one can be eliminated by a direct check. Finally it is easy to verify that the representations cited in Lemma 5.13 do satisfy $(C_{\frac{1}{2}})$. $\square$

5.15 Lemma *The only complex irreducible representation of real type of a compact connected semisimple nonsimple Lie group G satisfying condition (C_1) is the tensor product of the vector representations of $\mathbf{Sp}(1)$ and $\mathbf{Sp}(m)$.*

Proof. We already know that $k(\lambda) = 2$ (cf. proof of Lemmas 5.13 and 5.14), so $G = G_1 \times G_2$, where G_i is a simple group, and $\pi_\lambda = \pi_{\mu_1} \otimes \pi_{\mu_2}$ with $k(\mu_i) = 1$. Since each π_{μ_i} is self-dual, it must be of quaternionic type. Therefore, each $G_i = \mathbf{Sp}(n_i)$ and the corresponding π_{μ_i} is just the vector representation. Now one of the n_i must be 1, for otherwise condition (C_1) for π_λ is violated by consideration of a weight vector of type $v_1 \otimes v_2$, where v_i is an *intermediate* (that is, neither highest nor lowest) weight vector for π_{μ_i} . $\square$

5.16 Lemma *The complex irreducible representations π_λ of quaternionic type of compact connected simple Lie groups G satisfying $k(\lambda) = 3$ and condition $(C_{1\frac{1}{2}})$, but not condition (C_1) , are the following:*

G	π_λ
$\mathbf{Spin}(13)$	
$\mathbf{Sp}(2)$	

We postpone the proof of Lemma 5.16 to the end of Subsection 5.3.5, since the methods used to prove it better belong there.

5.3.1 The case where G has four simple factors

In this case we have:

5.17 Proposition *Let ρ be of real type and $G = G_1 \times G_2 \times G_3 \times G_4$, where G_i is a simple group. Then $G = \mathbf{SU}(2) \times \mathbf{SU}(2) \times \mathbf{SU}(2) \times \mathbf{Sp}(n)$, $n \geq 2$, and π is the tensor product of the vector representations of each of the factors.*

Proof. Set $\pi_\lambda = \pi_{\mu_1} \otimes \pi_{\mu_2} \otimes \pi_{\mu_3} \otimes \pi_{\mu_4}$, where π_{μ_i} is the representation of G_i of highest weight μ_i . We have $k(\mu_i) = 1$ for all i , and since each π_{μ_i} is self-dual, it must be of quaternionic type. Therefore, each $G_i = \mathbf{Sp}(n_i)$ and the corresponding π_{μ_i} is just the vector representation. Notice that $n_i = 1$ for all i gives the isotropy representation of a symmetric space. Suppose now that for two indices, say 3 and 4, we have $n_3 > 1$ and $n_4 > 1$. Take weight vectors v_i of π_{μ_i} such that: v_1 is the highest weight vector of π_{μ_1} , v_2 is the lowest weight vector of π_{μ_2} , and v_3 and v_4 are intermediate weight vectors of π_{μ_3} and π_{μ_4} , respectively. Then $v_1 \otimes v_2 \otimes v_3 \otimes v_4$ is a weight vector of π_λ which cannot be contained in $\mathcal{U}^2(\mathfrak{g}^e)v_\lambda + \mathcal{U}^2(\mathfrak{g}^e)\epsilon(v_\lambda)$, violating the condition (C_2) from Lemma 4.8. $\square$

5.3.2 The case where G has three simple factors

The classification is:

5.18 Proposition *Let ρ be of real type and $G = G_1 \times G_2 \times G_3$, where G_i is a simple group. Then ρ is one of the following:*

G	ρ	Conditions
$\mathbf{SU}(2) \times \mathbf{Sp}(n) \times \mathbf{SO}(2m)$		$m \geq 3, \quad n \geq 2$
$\mathbf{SU}(2) \times \mathbf{Sp}(n) \times \mathbf{SO}(2m+1)$		$n \geq 2$
$\mathbf{SU}(2) \times \mathbf{Sp}(n) \times \mathbf{G}_2$		—
$\mathbf{SU}(2) \times \mathbf{Sp}(n) \times \mathbf{Spin}(7)$		—
$\mathbf{SU}(2) \times \mathbf{SU}(2) \times \mathbf{Spin}(9)$		—

Proof. Let $\pi_\lambda = \pi_{\mu_1} \otimes \pi_{\mu_2} \otimes \pi_{\mu_3}$, where π_{μ_i} is the representation of G_i of highest weight μ_i , say with $k(\mu_1) = k(\mu_2) = 1$, $k(\mu_3) = 2$. Then π_{μ_1}, π_{μ_2} are each of quaternionic type and π_{μ_3} is of real type. So each of π_{μ_1}, π_{μ_2} is the vector representation of $\mathbf{Sp}(m)$ for some m . Moreover, one of them must have $m = 1$, say π_{μ_1} , for otherwise condition (C_2) for π_λ is violated by consideration of a weight vector of type $v_1 \otimes v_2 \otimes v_3$, where v_i is an intermediate weight vector for π_{μ_i} . Now we have $G_1 = \mathbf{SU}(2)$, $G_2 = \mathbf{Sp}(n)$, and π_{μ_1}, π_{μ_2} the respective vector representations. Next take a weight vector $v_1 \otimes v_2 \otimes v_3$ for π_λ such that v_1 is highest, v_2 is lowest and v_3 is intermediate. Condition (C_2) for π_λ forces π_{μ_3} to satisfy condition (C_1) . If $n \geq 2$, we could also take v_2 to be intermediate, and then condition (C_2) for π_λ implies something stronger, namely that π_{μ_3} must satisfy condition $(C_{\frac{1}{2}})$. Our table now follows from Lemmas 5.13 and 5.14. $\square$

5.3.3 The case where G has two simple factors and ρ has two factors of real type

Here the result is:

5.19 Proposition *Let ρ be of real type and $G = G_1 \times G_2$, where G_i is a simple group. Write $\rho^c = \pi_\lambda$, decompose $\pi_\lambda = \pi_{\mu_1} \otimes \pi_{\mu_2}$ and suppose that each π_{μ_i} is of real type. Then each π_{μ_i} is one of the following representations: the vector representation of $\mathbf{SO}(m)$ for $m \neq 2, 4$, or the 7-dimensional representation of $\mathbf{G}_2$, or the spin representation of $\mathbf{Spin}(7)$, or the spin representation of $\mathbf{Spin}(9)$; but the case where $G_1 = G_2 = \mathbf{Spin}(9)$ is excluded.*

Proof. Here $k(\mu_1) = k(\mu_2) = 2$. Let $v_1 \otimes v_2$ be a weight vector for π_λ such that each v_i is an intermediate weight vector for π_{μ_i} . Condition (C_2) for π_λ applied to $v_1 \otimes v_2$ forces each π_{μ_i} to satisfy (C_1) . In fact, one of the π_{μ_i} must even satisfy $(C_{\frac{1}{2}})$. The claim follows from Lemmas 5.13 and 5.14. $\square$

5.3.4 The case where G has two simple factors and ρ has two factors of quaternionic type

The classification is:

5.20 Proposition *Let ρ be of real type and $G = G_1 \times G_2$, where G_i is a simple group. Write $\rho^c = \pi_\lambda$, decompose $\pi_\lambda = \pi_{\mu_1} \otimes \pi_{\mu_2}$ and suppose that each π_{μ_i} is of quaternionic type. Then ρ is one of the following:*

G	ρ	Conditions
$\mathbf{Sp}(n) \times \mathbf{SU}(2)$		$n \geq 2$
$\mathbf{Sp}(n) \times \mathbf{SU}(6)$		$n \geq 2$
$\mathbf{Sp}(n) \times \mathbf{Spin}(11)$		—
$\mathbf{Sp}(n) \times \mathbf{Spin}(12)$		$n \geq 2$
$\mathbf{Sp}(1) \times \mathbf{Spin}(13)$		—
$\mathbf{Sp}(1) \times \mathbf{Sp}(2)$		—
$\mathbf{Sp}(n) \times \mathbf{Sp}(3)$		$n \geq 2$
$\mathbf{Sp}(n) \times \mathbf{E}_7$		$n \geq 2$

Proof. If $k(\lambda) = 2$, then $k(\mu_1) = k(\mu_2) = 1$ and each π_{μ_i} must be the vector representation of $\mathbf{Sp}(n_i)$ for some n_i , so we get a representation associated to a symmetric space. In the case $k(\lambda) = 4$ we can write $k(\mu_1) = 1$ and $k(\mu_2) = 3$. Now π_{μ_1} must be the vector representation of $\mathbf{Sp}(n)$ and there are two cases to consider:

- $n = 1$: here π_{μ_2} must satisfy condition $(C_{1\frac{1}{2}})$;
- $n \geq 2$: here π_{μ_2} must satisfy the stronger condition (C_1) .





The claim now follows from Lemmas 5.14 and 5.16. □

5.3.5 The case where G is a simple group

The representations of real type of simple groups that are candidates to a position in class $\mathcal{O}^2$ are:





5.21 Proposition *Let ρ be of real type and suppose G is a simple group. Then ρ is one of*














the following:






G	ρ
Spin(7)	
Spin(9)	
Spin(15)	
Spin(17)	









We will accomplish the proof of the above proposition by running through all the cases of representations of real type of a simple group with $k(\lambda) \leq 4$ given in Table B.1. In fact, according to that table, all representations with $k(\lambda) = 2$ are either associated to symmetric spaces or of cohomogeneity one. So we only need to worry about the case $k(\lambda) = 4$. A careful inspection of Table B.1 shows:

5.22 Lemma *Let π_λ be a representation of real type of a compact connected simple Lie group G with $k(\lambda) = 4$. Then (G, π_λ) is one of the following:*

	G	π_λ	Conditions
1	A₁		—
2	A₃		—
3	A₃		—
4	A₇		—
5	A_n		$n \geq 2$
6	A_n		$n \geq 4$

	G	π_λ	Conditions
7	B_3		—
8	B_3		—
9	B_3		—
10	B_4		—
11	B_4		—
12	B_4		—
13	B_7		—
14	B_8		—
15	B_n		$n \geq 2$
16	B_n		$n \geq 3$
17	B_n		$n \geq 3$
18	B_n		$n \geq 4$
19	B_n		$n \geq 5$

	G	π_λ	Conditions
20	C_n		$n \geq 2$
21	C_n		$n \geq 2$
22	C_n		$n \geq 3$
23	C_n		$n \geq 3$
24	C_n		$n \geq 4$

	G	π_λ	Conditions
25	$\mathbf{D}_4$		—
26	$\mathbf{D}_5$		—
27	$\mathbf{D}_8$		—
28	$\mathbf{D}_n$		$n \geq 4$
29	$\mathbf{D}_n$		$n \geq 4$
30	$\mathbf{D}_n$		$n \geq 4$
31	$\mathbf{D}_n$		$n \geq 5$
32	$\mathbf{D}_n$		$n \geq 6$

	G	π_λ
33	G_2	
34	G_2	
35	G_2	
36	F_4	
37	F_4	
38	F_4	
39	F_4	
40	E_6	
41	E_6	
42	E_7	
43	E_7	
44	E_8	
45	E_8	

To begin with, we note the representations listed in the tables of Lemma 5.22 associated to a symmetric space, namely, numbers 1, 2, 4, 15, 24 ($n = 4$), 27 and 28.

5.23 Lemma *Let π_λ be a representation of real type with $k(\lambda) = 4$. Assume that $\lambda = \frac{1}{2}(3\beta_i + \beta_j)$ where $\beta_i, \beta_j \in \mathcal{B}$ are distinct long roots and that there exists a Weyl group involution s such that $s\beta_i = \beta_j$. Then a real form of π_λ is not of class $\mathcal{O}^2$.*

Proof. We have that $\mu = ss_{\beta_i}(\lambda) = \frac{1}{2}(\beta_i - 3\beta_j)$ is a weight and $\lambda - \mu = \beta_i + 2\beta_j$, $\lambda + \mu = 2\beta_i - \beta_j$. Since β_i and β_j are orthogonal, we get that $\|\lambda \pm \mu\|^2 = 5$ (normalizing the length of a long root to be 1). On the other hand, for $\alpha', \alpha'' \in \Delta$ we have that $\|\alpha' + \alpha''\|^2 \leq (\|\alpha'\| + \|\alpha''\|)^2 \leq 4$, so $\lambda \pm \mu$ is neither a root nor a sum of two roots. Thus a μ -weight vector is not in $\mathcal{U}^2(\mathfrak{g}^c)v_\lambda + \mathcal{U}^2(\mathfrak{g}^c)\epsilon(v_\lambda)$, violating condition (C_2) . $\square$

One can use the last lemma together with the tables in Appendix A to get rid of the representations listed in Lemma 5.22 under the numbers 3, 12, 17, 21, 30 and 37.

On the other hand, we find that:

$$\begin{array}{c} \overset{1}{\circ} - \overset{1}{\circ} - \dots - \overset{1}{\circ} \otimes \overset{1}{\circ} - \dots - \overset{1}{\circ} = \\ \overset{1}{\circ} - \overset{1}{\circ} - \dots - \overset{1}{\circ} \oplus \overset{1}{\circ} - \dots - \overset{1}{\circ} \oplus (\text{trivial}), \end{array}$$

from where we deduce that the multiplicity of the 0-weight in π_λ is $\frac{n(n+1)}{2} - n - 1 = \frac{n^2 - n - 2}{2} > 2$, so we can apply Lemma 5.26. $\square$

5.28 Lemma *Representations numbers 18, 10 and 19 are not of class $\mathcal{O}^2$.*

Proof. We have that

$$\begin{array}{l} \overset{1}{\circ} - \overset{1}{\circ} - \dots - \overset{1}{\circ} = \Lambda^3(\overset{1}{\circ} - \overset{1}{\circ} - \dots - \overset{1}{\circ}), \quad n \geq 4, \\ \overset{1}{\circ} - \overset{1}{\circ} - \dots - \overset{1}{\circ} = \Lambda^4(\overset{1}{\circ} - \overset{1}{\circ} - \dots - \overset{1}{\circ}), \\ \overset{1}{\circ} - \overset{1}{\circ} - \dots - \overset{1}{\circ} = \Lambda^4(\overset{1}{\circ} - \overset{1}{\circ} - \dots - \overset{1}{\circ}), \quad n \geq 5. \end{array}$$

This shows that the multiplicity of the 0-weight is, respectively, n , 6 and $\binom{n}{2}$ (because the vector representation $\overset{1}{\circ} - \dots - \overset{1}{\circ}$ has weights $\theta_1, \theta_2, \dots, \theta_n, 0, -\theta_n, -\theta_{n-1}, \dots, -\theta_1$). But the integer $N = 3$ in all cases. $\square$

5.29 Lemma *Representation number 9 is not of class $\mathcal{O}^2$.*

Proof. The highest weight is $\lambda = \frac{1}{2}(3\theta_1 + 3\theta_2 + \theta_3)$. Now $\mu = \frac{1}{2}(3\theta_1 - 3\theta_2 + \theta_3)$ is a weight and $\lambda + \mu = 3\theta_1 + \theta_3$, $\lambda - \mu = 3\theta_2$ are neither a root nor a sum of two roots, so we are done as in the proof of Lemma 5.23. $\square$

5.30 Lemma *Representations numbers 22 and 23 are not of class $\mathcal{O}^2$. Representation number 24 is not of class $\mathcal{O}^2$ if $n \geq 5$.*

Proof. We first observe that:

- the vector representation $C_n : \overset{1}{\bullet} - \overset{1}{\bullet} - \dots - \overset{1}{\bullet} = \overset{1}{\circ}$ has weights $\pm\theta_i$ with multiplicity 1;
- the representation $\overset{1}{\bullet} - \overset{1}{\bullet} - \dots - \overset{1}{\bullet} = \overset{1}{\circ}$ is a subrepresentation of $\Lambda^2(\overset{1}{\bullet} - \dots - \overset{1}{\bullet})$ and has weights $\pm\theta_i \pm \theta_j$ with multiplicity 1, and 0 with multiplicity $n - 1$;
- the representation $\overset{1}{\bullet} - \overset{1}{\bullet} - \dots - \overset{1}{\bullet} = \overset{1}{\circ}$ is a subrepresentation of $\Lambda^3(\overset{1}{\bullet} - \dots - \overset{1}{\bullet})$ and has weights $\pm\theta_i \pm \theta_j \pm \theta_k$ with multiplicity 1, and $\pm\theta_i$ with multiplicity $n - 2$;
- the representation $\overset{1}{\bullet} - \overset{1}{\bullet} - \dots - \overset{1}{\bullet} = \overset{1}{\circ}$ is a subrepresentation of

$$\Lambda^4(\overset{1}{\bullet} - \dots - \overset{1}{\bullet})$$

and has weights $\pm\theta_i \pm \theta_j \pm \theta_k \pm \theta_l$ with multiplicity 1, and $\pm\theta_i \pm \theta_j$ with multiplicity $n - 1$, and 0 with multiplicity $\binom{n}{2} - n$.

If $n \geq 5$ then the multiplicity of the 0-weight in $\bullet \text{---} \bullet \text{---} \bullet \text{---} \overset{1}{\bullet} \cdots \text{---} \bullet \text{---} \circ$ is $\frac{1}{2}n(n-3) \geq 5$. Since $\lambda = \theta_1 + \theta_2 + \theta_3 + \theta_4$ can be written as a root or as a sum of two roots in $N = 3$ ways only, namely,

$$\begin{aligned}\lambda &= (\theta_1 + \theta_2) + (\theta_3 + \theta_4) \\ &= (\theta_1 + \theta_3) + (\theta_2 + \theta_4) \\ &= (\theta_1 + \theta_4) + (\theta_2 + \theta_3),\end{aligned}$$

Lemma 5.26 says that $\bullet \text{---} \bullet \text{---} \overset{1}{\bullet} \text{---} \cdots \text{---} \bullet \equiv \circ$, $n \geq 5$, cannot be of class $\mathcal{O}^2$.

Next, it follows from

$$\begin{array}{l}
\bullet \xrightarrow{1} \bullet \xrightarrow{1} \bullet \otimes \bullet \xrightarrow{1} \bullet \xrightarrow{1} \bullet = \bullet \xrightarrow{1} \bullet \xrightarrow{1} \bullet \oplus \bullet \xrightarrow{1} \bullet \xrightarrow{1} \bullet, \\
\bullet \xrightarrow{1} \bullet \xrightarrow{1} \bullet \cdots \bullet \xrightarrow{1} \bullet \otimes \bullet \xrightarrow{1} \bullet \xrightarrow{1} \bullet \cdots \bullet \xrightarrow{1} \bullet = \\
\bullet \xrightarrow{1} \bullet \xrightarrow{1} \bullet \cdots \bullet \xrightarrow{1} \bullet \oplus \bullet \xrightarrow{1} \bullet \xrightarrow{1} \bullet \cdots \bullet \xrightarrow{1} \bullet \oplus \bullet \xrightarrow{1} \bullet \xrightarrow{1} \bullet \cdots \bullet \xrightarrow{1} \bullet, \quad n \geq 4,
\end{array}$$

that the multiplicity of the 0-weight in $\overset{1}{\bullet} \text{---} \overset{1}{\bullet} \text{---} \bullet \cdots \text{---} \bullet \text{---} \circ$, $n \geq 3$, is given by $\frac{1}{2}(n - 2)(3n - 1) \geq 4$. But $\lambda = 2\theta_1 + \theta_2 + \theta_3$ can be written as a root or as sum of two roots in $N = 2$ ways only:

$$\begin{aligned}\lambda &= (\theta_1 + \theta_2) + (\theta_1 + \theta_3) \\ &= 2\theta_1 + (\theta_2 + \theta_3).\end{aligned}$$

Lemma 5.26 now implies that $\overset{1}{\bullet} \overset{1}{\text{---}} \underset{1}{\bullet} \text{---} \cdots \text{---} \bullet \text{---} \circ$, $n \geq 3$, is not of class $\mathcal{O}^2$.

One can also see that $S^2(\bullet\text{---}\bullet\cdots\text{---}\bullet\text{=}\circ)$, $n \geq 3$, has 0 as a weight of multiplicity $\frac{3}{2}n(n-1)$. Now

$$\begin{aligned}
S^2(\overset{1}{\bullet} \text{---} \overset{1}{\bullet} \text{---} \bigcirc) &= \overset{2}{\bullet} \text{---} \overset{2}{\bullet} \text{---} \bigcirc \oplus \overset{1}{\bullet} \text{---} \overset{1}{\bullet} \text{---} \bigcirc \oplus (\text{trivial}), \\
S^2(\overset{1}{\bullet} \text{---} \overset{1}{\bullet} \text{---} \dots \text{---} \bullet \text{---} \bigcirc) &= \\
\overset{2}{\bullet} \text{---} \overset{2}{\bullet} \text{---} \dots \text{---} \bullet \text{---} \bigcirc \oplus \overset{1}{\bullet} \text{---} \overset{1}{\bullet} \text{---} \dots \text{---} \bullet \text{---} \bigcirc \oplus \dots \oplus \overset{1}{\bullet} \text{---} \dots \text{---} \bullet \text{---} \bigcirc \oplus (\text{trivial}), \quad n \geq 4,
\end{aligned}$$

so the multiplicity of the 0-weight in $\bullet \overset{2}{\bullet} \cdots \bullet \text{---} \bullet \text{---} \circ$, $n \geq 3$, is $n(n-1) \geq 6$. But $\lambda = 2\theta_1 + 2\theta_2$ can be written as a root or as sum of two roots in $N = 2$ ways only:

$$\begin{aligned}\lambda &= (\theta_1 + \theta_2) + (\theta_1 + \theta_2) \\ &= 2\theta_1 + 2\theta_2.\end{aligned}$$

Lemma 5.26 implies again that $\overset{2}{\bullet} \text{---} \bullet \text{---} \cdots \text{---} \bullet \text{=}\circ$, $n \geq 3$, is not of class $\mathcal{O}^2$.

5.31 Lemma *Representations numbers 26 and 32 are not of class $\mathcal{O}^2$.*

Proof. These representations are just

$$\mathbf{D}_n : \Lambda^4(\overset{1}{\circ} \cdots \overset{1}{\circ} \begin{array}{c} \circ \\ \circ \end{array}), \quad n \geq 5,$$

where $\overset{1}{\circ} \cdots \overset{1}{\circ} \begin{array}{c} \circ \\ \circ \end{array}$ is the vector representation. Therefore the multiplicity of the 0-weight is $\binom{n}{2} \geq 10$. But the highest weight $\lambda = \theta_1 + \theta_2 + \theta_3 + \theta_4$ decomposes as a sum of two roots in $N = 3$ ways, so we can use Lemma 5.26. $\square$

5.32 Lemma *Representation number 35 is not of class $\mathcal{O}^2$.*

Proof. Here $\lambda = \lambda_1 + \lambda_2 = 5\alpha_1 + 3\alpha_2$ decomposes only as $(2\alpha_1 + \alpha_2) + (3\alpha_1 + 2\alpha_2)$, so that $N = 1$. Now we find that:

$$\overset{1}{\bullet} \equiv \overset{1}{\circ} \otimes \overset{1}{\bullet} \equiv \overset{1}{\bullet} \oplus \overset{1}{\bullet} \oplus \overset{2}{\bullet} \oplus \overset{1}{\bullet},$$

where $\overset{1}{\bullet} \equiv \overset{1}{\circ}$ is the vector representation and $S^2(\overset{1}{\bullet} \equiv \overset{1}{\circ}) = \overset{2}{\bullet} \oplus (\text{trivial})$. Therefore the multiplicity of the 0-weight in $\overset{1}{\bullet} \equiv \overset{1}{\circ}$ is 4 and we can use Lemma 5.26. $\square$

5.33 Lemma *Representations numbers 38 and 39 are not of class $\mathcal{O}^2$.*

Proof. First we observe that the 26-dimensional representation $\mathbf{F}_4 : \overset{1}{\circ} \cdots \overset{1}{\bullet} \equiv \overset{1}{\bullet}$ has weights $\frac{1}{2}(\pm\theta_1 \pm \theta_2 \pm \theta_3 \pm \theta_4)$ and $\pm\theta_i$ with multiplicity 1, and 0 with multiplicity 2. Therefore $\Lambda^2(\overset{1}{\circ} \cdots \overset{1}{\bullet} \equiv \overset{1}{\bullet})$ has 0 as a weight of multiplicity 13. Now

$$\Lambda^2(\overset{1}{\circ} \cdots \overset{1}{\bullet} \equiv \overset{1}{\bullet}) = \overset{1}{\circ} \cdots \overset{1}{\bullet} \equiv \overset{1}{\bullet} \oplus \overset{1}{\circ} \cdots \overset{1}{\bullet} \equiv \overset{1}{\bullet},$$

where $\overset{1}{\circ} \cdots \overset{1}{\bullet} \equiv \overset{1}{\bullet}$ is the adjoint representation. It follows that $\pi_\lambda : \overset{1}{\circ} \cdots \overset{1}{\bullet} \equiv \overset{1}{\bullet}$ has 0 as a weight of multiplicity 9. But its highest weight $\lambda = \frac{1}{2}(3\theta_1 + \theta_2 + \theta_3 + \theta_4)$ is not a root and decomposes as a sum of two roots as follows:

$$\begin{aligned} \lambda &= \frac{1}{2}(\theta_1 + \theta_2 + \theta_3 + \theta_4) + \theta_1 \\ &= \frac{1}{2}(\theta_1 - \theta_2 + \theta_3 + \theta_4) + (\theta_1 + \theta_2) \\ &= \frac{1}{2}(\theta_1 + \theta_2 - \theta_3 + \theta_4) + (\theta_1 + \theta_3) \\ &= \frac{1}{2}(\theta_1 + \theta_2 + \theta_3 - \theta_4) + (\theta_1 + \theta_4). \end{aligned}$$

Thus the integer of Lemma 5.26 is $N = 4$ and $\overset{1}{\circ} \cdots \overset{1}{\bullet} \equiv \overset{1}{\bullet}$ is out.

Similarly, the multiplicity of the 0 weight in $S^2(\overset{1}{\circ} \cdots \overset{1}{\bullet} \equiv \overset{1}{\bullet})$ is 15, and

$$S^2(\overset{1}{\circ} \cdots \overset{1}{\bullet} \equiv \overset{1}{\bullet}) = \overset{2}{\bullet} \oplus \overset{1}{\circ} \cdots \overset{1}{\bullet} \equiv \overset{1}{\bullet} \oplus (\text{trivial}).$$


It follows that $\pi_\lambda : \circ \text{---} \circ \text{---} \overset{2}{\bullet} \text{---} \bullet$ has 0 as a weight of multiplicity 12. But its highest weight $\lambda = 2\theta_1$ is not a root and decomposes as a sum of two roots as follows:

$$\begin{aligned}\lambda &= (\theta_1) + (\theta_1) \\ &= (\theta_1 + \theta_2) + (\theta_1 - \theta_2) \\ &= (\theta_1 + \theta_3) + (\theta_1 - \theta_3) \\ &= (\theta_1 + \theta_4) + (\theta_1 - \theta_4).\end{aligned}$$

Thus the integer of Lemma 5.26 is $N = 4$ and $\circ \text{---} \circ \text{---} \overset{2}{\bullet} \text{---} \bullet$ is out, too. $\square$

5.34 Lemma *Representation number 40 is not of class $\mathcal{O}^2$.*

Proof. We have that



are respectively the 27-dimensional representation and its dual. Therefore the multiplicity of the 0-weight in the tensor product of these representations is at least 27. But

$$\begin{array}{c} 1 \\ \circ \text{---} \circ \text{---} \circ \text{---} \circ \\ | \\ \circ \end{array} \otimes \begin{array}{c} 1 \\ \circ \text{---} \circ \text{---} \circ \text{---} \circ \\ | \\ \circ \end{array} = \begin{array}{c} 1 \\ \circ \text{---} \circ \text{---} \circ \text{---} \circ \\ | \\ \circ \end{array} \oplus \begin{array}{c} 1 \\ \circ \text{---} \circ \text{---} \circ \text{---} \circ \\ | \\ \circ \end{array} \oplus (\text{trivial})$$

where the second summand on the right hand side is the adjoint representation. It follows

that the multiplicity of the 0-weight in $\pi_\lambda : \begin{array}{c} 1 \\ \circ \text{---} \circ \text{---} \circ \text{---} \circ \\ | \\ \circ \end{array}$ is at least 20. But its highest weight $\lambda = \theta_8 - \theta_7 - \theta_6 + \theta_5$ is not a root and decomposes as a sum of two roots as follows:

$$\begin{aligned}\lambda &= \frac{1}{2}(\theta_8 - \theta_7 - \theta_6 + \theta_5 + \theta_4 + \theta_3 + \theta_2 + \theta_1) + \frac{1}{2}(\theta_8 - \theta_7 - \theta_6 + \theta_5 - \theta_4 - \theta_3 - \theta_2 - \theta_1) \\ &= \frac{1}{2}(\theta_8 - \theta_7 - \theta_6 + \theta_5 + \theta_4 + \theta_3 - \theta_2 - \theta_1) + \frac{1}{2}(\theta_8 - \theta_7 - \theta_6 + \theta_5 - \theta_4 - \theta_3 + \theta_2 + \theta_1) \\ &= \frac{1}{2}(\theta_8 - \theta_7 - \theta_6 + \theta_5 + \theta_4 - \theta_3 + \theta_2 - \theta_1) + \frac{1}{2}(\theta_8 - \theta_7 - \theta_6 + \theta_5 - \theta_4 + \theta_3 - \theta_2 + \theta_1) \\ &= \frac{1}{2}(\theta_8 - \theta_7 - \theta_6 + \theta_5 - \theta_4 + \theta_3 + \theta_2 - \theta_1) + \frac{1}{2}(\theta_8 - \theta_7 - \theta_6 + \theta_5 + \theta_4 - \theta_3 - \theta_2 + \theta_1).\end{aligned}$$

Thus the integer of Lemma 5.26 is $N = 4$ and we are done. $\square$

5.35 Lemma *Representation number 43 is not of class $\mathcal{O}^2$.*

Proof. We have that $\begin{array}{c} 1 \\ \circ \text{---} \circ \text{---} \circ \text{---} \circ \\ | \\ \circ \end{array}$ is the 56-dimensional representation. It is self-dual and does not have 0 as a weight. Therefore the multiplicity of the 0-weight in the exterior square of this representation is 28. But

$$\Lambda^2 \left(\begin{array}{c} 1 \\ \circ \text{---} \circ \text{---} \circ \text{---} \circ \\ | \\ \circ \end{array} \right) = \begin{array}{c} 1 \\ \circ \text{---} \circ \text{---} \circ \text{---} \circ \\ | \\ \circ \end{array} \oplus (\text{trivial}).$$

Therefore the multiplicity of the 0-weight in π_λ is 27. But its highest weight $\lambda = \theta_8 - \theta_7 + \theta_6 + \theta_5$ is not a root and one can easily check that it decomposes as a sum of two roots in exactly $N = 17$ different ways. $\square$

5.36 Lemma *Representation number 44 is not of class $\mathcal{O}^2$.*

Proof. Here we slightly change the argument used in the previous lemmas. The highest weight $\lambda = 2\theta_8$ and $(\lambda, \theta_8 - \theta_7) \neq 0$, so $\nu = \lambda - (\theta_8 - \theta_7) = \theta_8 + \theta_7$ is a weight. Now $\theta_8 + \theta_7$ and $\theta_6 + \theta_5$ are long roots of $\mathbf{E}_8$, hence in the same Weyl orbit. It follows that $\mu = \theta_6 + \theta_5$ is a weight, too. Observe that each of $\lambda \pm \mu$ is not a root and can be decomposed as a sum of two roots in only one way:

$$\begin{aligned}\lambda + \mu &= (\theta_8 + \theta_6) + (\theta_8 + \theta_5), \\ \lambda - \mu &= (\theta_8 - \theta_6) + (\theta_8 - \theta_5).\end{aligned}$$

The proof will be complete if we show that the multiplicity of μ as a weight is greater than 2, since then we will have that condition (C_2) is violated.

In fact, the multiplicity of μ is the same as the multiplicity of ν . We use Freudenthal's formula as it is stated in [18], namely, the multiplicity of ν as a weight of π_λ is given by the following formula:

$$m_\nu = \frac{2}{c(\nu)} \sum_{\alpha \in \Delta^+} \sum_{k \geq 1} (\nu + k\alpha, \alpha) m_{\nu+k\alpha},$$

where $c(\nu) = \|\lambda + \rho\|^2 - \|\nu + \rho\|^2$, $\rho = \frac{1}{2} \sum_{\alpha \in \Delta^+} \alpha$ and $m_{\nu+k\alpha}$ is the multiplicity of $\nu + k\alpha$ as a weight of π_λ .

We content ourselves with an estimate. Using that $\rho = \theta_2 + 2\theta_3 + 3\theta_4 + 4\theta_5 + 5\theta_6 + 6\theta_7 + 23\theta_8$, we compute that $c(\nu) = 36\|\theta_1\|^2$. Since $\nu > 0$, all terms in Freudenthal's formula are nonnegative. Consider the roots $\alpha = \frac{1}{2}(\theta_8 - \theta_7 + \sum_{i=1}^6 \epsilon_i \theta_i)$, $\beta = \frac{1}{2}(\theta_8 - \theta_7 - \sum_{i=1}^6 \epsilon_i \theta_i)$, where $\prod_{i=1}^6 \epsilon_i = -1$. Note that $\alpha + \beta = \theta_8 - \theta_7$. Since $(\lambda, \alpha) \neq 0$, we have that $\lambda - \alpha$ is a weight. Now we have two strings of weights starting at ν , namely

$$\begin{aligned}\nu, \quad \nu + (\theta_8 - \theta_7) = \lambda \quad \text{and} \\ \nu, \quad \nu + \beta = \lambda - \alpha.\end{aligned}$$

Note that in fact there are 32 different possible choices of the signs $\epsilon_i = \pm 1$, each of which gives rise to a different string of weights of the type ν , $\nu + \beta$. Freudenthal's formula gives

$$\begin{aligned}m_\nu &\geq \frac{2}{36\|\theta_1\|^2} ((\lambda, \theta_8 - \theta_7) m_\lambda + 32(\lambda - \alpha, \beta) m_{\lambda - \alpha}) \\ &= \frac{1}{18} (2 \cdot 1 + 32 \cdot 2 \cdot 1) = \frac{33}{9} > 3,\end{aligned}$$

so $m_\nu \geq 4$ and we are done. $\square$

5.37 Lemma *Representations numbers 25 and 31 are not of class $\mathcal{O}^2$.*

Proof. Here we will use the sharper version of condition (C_2) given by Lemma 4.8, item (a). These representations are just

$$\pi_\lambda : \Lambda^3(\overset{1}{\circ} - \cdots - \underset{\circ}{\circ} \begin{array}{l} \nearrow \circ \\ \searrow \circ \end{array}), \quad n \geq 4,$$

where $\overset{1}{\circ} - \cdots - \underset{\circ}{\circ} \begin{array}{l} \nearrow \circ \\ \searrow \circ \end{array}$ is the vector representation. The vector representation has weights $\pm\theta_i$, $1 \leq i \leq n$; let $\{e_i, e_{n+i} : 1 \leq i \leq n\}$ be a basis of weight vectors such that e_i (resp., e_{n+i}) corresponds to the θ_i -weight (resp., $-\theta_i$ -weight) weight and $\epsilon(e_i) = e_{n+i}$, where ϵ is a invariant real structure on the representation space.

We have that $v_\mu = e_1 \wedge e_3 \wedge e_{n+3}$ is a $(\mu = \theta_1)$ -weight vector for π_λ and $\epsilon(v_\mu) = e_{n+1} \wedge e_{n+3} \wedge e_3$. We want to show that the condition (a) of Lemma 4.8 is violated by proving that the θ_2 -weight vector $e_2 \wedge e_4 \wedge e_{n+4}$ is not in the complex span of

$$v_\mu, \quad \epsilon(v_\mu), \quad X_\alpha v_\mu, \quad X_\alpha \epsilon(v_\mu), \quad X_\beta X_\alpha(v_\mu), \quad X_\beta X_\alpha(\epsilon(v_\mu)),$$

for $\alpha, \beta \in \Delta$. But this follows immediately from the description of the action of a root vector X_α , $\alpha \in \Delta = \{\pm(\theta_i \pm \theta_j) : 1 \leq i < j \leq n\}$, on the weight vectors $\{e_i, e_{n+i} : 1 \leq i \leq n\}$ under the vector representation, given by the following matrices:

$$\begin{aligned} X_{\theta_i + \theta_j} &= E_{i,n+j} - E_{j,n+i}, \\ X_{-\theta_i - \theta_j} &= E_{n+i,j} - E_{n+j,i}, \\ X_{\theta_i - \theta_j} &= E_{i,j} - E_{n+j,n+i}, \end{aligned}$$

where E_{ij} denotes the $2n \times 2n$ -matrix with 1 at the (i, j) -position and 0 elsewhere. □

5.38 Lemma *Representation number 7 is not of class $\mathcal{O}^2$.*

Proof. We use an argument somewhat similar to the one used in the previous lemma. The representation is $\pi_\lambda : \Lambda^3(\overset{1}{\circ} - \overset{1}{\circ} \text{---} \bullet)$, where $\overset{1}{\circ} - \overset{1}{\circ} \text{---} \bullet$ is the vector representation. Let $e_1, e_2, e_3, e_4 = \epsilon(e_1), e_5 = \epsilon(e_2), e_6 = \epsilon(e_3), e_7 = \epsilon(e_4)$ be weight vectors of the vector representation corresponding to the weights $\theta_1, \theta_2, \theta_3, -\theta_1, -\theta_2, -\theta_3, 0$, respectively, where ϵ is the invariant real structure on the representation space.

We have that $v_\mu = e_1 \wedge e_4 \wedge e_7$ is a $(\mu = 0)$ -weight vector for π_λ and $\epsilon(v_\mu) = -v_\mu$. Now condition (a) of Lemma 4.8 is violated because the θ_2 -weight vector $e_2 \wedge e_3 \wedge e_6$ is not in the complex span of

$$v_\mu, \quad X_\alpha v_\mu, \quad X_\beta X_\alpha(v_\mu),$$

for $\alpha, \beta \in \Delta$. □

5.39 Lemma *Representation number 33 is not of class $\mathcal{O}^2$.*

Proof. We start with a description of the Lie algebra of $\mathbf{G}_2$ and of its 7-dimensional representation, as it is done in [18]. Let α_1, α_2 be the simple roots. We label the other positive roots as $\alpha_3 = \alpha_1 + \alpha_2$, $\alpha_4 = 2\alpha_1 + \alpha_2$, $\alpha_5 = 3\alpha_1 + \alpha_2$, $\alpha_6 = 3\alpha_1 + 2\alpha_2$. Choose root vectors X_i, Y_i for $\alpha_i, -\alpha_i$, respectively, $i = 1, 2$, such that $[H_i, X_i] = 2Y_i$, $[H_i, Y_i] = -2X_i$, where $H_i = [X_i, Y_i]$, $i = 1, 2$. Next define

$$\begin{aligned} X_3 &= [X_1, X_2], & Y_3 &= -[Y_1, Y_2], \\ X_4 &= \frac{1}{2}[X_1, X_3], & Y_4 &= -\frac{1}{2}[Y_1, Y_3], \\ X_5 &= -\frac{1}{3}[X_1, X_4], & Y_5 &= \frac{1}{3}[Y_1, Y_4], \\ X_6 &= -[X_2, X_5], & Y_6 &= [Y_2, Y_5], \end{aligned}$$

and $H_i = [X_i, Y_i]$ for $i = 1, \dots, 6$. Then $[H_i, X_i] = 2Y_i$, $[H_i, Y_i] = -2X_i$ for $i = 3, \dots, 6$, and X_i, Y_i are root vectors for $\alpha_i, -\alpha_i$, respectively, $i = 1, \dots, 6$.

The 7-dimensional representation of $\mathbf{G}_2$ has weights $\alpha_1, -\alpha_1, \alpha_3, -\alpha_3, \alpha_4, -\alpha_4, 0$ with respective weight vectors $v_1, w_1, v_3, w_3, v_4, w_4, u$, such that the action of the basis vectors of the Lie algebra of $\mathbf{G}_2$ is described in the following table:

	H_1	H_2	X_1	X_2	X_3	X_4	X_5	X_6	Y_1	Y_2	Y_3	Y_4	Y_5	Y_6
u	0	0	$2v_1$	0	$2v_3$	$2v_4$	0	0	$2w_1$	0	$2w_3$	$2w_4$	0	0
v_4	v_4	0	0	0	0	0	0	0	v_3	0	$-v_1$	u	w_1	w_3
v_1	$2v_1$	$-v_1$	0	$-v_3$	$-v_4$	0	0	0	u	0	0	w_3	$-w_4$	0
v_3	$-v_3$	v_3	v_4	0	0	0	0	0	0	$-v_1$	u	$-w_1$	0	$-w_4$
w_4	$-w_4$	0	$-w_3$	0	w_1	u	$-v_1$	$-v_3$	0	0	0	0	0	0
w_1	$-2w_1$	w_1	u	0	0	$-v_3$	v_4	0	0	w_3	w_4	0	0	0
w_3	w_3	$-w_3$	0	w_1	u	v_1	0	v_4	$-w_4$	0	0	0	0	0

Now we have the equation

$$S^2(\overset{1}{\bullet \equiv \circ}) = \overset{2}{\bullet \equiv \circ} \oplus (\text{trivial}),$$

from which we learn three things. First, $\pi_\lambda : \overset{2}{\bullet \equiv \circ}$ is a subrepresentation of the symmetric square of the 7-dimensional representation, so we can read it off the table above. Second, the multiplicity of the weights of π_λ ; in particular, α_4 is a weight of multiplicity 2. And third, the trivial summand above is spanned by $-\frac{1}{2}u^2 + 2v_1w_1 + 2v_3w_3 + 2v_4w_4$, so that an invariant real structure on the representation space of the 7-dimensional representation is given by $\epsilon(v_i) = w_i$, $i = 1, 2, 3$ and $\epsilon(u) = -u$.

Let $v_\mu = 2u^2 - 8v_1w_1 + 6v_3w_3 + 6v_4w_4$, which is a real, $(\mu = 0)$ -weight vector for π_λ . We next show that condition (a) of Lemma 4.8 is violated because the α_4 -weight space is not contained in the complex span of

$$(5.40) \quad v_\mu, \quad X_\alpha v_\mu, \quad X_\beta X_\alpha(v_\mu),$$

for $\alpha, \beta \in \Delta$. In fact, $\alpha_4 - 0$ can be written as a root or as a sum of two roots as follows:

$$\begin{aligned}
\alpha_4 - 0 &= 2\alpha_1 + \alpha_2 \\
&= \alpha_1 + (\alpha_1 + \alpha_2) \\
&= -\alpha_1 + (3\alpha_1 + \alpha_2) \\
&= (3\alpha_1 + 2\alpha_2) - (\alpha_1 + \alpha_2)
\end{aligned}$$

Now applying the corresponding root vectors to v_μ we get:

$$\begin{aligned}
X_4(v_\mu) &= 14(uv_4 + v_1v_3), \\
X_3X_1(v_\mu) &= X_3(0) = 0, \\
X_5Y_1(v_\mu) &= X_5(0) = 0, \\
Y_3X_6(v_\mu) &= Y_3(0) = 0.
\end{aligned}$$

This shows that the intersection of the complex span of (5.40) with the α_4 -weight space has dimension 1; but the weight space itself has dimension 2. $\square$

Proof of Lemma 5.16. We eliminate the possibilities given by Table B.1 and not listed in the table.

Let $G = \mathbf{C}_n$, $\pi_\lambda : \overset{3}{\bullet} \text{---} \bullet \text{---} \cdots \text{---} \bullet \text{---} \bullet \text{---} \circ$, $n \geq 2$. Then $\lambda = 3\theta_1$, $\mu = -3\theta_2$ is a weight and $\lambda \pm \mu$ are neither roots nor a sum of two roots, so π_λ does not even satisfy (C_2) .

Let $G = \mathbf{C}_n$, $\pi_\lambda : \overset{1}{\bullet} \text{---} \overset{1}{\bullet} \text{---} \cdots \text{---} \bullet \text{---} \bullet \text{---} \circ$, $n \geq 3$. We have

$$\begin{aligned}
&\overset{1}{\bullet} \text{---} \overset{1}{\bullet} \text{---} \cdots \text{---} \bullet \text{---} \bullet \text{---} \circ \otimes \overset{1}{\bullet} \text{---} \bullet \text{---} \cdots \text{---} \bullet \text{---} \bullet \text{---} \circ = \\
&\quad \overset{1}{\bullet} \text{---} \overset{1}{\bullet} \text{---} \cdots \text{---} \bullet \text{---} \bullet \text{---} \circ \oplus \overset{1}{\bullet} \text{---} \bullet \text{---} \cdots \text{---} \bullet \text{---} \bullet \text{---} \circ \oplus \overset{1}{\bullet} \text{---} \bullet \text{---} \cdots \text{---} \bullet \text{---} \bullet \text{---} \circ,
\end{aligned}$$

from where we deduce that the multiplicity of $\mu = \theta_3$ as a weight of $\overset{1}{\bullet} \text{---} \overset{1}{\bullet} \text{---} \cdots \text{---} \bullet \text{---} \bullet \text{---} \circ$ is $2n - 2 \geq 4$. Now $\lambda - \mu$ is not a root and can be written as a sum of two roots in two ways,

$$\begin{aligned}
\lambda - \mu &= (2\theta_1) + (\theta_2 - \theta_3) \\
&= (\theta_1 + \theta_2) + (\theta_1 - \theta_3),
\end{aligned}$$

and $\lambda + \mu = 2\theta_1 + \theta_2 + \theta_3$ is not a root. It follows that the dimension of the intersection of the μ -weight space with $\mathcal{U}^2(\mathfrak{g}^c)v_\lambda + \mathcal{U}^1(\mathfrak{g}^c)\epsilon(v_\lambda)$ is at most 2, so that condition $(C_{1\frac{1}{2}})$ is not satisfied.

Let $G = \mathbf{C}_n$, $\pi_\lambda : \overset{1}{\bullet} \text{---} \bullet \text{---} \bullet \text{---} \cdots \text{---} \bullet \text{---} \bullet \text{---} \circ$, $n \geq 4$. Then $\lambda = \theta_1 + \theta_2 + \theta_3$ and this representation is a subrepresentation of the cubic exterior power of the vector representation. Also, $\mu = -\theta_1$ is a weight of multiplicity $n - 2 \geq 2$. But one can check by direct computation that the dimension of the intersection of the μ -weight space with $\mathcal{U}^2(\mathfrak{g}^c)v_\lambda + \mathcal{U}^1(\mathfrak{g}^c)\epsilon(v_\lambda)$ is 1, so condition $(C_{1\frac{1}{2}})$ is not satisfied. $\square$

6 The proof of the converse to a Theorem of Bott and Samelson

In this section we classify variationally complete representations. The strategy is as follows. We first exclude many irreducible representations in class $\mathcal{O}^2$ from being variationally complete by making use of Propositions 3.7 and 3.8. We then observe that the remaining irreducible representations are all orbit equivalent to isotropy representations of symmetric spaces, hence variationally complete by Proposition 3.3. Using Proposition 3.4, item (d), we see that arbitrary variationally complete representations are precisely those that are orbit equivalent to isotropy representations of symmetric spaces. This result can be viewed as a converse to Theorem 2.2 of Bott and Samelson. Finally we classify arbitrary variationally complete representations, mainly as a repeated application of Proposition 3.4, item (b).

Recall that the irreducible representations which according to the results in Section 5 can belong to class $\mathcal{O}^2$ without being either the isotropy representation of a symmetric space or of cohomogeneity one are listed in the tables of Propositions 5.1, 5.2, 5.17, 5.18, 5.19, 5.20 and 5.21. For the sake of our forthcoming arguments, these representations are rearranged in a more systematic way in Tables C.1, C.2, C.3 and C.4 in Appendix C and in the table of Proposition 6.3.

6.1 Proposition *The representations listed in Tables C.1, C.2 and C.3 are not variationally complete.*

Proof. For each representation ρ of the compact connected Lie group G on the real vector space V listed in these tables, we need to exhibit a compact symmetric space $X = L/K$ such that:

- (a) K contains G as a closed subgroup;
- (b) the isotropy representation $\tilde{\rho}$ of X restricts to ρ on G ;
- (c) there exists $p \in V$ such that the orbits of $\tilde{\rho}$ and ρ through p are the equal;
- (d) there exists $q \in V$ such that the orbits of $\tilde{\rho}$ and ρ through q are different.

It will then follow from Proposition 3.7 that ρ is not variationally complete.

Let $G = \mathbf{SO}(2) \times \mathbf{Spin}(9)$. Then $X = \mathbf{SO}(18)/\mathbf{SO}(2) \times \mathbf{SO}(16)$, $\tilde{\rho}$ is the tensor product of the vector representations of $\mathbf{SO}(2)$, $\mathbf{SO}(16)$ on $\mathbf{R}^2$, $\mathbf{R}^{16}$, respectively, and we take $p = v_1 \otimes v_2 \in \mathbf{R}^2 \otimes \mathbf{R}^{16}$. Note that the orbits of $\tilde{\rho}$ and ρ through p are equal because $\mathbf{SO}(2)$, $\mathbf{Spin}(9)$ and $\mathbf{SO}(16)$ are all transitive on unit spheres, but the orbits through a common regular point q are different because $\tilde{\rho}$ is of cohomogeneity two and ρ is of cohomogeneity three.

Let $G = \mathbf{U}(2) \times \mathbf{Sp}(n)$, $n \geq 2$. Then $X = \mathbf{SU}(2 + 2n)/\mathbf{S}(\mathbf{U}(2) \times \mathbf{U}(2n))$, $\tilde{\rho}$ can be viewed as the realification of the complex tensor product of the vector representations of $\mathbf{U}(2)$ and $\mathbf{SU}(2n)$ on $\mathbf{C}^2$, $\mathbf{C}^{2n}$, respectively, and we take $p = v_1 \otimes v_2 \in \mathbf{C}^2 \otimes \mathbf{C}^{2n}$. The orbits of $\tilde{\rho}$ and ρ through p are equal because $\mathbf{U}(2)$, $\mathbf{Sp}(n)$ and $\mathbf{SU}(2n)$ are all transitive on unit spheres, but the orbits through a common regular point q are different because $\tilde{\rho}$ is of cohomogeneity two and ρ is of cohomogeneity three.

Let $G = \mathbf{SU}(2) \times \mathbf{Sp}(n)$, $n \geq 2$. Then $X = \mathbf{Sp}(2+n)/\mathbf{Sp}(2) \times \mathbf{Sp}(n)$, $\tilde{\rho}$ is the quaternionic tensor product of the vector representations of $\mathbf{Sp}(2)$, $\mathbf{Sp}(n)$ on $\mathbf{H}^2$, $\mathbf{H}^n$, respectively, and we take $p = e_1 \otimes f_1 + e_2 \otimes f_2 \in \mathbf{H}^2 \otimes_{\mathbf{H}} \mathbf{H}^n$, where $\{e_1, e_2\} \subset \mathbf{H}^2$ and $\{f_1, f_2\} \subset \mathbf{H}^n$ are orthonormal. The orbits of $\tilde{\rho}$ and ρ through p are equal because both of them are $8n - 6$ -dimensional, but the orbits through a common regular point q are different because $\tilde{\rho}$ is of cohomogeneity two and ρ is of cohomogeneity three.

We omit the details for the representations of Tables C.2 and C.3 because they are analogous to the above (For instance, the reason for the existence of the common orbit (item (c)) is that as in the first two cases above we are dealing with tensor products of representations transitive on unit spheres.) Moreover, we will prove something stronger later in Propositions 7.5 and 7.6, namely that these representations are not taut. $\square$

6.2 Proposition *The representations listed in Table C.4 are not taut.*

Proof. Let π_λ be a representation of the compact connected Lie group G on V_λ which is in the table. Notice that π_λ is of real type. Let ρ be a real form acting on $V \subset V_\lambda$ and let ϵ be the invariant real structure on V_λ . We shall go case by case and show that the slice representation of ρ at a specific point $p \in V$ is not taut, and then use Proposition 3.8 to conclude that ρ is not taut. In fact, we use $p = v_\lambda + \epsilon(v_\lambda) \in V$ where v_λ is a highest weight vector for π_λ . In the following, we omit the tedious and sometimes lengthy calculations involved and simply state the final results.

Let $G = \mathbf{Sp}(n) \times \mathbf{SU}(6)$. Then the isotropy subalgebra at p is

$$\mathfrak{u}(1) + \mathfrak{su}(3) + \mathfrak{su}(3) + \mathfrak{sp}(n-1), \quad n \geq 2$$

The dimension of the orbit through p is $\dim G(p) = 4n + 17$, the dimension of the normal space at p is $36n - 17$, and the complexified slice representation at p is given by:

$$\begin{aligned} & (\text{trivial}) \\ & \oplus (x^4 \otimes \overset{1}{\circ} - \overset{1}{\circ} \otimes \overset{1}{\circ} - \overset{1}{\circ} \otimes \overset{1}{\circ}) \oplus (x^{-4} \otimes \overset{1}{\circ} - \overset{1}{\circ} \otimes \overset{1}{\circ} - \overset{1}{\circ} \otimes \overset{1}{\circ}) \otimes (\text{trivial}) \\ & \oplus (x \otimes \overset{1}{\circ} - \overset{1}{\circ} \otimes \overset{1}{\circ} - \overset{1}{\circ} \otimes \overset{1}{\circ}) \oplus (x^{-1} \otimes \overset{1}{\circ} - \overset{1}{\circ} \otimes \overset{1}{\circ} - \overset{1}{\circ} \otimes \overset{1}{\circ}) \otimes \bullet - \dots - \bullet = \bullet. \end{aligned}$$

Therefore, the slice representation contains as a summand the realification of

$$x \otimes \overset{1}{\circ} - \overset{1}{\circ} \otimes \overset{1}{\circ} - \overset{1}{\circ} \otimes \bullet - \dots - \bullet = \bullet.$$

But this is a representation of complex type with Dadok invariant 3. By Proposition 4.9, it cannot be of class $\mathcal{O}^2$. In particular, it is not taut.

Let $G = \mathbf{Sp}(n) \times \mathbf{Spin}(12)$. Then the isotropy subalgebra at p is

$$\mathfrak{u}(1) + \mathfrak{su}(6) + \mathfrak{sp}(n-1), \quad n \geq 2$$

The dimension of the orbit through p is $\dim G(p) = 4n + 29$, the dimension of the normal space at p is $60n - 29$, and the complexified slice representation at p is given by:

$$\begin{aligned} & (\text{trivial}) \\ & \oplus (x^4 \otimes \overset{1}{\circ} - \overset{1}{\circ} \otimes \overset{1}{\circ} - \overset{1}{\circ} \otimes \overset{1}{\circ} - \overset{1}{\circ} \otimes \overset{1}{\circ}) \oplus (x^{-4} \otimes \overset{1}{\circ} - \overset{1}{\circ} \otimes \overset{1}{\circ} - \overset{1}{\circ} \otimes \overset{1}{\circ} - \overset{1}{\circ} \otimes \overset{1}{\circ}) \otimes (\text{trivial}) \\ & \oplus (x \otimes \overset{1}{\circ} - \overset{1}{\circ} \otimes \overset{1}{\circ} - \overset{1}{\circ} \otimes \overset{1}{\circ} - \overset{1}{\circ} \otimes \overset{1}{\circ}) \oplus (x^{-1} \otimes \overset{1}{\circ} - \overset{1}{\circ} \otimes \overset{1}{\circ} - \overset{1}{\circ} \otimes \overset{1}{\circ} - \overset{1}{\circ} \otimes \overset{1}{\circ}) \otimes \bullet - \dots - \bullet = \bullet. \end{aligned}$$

Therefore, the slice representation contains as a summand the realification of

$$x \otimes \overset{1}{\circ} - \overset{1}{\circ} - \overset{1}{\circ} - \overset{1}{\circ} - \overset{1}{\circ} \otimes \bullet \cdots \bullet = \bullet.$$

But this is a representation of complex type with Dadok invariant 3. By Proposition 4.9, it cannot be of class $\mathcal{O}^2$. In particular, it is not taut.

Let $G = \mathbf{Sp}(n) \times \mathbf{Sp}(3)$. Then the isotropy subalgebra at p is

$$\mathfrak{u}(1) + \mathfrak{su}(3) + \mathfrak{sp}(n-1), \quad n \geq 2$$

The dimension of the orbit through p is $\dim G(p) = 4n + 11$, the dimension of the normal space at p is $24n - 11$, and the complexified slice representation at p is given by:

$$\begin{aligned} & (\text{trivial}) \\ & \oplus (x^4 \otimes \overset{2}{\circ} - \overset{2}{\circ} \oplus x^{-4} \otimes \overset{2}{\circ} - \overset{2}{\circ}) \otimes (\text{trivial}) \\ & \oplus (x \otimes \overset{2}{\circ} - \overset{2}{\circ} \oplus x^{-1} \otimes \overset{2}{\circ} - \overset{2}{\circ}) \otimes \overset{1}{\bullet} \cdots \bullet = \bullet. \end{aligned}$$

Therefore, the slice representation contains as a summand the realification of

$$x \otimes \overset{2}{\circ} - \overset{2}{\circ} \otimes \overset{1}{\bullet} \cdots \bullet = \bullet.$$

But this is a representation of complex type with Dadok invariant 3. By Proposition 4.9, it cannot be of class $\mathcal{O}^2$. In particular, it is not taut.

Let $G = \mathbf{Sp}(n) \times \mathbf{E}_7$. Then the isotropy subalgebra at p is

$$\mathfrak{u}(1) + \mathfrak{e}_6 + \mathfrak{sp}(n-1), \quad n \geq 2$$

The dimension of the orbit through p is $\dim G(p) = 4n + 53$, the dimension of the normal space at p is $108n - 53$, and the complexified slice representation at p is given by:

$$\begin{aligned} & (\text{trivial}) \\ & \oplus (x^4 \otimes \overset{1}{\circ} - \overset{1}{\circ} - \overset{1}{\circ} - \overset{1}{\circ} - \overset{1}{\circ} \oplus x^{-4} \otimes \overset{1}{\circ} - \overset{1}{\circ} - \overset{1}{\circ} - \overset{1}{\circ} - \overset{1}{\circ}) \otimes (\text{trivial}) \\ & \oplus (x \otimes \overset{1}{\circ} - \overset{1}{\circ} - \overset{1}{\circ} - \overset{1}{\circ} - \overset{1}{\circ} \oplus x^{-1} \otimes \overset{1}{\circ} - \overset{1}{\circ} - \overset{1}{\circ} - \overset{1}{\circ} - \overset{1}{\circ}) \otimes \overset{1}{\bullet} \cdots \bullet = \bullet. \end{aligned}$$

Therefore, the slice representation contains as a summand the realification of

$$x \otimes \overset{1}{\circ} - \overset{1}{\circ} - \overset{1}{\circ} - \overset{1}{\circ} - \overset{1}{\circ} \otimes \overset{1}{\bullet} \cdots \bullet = \bullet.$$

But this is a representation of complex type with Dadok invariant 3. By Proposition 4.9, it cannot be of class $\mathcal{O}^2$. In particular, it is not taut.

Let $G = \mathbf{Sp}(1) \times \mathbf{Spin}(13)$. Then the isotropy subalgebra at p is

$$\mathfrak{u}(1) + \mathfrak{su}(6).$$

The dimension of the orbit through p is $\dim G(p) = 45$, the dimension of the normal space at p is 83, and the complexified slice representation at p is given by:

$$\begin{aligned} & \text{(trivial)} \\ & \oplus (x^5 \otimes \overset{1}{\circ} \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \oplus x^{-5} \otimes \circ \text{---} \circ \text{---} \circ \text{---} \circ \overset{1}{\circ}) \\ & \oplus (x^4 \otimes \overset{1}{\circ} \text{---} \circ \text{---} \circ \text{---} \circ \oplus x^{-4} \otimes \circ \text{---} \circ \text{---} \circ \text{---} \circ \overset{1}{\circ}) \\ & \oplus (x^3 \oplus x^{-3}) \otimes \circ \text{---} \overset{1}{\circ} \text{---} \circ \text{---} \circ. \end{aligned}$$

Therefore, the slice representation contains as a summand the realification of

$$x^3 \otimes \circ \text{---} \overset{1}{\circ} \text{---} \circ \text{---} \circ.$$

But this is a representation of complex type with Dadok invariant 3. By Proposition 4.9, it cannot be of class $\mathcal{O}^2$. In particular, it is not taut.

Let $G = \mathbf{Sp}(1) \times \mathbf{Sp}(2)$. Then the isotropy subalgebra at p is

$$\mathfrak{u}(1) + \mathfrak{u}(1).$$

The dimension of the orbit through p is $\dim G(p) = 11$, the dimension of the normal space at p is 21, and the slice representation at p is given by the realification of:

$$\begin{aligned} & \text{(trivial)} \\ & \oplus (x^{-5} \oplus x^{-2} \oplus x^1) \otimes x^2 \\ & \oplus (x^5 \oplus x^{-4} \oplus 2x^{-1} \oplus 2x^2) \otimes x^4 \\ & \oplus x^3 \otimes x^6. \end{aligned}$$

Therefore, the slice representation contains as a summand the realification of $x^{-2} \otimes x^2 \oplus x^{-4} \otimes x^4$ which is orbit equivalent to the circle action

$$x^2 \oplus x^4$$

on $\mathbf{R}^2 \oplus \mathbf{R}^2$. But the orbit of the circle action through $((1, 0), (0, 1))$ is a torus-knot which is not taut.

Let $G = \mathbf{Spin}(15)$. Then the isotropy subalgebra at p is $\mathfrak{su}(7)$. The dimension of the orbit through p is $\dim G(p) = 57$, the dimension of the normal space at p is 71, and the complexified slice representation at p is given by:

$$\text{(trivial)} \oplus \circ \text{---} \overset{1}{\circ} \text{---} \circ \text{---} \circ \text{---} \circ \oplus \circ \text{---} \overset{1}{\circ} \text{---} \circ \text{---} \circ$$

Therefore, the slice representation contains as a summand the realification of

$$\circ \text{---} \overset{1}{\circ} \text{---} \circ \text{---} \circ$$

But this is a representation of complex type with Dadok invariant 3. By Proposition 4.9, it cannot be of class $\mathcal{O}^2$. In particular, it is not taut.

Let $G = \mathbf{Spin}(17)$. Then the isotropy subalgebra at p is $\mathfrak{su}(8)$. The dimension of the orbit through p is $\dim G(p) = 73$, the dimension of the normal space at p is 183, and the complexified slice representation at p is given by:

$$\begin{array}{c} \text{(trivial)} \oplus \text{---} \overset{1}{\circ} \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \\ \oplus \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \overset{1}{\circ} \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \oplus \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \overset{1}{\circ} \text{---} \circ \text{---} \circ \end{array}$$

Therefore, the slice representation contains as a summand the realification of

$$\text{---} \circ \text{---} \circ \text{---} \overset{1}{\circ} \text{---} \circ \text{---} \circ \text{---} \circ$$

But this is a representation of complex type with Dadok invariant 3. By Proposition 4.9, it cannot be of class $\mathcal{O}^2$. In particular, it is not taut.

Let $G = \mathbf{Spin}(7)$. Then the isotropy subalgebra at p is

$$\mathfrak{u}(1) + \mathfrak{su}(2)$$

The dimension of the orbit through p is $\dim G(p) = 17$, the dimension of the normal space at p is 31, and the slice representation at p is given by:

$$\begin{array}{l} \text{(trivial)} \\ \oplus (2(x^4 \oplus x^{-4}) \oplus (x^2 \oplus x^{-2})) \otimes \text{(trivial)} \\ \oplus (2(x^1 \oplus x^{-1}) \oplus (x^7 \oplus x^{-7})) \otimes \overset{1}{\circ} \\ \oplus ((x^2 \oplus x^{-2}) \oplus (x^4 \oplus x^{-4})) \otimes \overset{2}{\circ} \end{array}$$

Therefore, the slice representation contains as a summand the realification of

$$x^4 \oplus x^2$$

acting on $\mathbf{R}^2 \oplus \mathbf{R}^2$. But the orbit through $((1, 0), (0, 1))$ is a torus-knot which is not taut.

Let $G = \mathbf{Spin}(9)$. Then the isotropy subalgebra at p is

$$\mathfrak{u}(1) + \mathfrak{su}(3)$$

The dimension of the orbit through p is $\dim G(p) = 27$, the dimension of the normal space at p is 101, and the complexified slice representation at p is given by:

$$\begin{array}{l} \text{(trivial)} \\ \oplus ((x^3 \oplus x^{-3}) \oplus 2(x^2 \oplus x^{-2}) \oplus (x \oplus x^{-1})) \otimes \text{(trivial)} \\ \oplus ((x \oplus x^{-1}) \oplus (x^2 \oplus x^{-2})) \otimes \overset{1}{\circ} \text{---} \overset{1}{\circ} \\ \oplus ((3x \oplus 2x^0 \oplus x^{-3}) \otimes \overset{1}{\circ} \text{---} \overset{1}{\circ} \oplus (3x^{-1} \oplus 2x^0 \oplus x^3) \otimes \overset{1}{\circ} \text{---} \overset{1}{\circ}) \\ \oplus ((x^0 \oplus x^{-1}) \otimes \overset{2}{\circ} \text{---} \overset{2}{\circ} \oplus (x^0 \oplus x^1) \otimes \overset{2}{\circ} \text{---} \overset{2}{\circ}) \end{array}$$

$$x \otimes \overset{1}{\circ} \overset{1}{\circ}$$

Let $G = \mathbf{Sp}(n) \times \mathbf{Spin}(11)$. Then the isotropy subalgebra at p is

$$\mathfrak{u}(1) + \mathfrak{su}(5) + \mathfrak{sp}(n-1), \quad n \geq 2$$

(trivial)

[illegible]





For $n = 1$ we need a different argument. In this case the connected component K of the isotropy group at p is locally isomorphic to $\mathbf{U}(5)$, and the slice representation at p contains as a summand the realification of

$$x^8 \otimes \overset{1}{\bigcirc} \text{---} \bigcirc \text{---} \bigcirc \text{---} \bigcirc \oplus x^6 \otimes \overset{1}{\bigcirc} \text{---} \bigcirc \text{---} \bigcirc \text{---} \bigcirc.$$

Denote by V_1, V_2 the respective representation spaces of the above summands. Then $V_1 = \mathbf{C}^5$ and $V_2 = \Lambda^2 \mathbf{C}^5$. Choose an orthonormal basis $\{e_1, \dots, e_5\}$ for $\mathbf{C}^5$. Let $q_2 = ae_1 \wedge e_2 + be_3 \wedge e_4 \in V_2$ be a regular point, where a, b are distinct positive real numbers. We have that the isotropy K_{q_2} is locally isomorphic to $\mathbf{SU}(2) \times \mathbf{SU}(2) \times \mathbf{U}(1)$ sitting diagonally in K . Let $q_1 = e_1 + e_3 \in V_1$. Now the orbit $K_{q_2}(q_1)$ is diffeomorphic to $S^3 \times S^3$, whereas the orbit $K(q_1, q_2)$ is diffeomorphic to $\mathbf{SU}(5)$. It is known that the third Betti number of a compact connected simple Lie group is one. It follows from Proposition 3.11 that the slice representation at p is not taut. $\square$

6.3 Proposition *The following is a list of pairs of irreducible representations ρ, ρ' of compact connected Lie groups G, G' such that ρ' is the isotropy representation of a symmetric*

space, G is a subgroup of G' and $\rho = \rho'|_G$ is orbit equivalent to ρ' .

G and G'	ρ and ρ'
$\mathbf{SU}(n) \times \mathbf{SU}(m), n \neq m$ and $S^1 \times \mathbf{SU}(n) \times \mathbf{SU}(m)$	$(\bigcirc - \dots - \overset{1}{\bigcirc} \otimes \overset{1}{\bigcirc} - \dots - \bigcirc)$ $\oplus (\bigcirc - \dots - \overset{1}{\bigcirc} \otimes \bigcirc - \dots - \overset{1}{\bigcirc})$ and $(\bigcirc - \dots - \overset{1}{\bigcirc} \otimes x \otimes \overset{1}{\bigcirc} - \dots - \bigcirc)$ $\oplus (\bigcirc - \dots - \overset{1}{\bigcirc} \otimes x^{-1} \otimes \bigcirc - \dots - \overset{1}{\bigcirc})$
$\mathbf{SU}(n), n \text{ odd, and}$ $\mathbf{U}(n)$	$\bigcirc - \overset{1}{\bigcirc} - \dots - \bigcirc \oplus \bigcirc - \dots - \overset{1}{\bigcirc} - \bigcirc$ and $(x \otimes \bigcirc - \overset{1}{\bigcirc} - \dots - \bigcirc) \oplus (x^{-1} \otimes \bigcirc - \dots - \overset{1}{\bigcirc} - \bigcirc)$
$\mathbf{Spin}(10)$ and $\mathbf{SO}(2) \times \mathbf{Spin}(10)$	 and 
$\mathbf{SO}(2) \times \mathbf{Spin}(7)$ and $\mathbf{SO}(2) \times \mathbf{SO}(8)$	$(x \oplus x^{-1}) \otimes \bigcirc - \overset{1}{\bullet} - \bigcirc$ and 
$\mathbf{SO}(2) \times \mathbf{G}_2$ and $\mathbf{SO}(2) \times \mathbf{SO}(7)$	$(x \oplus x^{-1}) \otimes \bullet - \overset{1}{\bullet} - \bigcirc$ and $(x \oplus x^{-1}) \otimes \bigcirc - \overset{1}{\bullet} - \bullet$
$\mathbf{SO}(3) \times \mathbf{Spin}(7)$ and $\mathbf{SO}(3) \times \mathbf{SO}(8)$	$\overset{2}{\bigcirc} \otimes \bigcirc - \overset{1}{\bullet} - \bullet$ and 

Proof. The first three cases in the table were treated in Lemma 5.4, item (a). It remains to analyze the other three cases. It suffices to show the existence of a common principal orbit by Lemma 3.6.

Let $G = \mathbf{SO}(2) \times \mathbf{Spin}(7)$. Then ρ is the tensor product of the vector representation of $\mathbf{SO}(2)$ on $\mathbf{R}^2$ and the spin representation of $\mathbf{Spin}(7)$ on $\mathbf{R}^8$. Let $p = a_1 e_1 \otimes f_1 + a_2 e_2 \otimes f_2 \in \mathbf{R}^2 \otimes \mathbf{R}^8$, where $e_1, e_2 \in \mathbf{R}^2$ are orthonormal, $f_1, f_2 \in \mathbf{R}^8$ are orthonormal, and a_1, a_2 are distinct positive numbers. Then the orbit $G'p$ is principal and of dimension 14. Since the isotropy subalgebra of G at p is easily seen to be $\mathfrak{su}(3)$, it follows that Gp is also an orbit of dimension 14. Hence, it coincides with $G'p$. It follows from Lemma 3.6 that ρ and ρ' are orbit equivalent.

Similarly, one sees that in the last two cases in the table ρ has respectively an orbit of dimension 12, 21, and therefore is orbit equivalent to ρ' . $\square$

6.4 Theorem (Converse to a Theorem of Bott and Samelson) *A variationally complete representation of a compact connected Lie group is orbit equivalent to the isotropy representation of a symmetric space.*

Proof. We first assume that the given representation is irreducible. A variationally complete, irreducible representation ρ of a compact connected Lie group G is of class $\mathcal{O}^2$. If ρ is not the isotropy representation of one of the irreducible symmetric spaces and has cohomogeneity greater than one, we have obtained all possibilities for it in Propositions 5.1, 5.2, 5.17, 5.18, 5.19, 5.20 and 5.21, and these are listed in the table of Proposition 6.3 and in Tables C.1, C.2, C.3 and C.4 in Appendix C. Now Propositions 6.1 and 6.2 eliminate all possibilities for ρ , except those listed in the table of Proposition 6.3. But the same proposition asserts that ρ is orbit equivalent to another representation ρ' which is the isotropy representation of a symmetric space.

Now assume that the representation is reducible. Each of its irreducible factors is variationally complete by part (a) of Proposition 3.4 and hence orbit equivalent to the isotropy representation of an irreducible symmetric space by what we have just proved. Part (d) of the same proposition now implies that the reducible representation is orbit equivalent to the isotropy representation of the symmetric space that is the product of the irreducible symmetric spaces corresponding to the irreducible summands of the representation. $\square$

6.5 Corollary *A variationally complete representation of a compact connected Lie group is polar.*

The first paragraph in the proof of Theorem 6.4 combined with Theorem 8.2 to be proved in Section 8 also implies:

6.6 Theorem (Compare [15]) *Let $\rho : G \rightarrow \mathbf{O}(V)$ be a variationally complete irreducible representation of a compact connected Lie group G . Then there are two cases:*

- (a) *The image $\rho(G)$ is the maximal compact connected Lie group with its orbits. In this case ρ is the isotropy representation of an irreducible symmetric space.*
- (b) *The image $\rho(G)$ is not the maximal compact connected Lie group with its orbits. In this case we have:*
 - (i) *If ρ has cohomogeneity greater than one, then there exists a larger group $G' \supset G$ and an isotropy representation of an irreducible symmetric space $\rho' : G' \rightarrow \mathbf{O}(V)$ which is an extension of ρ . Moreover, the possibilities for the pair (ρ, ρ') are listed in the table of Proposition 6.3.*
 - (ii) *If ρ has cohomogeneity one, then it is listed in the tables of Theorem 8.2.*

In order to complete the classification of variationally complete representations, we finally discuss the reducible case and state a result in Theorem 6.8 which, in view of Theorem 6.4, is equivalent to a result of Bergmann in [1]. First note that there are some trivial ways to produce variationally complete reducible representations from known variationally complete representations.

Let $\rho : G \rightarrow \mathbf{O}(V)$ be a representation of a compact connected Lie group G .

- (a) Assume $G = G' \times K$ where G', K are compact connected Lie groups. We say that K is *nonessential* for ρ if the restriction $\rho|_{G'}$ is orbit equivalent to the original representation ρ . Otherwise, we say that K is *essential* for ρ .
- (b) Assume ρ reducible, namely $\rho = \rho_1 \oplus \rho_2$ where $\rho_i : G \rightarrow \mathbf{O}(V_i)$ and $\dim V_i > 0$ ($i = 1, 2$). We say that ρ is *splitting* if we can write $G = G_1 \times G_2$ for compact connected Lie groups G_i ($i = 1, 2$) such that ρ is equivalent to the outer direct sum representation $\rho_1|_{G_1} \hat{\oplus} \rho_2|_{G_2} : G_1 \times G_2 \rightarrow \mathbf{O}(V_1 \oplus V_2)$. We say that ρ is *almost splitting* if we can write $G = G_1 \times G_2 \times K$ for compact connected Lie groups K, G_i ($i = 1, 2$) such that K is nonessential at least for one of $\rho_1|_{G_1 \times K}, \rho_2|_{G_2 \times K}$ and $\rho|_{G_1 \times G_2}$ is equivalent to the outer direct sum representation $\rho_1|_{G_1} \hat{\oplus} \rho_2|_{G_2}$.

It is immediate from these definitions that a representation with a trivial summand is automatically splitting, and that splitting implies almost splitting. We also have:

6.7 Proposition *An almost splitting representation $\rho = \rho_1 \oplus \rho_2$ is variationally complete if and only if each one of its summands ρ_i is variationally complete⁵.*

Proof. One direction follows from Proposition 3.4, item (a); we prove that the summands being variationally complete implies the same for the sum. We may assume that K is nonessential for $\rho_1|_{G_1 \times K}$. Then ρ_1 is orbit equivalent to $\rho_1|_{G_1}$ and ρ_2 is orbit equivalent to $\rho_2|_{G_2 \times K}$, so $\rho_1|_{G_1}$ and $\rho_2|_{G_2 \times K}$ are variationally complete. Now ρ is orbit equivalent to $\rho_1|_{G_1} \hat{\oplus} \rho_2|_{G_2 \times K}$ and hence variationally complete, because the outer direct sum of two variationally complete representations is also variationally complete, as is easy to see. $\square$

We are left with the classification of variationally complete reducible representations which are not almost splitting. This is given in the following theorem.

6.8 Theorem (Compare [1]) *Let ρ be a variationally complete reducible representation of a compact connected Lie group G which is not almost splitting. Then ρ is one of the following*

⁵Note that the isotropy representations of reducible symmetric spaces are splitting.

orthogonal representations:

G	ρ
SU(4)	$(\overset{1}{\circ} - \overset{1}{\circ} - \overset{1}{\circ} \oplus \overset{1}{\circ} - \overset{1}{\circ} - \overset{1}{\circ}) \oplus \overset{1}{\circ} - \overset{1}{\circ} - \overset{1}{\circ}$
U(4)	$(x \otimes \overset{1}{\circ} - \overset{1}{\circ} - \overset{1}{\circ} \oplus x^{-1} \otimes \overset{1}{\circ} - \overset{1}{\circ} - \overset{1}{\circ}) \oplus \overset{1}{\circ} - \overset{1}{\circ} - \overset{1}{\circ}$
Spin(7)	$\overset{1}{\circ} - \overset{1}{\circ} - \bullet \oplus \overset{1}{\circ} - \overset{1}{\circ} - \bullet$
SO(2) $\times$ Spin(7)	$(x \oplus x^{-1}) \otimes \overset{1}{\circ} - \overset{1}{\circ} - \bullet \oplus \overset{1}{\circ} - \overset{1}{\circ} - \bullet$
Spin(8)	$\overset{1}{\circ} - \overset{1}{\circ} - \overset{1}{\circ} \oplus \overset{1}{\circ} - \overset{1}{\circ} - \overset{1}{\circ}$
Spin(8)	$\overset{1}{\circ} - \overset{1}{\circ} - \overset{1}{\circ} \oplus \overset{1}{\circ} - \overset{1}{\circ} - \overset{1}{\circ}$
SO(2) $\times$ Spin(8)	$(x \oplus x^{-1}) \otimes \overset{1}{\circ} - \overset{1}{\circ} - \overset{1}{\circ} \oplus \overset{1}{\circ} - \overset{1}{\circ} - \overset{1}{\circ}$
SO(2) $\times$ Spin(8)	$(x \oplus x^{-1}) \otimes \overset{1}{\circ} - \overset{1}{\circ} - \overset{1}{\circ} \oplus \overset{1}{\circ} - \overset{1}{\circ} - \overset{1}{\circ}$
SO(3) $\times$ Spin(8)	$\overset{2}{\circ} \otimes \overset{1}{\circ} - \overset{1}{\circ} - \overset{1}{\circ} \oplus \overset{1}{\circ} - \overset{1}{\circ} - \overset{1}{\circ}$
SO(3) $\times$ Spin(8)	$\overset{2}{\circ} \otimes \overset{1}{\circ} - \overset{1}{\circ} - \overset{1}{\circ} \oplus \overset{1}{\circ} - \overset{1}{\circ} - \overset{1}{\circ}$

Sketch of a proof. Bergmann proves her theorem in [1] using Theorem 6.6 as it is stated in [15]. We sketch here an argument in a similar vein, but divided into cases in a different way. Since summands of almost faithful reducible representations need not be almost faithful, this is the only place in this paper where we have to exercise some care in distinguishing between a group G and its image $\rho(G)$ under a representation $\rho : G \rightarrow \mathbf{O}(V)$.

G IS A SIMPLE GROUP. Let $\rho = \rho_1 \oplus \rho_2$ be a decomposition where ρ_1 is irreducible. Then ρ_1, ρ_2 are variationally complete. Denote by V_1, V_2 the representation spaces and by H_1, H_2 the connected components of the principal isotropy subgroups. We have that ρ_1 is an almost faithful variationally complete irreducible representation of a simple group, which is orbit equivalent to the restriction $\rho_1|_{H_2}$, where H_2 is a proper subgroup of G ; see Proposition 3.4, item (b). Since $\rho_1(H_2) \subset \mathbf{O}(V_1)$ is not the maximal compact connected group with its orbits, $\rho_1 : H_2 \rightarrow \mathbf{O}(V_1)$ is not the isotropy representation of a symmetric space. It follows from Theorem 6.6 that ρ_1 has cohomogeneity one, because the larger of the two groups for each pair in the table of Proposition 6.3 is never a simple group. Since ρ_1 is an arbitrary irreducible summand of ρ , we conclude that every irreducible summand of ρ must have cohomogeneity one. Moreover, we know from Proposition 3.4, item (c), that there cannot be two equivalent summands. The only simple groups admitting at least two inequivalent cohomogeneity one representations are **Spin**(n) for $n \in \{3, 5, 6, 7, 8, 9\}$. Using

the fact that ρ_2 must be orbit equivalent to $\rho_2|_{H_1}$ we exclude the groups **Spin**(3), **Spin**(5), **Spin**(9) and get the four examples corresponding to simple groups in the table.

G IS NOT SEMISIMPLE. Let ρ_1 be an irreducible summand of ρ such that $\rho_1(G)$ is locally isomorphic to $S^1 \times G'$, where G' is a nontrivial semisimple factor of G . Let ρ_2 be another irreducible summand of ρ with the connected component of the principal isotropy group being H_2 . If $\rho_1(H_2) = \rho_1(G)$ for all such ρ_2 , then ρ is splitting. So we assume there exists such a ρ_2 with $\rho_1(H_2) \subsetneq \rho_1(G)$. Now we can view ρ_1 as an almost faithful variationally complete irreducible representation of $S^1 \times G'$, which is orbit equivalent to the restriction $\rho_1|_{H'_2}$, where H'_2 is a compact connected proper subgroup of $S^1 \times G'$ and $\rho_1(H'_2) = \rho_1(H_2)$. It follows from Theorem 6.6, item (b), that the possible local isomorphism classes of H'_2 ,

$S^1 \times G'$ and the representations $\rho_1|_{H'_2}$, $\rho_1|_{S^1 \times G'}$ are given as follows:

	H'_2 and $S^1 \times G'$	$\rho_1 _{H'_2}$ and $\rho_1 _{S^1 \times G'}$
1	$\mathbf{SO}(2) \times \mathbf{G}_2$ and $\mathbf{SO}(2) \times \mathbf{SO}(7)$	$(x \oplus x^{-1}) \otimes \overset{1}{\bullet \equiv \circ}$ and $(x \oplus x^{-1}) \otimes \overset{1}{\circ \equiv \bullet}$
2	$\mathbf{SO}(2) \times \mathbf{Spin}(7)$ and $\mathbf{SO}(2) \times \mathbf{SO}(8)$	$(x \oplus x^{-1}) \otimes \overset{1}{\circ \equiv \bullet}$ and $(x \oplus x^{-1}) \otimes \overset{1}{\circ \text{---} \circ \text{---} \circ}$
3	$\mathbf{Spin}(10)$ and $\mathbf{SO}(2) \times \mathbf{Spin}(10)$	$\overset{1}{\circ \text{---} \circ \text{---} \circ}$ and $(x \oplus x^{-1}) \otimes \overset{1}{\circ \text{---} \circ \text{---} \circ}$
4	$\mathbf{SU}(n)$, n odd, and $S^1 \times \mathbf{SU}(n)$	$\overset{1}{\circ \text{---} \circ} \dots \overset{1}{\circ \text{---} \circ} \oplus \overset{1}{\circ \text{---} \circ} \dots \overset{1}{\circ \text{---} \circ}$ and $(x \otimes \overset{1}{\circ \text{---} \circ} \dots \overset{1}{\circ \text{---} \circ}) \oplus (x^{-1} \otimes \overset{1}{\circ \text{---} \circ} \dots \overset{1}{\circ \text{---} \circ})$
5	$\mathbf{SU}(n) \times \mathbf{SU}(m)$, $n \neq m$ and $S^1 \times \mathbf{SU}(n) \times \mathbf{SU}(m)$	$(\overset{1}{\circ \text{---} \dots \text{---} \circ} \otimes \overset{1}{\circ \text{---} \dots \text{---} \circ})$ $\oplus (\overset{1}{\circ \text{---} \dots \text{---} \circ} \otimes \overset{1}{\circ \text{---} \dots \text{---} \circ})$ and $(\overset{1}{\circ \text{---} \dots \text{---} \circ} \otimes x \otimes \overset{1}{\circ \text{---} \dots \text{---} \circ})$ $\oplus (\overset{1}{\circ \text{---} \dots \text{---} \circ} \otimes x^{-1} \otimes \overset{1}{\circ \text{---} \dots \text{---} \circ})$
6	$\mathbf{Sp}(n)$ and $S^1 \times \mathbf{SU}(2n)$	$\overset{1}{\bullet \text{---} \dots \text{---} \bullet \equiv \circ}$ and $x \otimes \overset{1}{\circ \text{---} \dots \text{---} \circ} \oplus x^{-1} \otimes \overset{1}{\circ \text{---} \dots \text{---} \circ}$
7	$\mathbf{SU}(n)$ and $S^1 \times \mathbf{SU}(n)$	$\overset{1}{\circ \text{---} \dots \text{---} \circ} \oplus \overset{1}{\circ \text{---} \dots \text{---} \circ}$ and $x \otimes \overset{1}{\circ \text{---} \dots \text{---} \circ} \oplus x^{-1} \otimes \overset{1}{\circ \text{---} \dots \text{---} \circ}$
8	$S^1 \times \mathbf{Sp}(n)$ and $S^1 \times \mathbf{SU}(2n)$	$(x \oplus x^{-1}) \otimes \overset{1}{\bullet \text{---} \dots \text{---} \bullet \equiv \circ}$ and $x \otimes \overset{1}{\circ \text{---} \dots \text{---} \circ} \oplus x^{-1} \otimes \overset{1}{\circ \text{---} \dots \text{---} \circ}$
9	$\mathbf{Sp}(n)$ and $S^1 \times \mathbf{Sp}(n)$	$\overset{1}{\bullet \text{---} \dots \text{---} \bullet \equiv \circ}$ and $(x \oplus x^{-1}) \otimes \overset{1}{\bullet \text{---} \dots \text{---} \bullet \equiv \circ}$

In the following we examine different possibilities for ρ_2 . Recall that a variationally complete representation does not admit two equivalent nontrivial irreducible summands.

- (i) ρ_2 is as in case 1. Here ρ_2 is trivial on $\mathbf{SO}(2)$, so we can view ρ_2 as an almost faithful representation $\mathbf{Spin}(7) \times K \rightarrow \mathbf{O}(V_2)$ (K is a compact connected Lie group) which

is variationally complete, irreducible and such that the projection of the connected component of its principal isotropy subgroup in $\mathbf{Spin}(7)$ is $\mathbf{G}_2$. The only possibility for ρ_2 is $\circ \text{---} \circ \text{---} \bullet$, $K = \{1\}$. Any other irreducible summand ρ_3 of ρ either would satisfy $\rho_1(H_3) = \rho_1(G)$ (where H_3 is the connected component of the principal isotropy subgroup of ρ_3) or would have to be in the same case as ρ_2 . The first possibility would make ρ splitting and the second possibility is excluded because ρ cannot have two equivalent irreducible summands. It follows that ρ has exactly two irreducible summands and it is $(x \oplus x^{-1}) \otimes \circ \text{---} \circ \text{---} \bullet \oplus \circ \text{---} \circ \text{---} \bullet$.

- (ii) ρ_2 is as in case 2. Here ρ_2 is trivial on $\mathbf{SO}(2)$, so we can view ρ_2 as an almost faithful representation $\mathbf{Spin}(8) \times K \rightarrow \mathbf{O}(V_2)$ (K is a compact connected Lie group) which is variationally complete, irreducible and such that the projection of the connected component of its principal isotropy subgroup in $\mathbf{Spin}(8)$ is $\mathbf{Spin}(7)$. Here $\mathbf{Spin}(7)$ is in the conjugacy class such that the vector representation of $\mathbf{Spin}(8)$ restricts to the spin representation of $\mathbf{Spin}(7)$. The only possibilities for ρ_2 are the half-spin representations of $\mathbf{Spin}(8)$, $K = \{1\}$. Similar considerations to those in (i) show that ρ must have exactly two irreducible summands and that it is either one of

$$(x \oplus x^{-1}) \otimes \circ \text{---} \circ \text{---} \circ \oplus \circ \text{---} \circ \text{---} \circ, \quad (x \oplus x^{-1}) \otimes \circ \text{---} \circ \text{---} \circ \oplus \circ \text{---} \circ \text{---} \circ.$$

- (iii) ρ_2 is as in one of the cases 3, 4, 5 or 9. Here S^1 is not essential for ρ_1 and $G' \subset \ker \rho_2$. Also, for any other irreducible summand ρ_3 either $\rho_1(H_3) = \rho_1(G)$ (where H_3 is the connected component of the principal isotropy subgroup of ρ_3) or $G' \subset \ker \rho_3$. It follows that ρ is almost splitting.
- (iv) ρ_2 is as in case 6. This case cannot occur, for otherwise $\rho_2 : \mathbf{U}(2n) \times K \rightarrow \mathbf{O}(V_2)$ (K a compact connected Lie group) is an almost faithful variationally complete irreducible representation such that the projection of the connected component of its principal isotropy subgroup in $\mathbf{U}(2n)$ is $\mathbf{Sp}(n)$. But such a ρ_2 does not exist.
- (v) ρ_2 is as in case 7. Here S^1 is not essential for ρ_1 and $G' \subset \ker \rho_2$. Also, for any other irreducible summand ρ_3 we have either one of the following: $\rho_1(H_3) = \rho_1(G)$ (where H_3 is the connected component of the principal isotropy subgroup of ρ_3); or $G' \subset \ker \rho_3$; or ρ_3 is as in case 8. In the first two possibilities ρ is almost splitting and in the third one we go to the next case.
- (vi) ρ_2 is as in case 8. Here ρ_2 is trivial on S^1 , so we can view ρ_2 as an almost faithful representation $\mathbf{SU}(2n) \times K \rightarrow \mathbf{O}(V_2)$ (K is a compact connected Lie group) which is variationally complete, irreducible and such that the projection of the connected component of its principal isotropy subgroup in $\mathbf{SU}(2n)$ is $\mathbf{Sp}(n)$. The only possibility for ρ_2 is $\circ \text{---} \circ \text{---} \circ$, $K = \{1\}$, $n = 2$. If ρ_3 is a third irreducible summand of ρ then either $\rho_1(H_3) = \rho_1(G)$ (where H_3 is the connected component of the principal isotropy subgroup of ρ_3) or $\rho_1(H_3)$ must be as in case 7, therefore ρ_3 is trivial on $\mathbf{SU}(4)$. So if

there is any irreducible summand besides ρ_1 and ρ_2 then ρ is almost splitting. Therefore there are no other irreducible summands and ρ is

$$(x \otimes \overset{1}{\circ} \text{---} \overset{1}{\circ} \text{---} \overset{1}{\circ} \oplus x^{-1} \otimes \overset{1}{\circ} \text{---} \overset{1}{\circ} \text{---} \overset{1}{\circ}) \oplus \overset{1}{\circ} \text{---} \overset{1}{\circ} \text{---} \overset{1}{\circ}.$$

G IS SEMISIMPLE BUT NOT SIMPLE. Let ρ_1 be an irreducible summand of ρ such that $\rho_1(G)$ is locally isomorphic to G' , a semisimple but not simple factor of G . Let ρ_2 be another irreducible summand of ρ with connected component of the principal isotropy group H_2 . If $\rho_1(H_2) = \rho_1(G)$ for all such ρ_2 , then ρ is splitting. So we assume there exists such a ρ_2 with $\rho_1(H_2) \subsetneq \rho_1(G)$. Now we can view ρ_1 as an almost faithful variationally complete irreducible representation of G' , which is orbit equivalent to the restriction $\rho_1|_{H'_2}$, where H'_2 is a compact connected proper subgroup of G' and $\rho_1(H'_2) = \rho_1(H_2)$. It follows from Theorem 6.6, item (b), that the possible local isomorphism classes of H'_2 , G' and the representations $\rho_1|_{H'_2}$, $\rho_1|_{G'}$ are given as follows:

	H'_2 and G'	$\rho_1 _{H'_2}$ and $\rho_1 _{G'}$
10	$\mathbf{Sp}(n)$ and $\mathbf{Sp}(1) \times \mathbf{Sp}(n)$	$\overset{1}{\bullet} \text{---} \cdots \text{---} \bullet \text{---} \bullet \text{---} \bullet$ and $\overset{1}{\circ} \otimes \overset{1}{\bullet} \text{---} \cdots \text{---} \bullet \text{---} \bullet \text{---} \bullet$
11	$\mathbf{SO}(3) \times \mathbf{Spin}(7)$ and $\mathbf{SO}(3) \times \mathbf{SO}(8)$	$\overset{2}{\circ} \otimes \overset{1}{\circ} \text{---} \bullet \text{---} \bullet \text{---} \bullet$ and $\overset{2}{\circ} \otimes \overset{1}{\circ} \text{---} \bullet \text{---} \bullet \text{---} \bullet$

If ρ_2 is an in case 10, then ρ is almost splitting by the same argument as in item (iii). If ρ_2 is as in case 11, then by an argument similar to that in item (ii) we show that ρ must be either one of

$$\overset{2}{\circ} \otimes \overset{1}{\circ} \text{---} \bullet \text{---} \bullet \text{---} \bullet \oplus \overset{1}{\circ} \text{---} \bullet \text{---} \bullet \text{---} \bullet, \quad \overset{2}{\circ} \otimes \overset{1}{\circ} \text{---} \bullet \text{---} \bullet \text{---} \bullet \oplus \overset{1}{\circ} \text{---} \bullet \text{---} \bullet \text{---} \bullet.$$

□

7 Taut representations

The main result of this section is the following classification theorem.

7.1 Theorem *A taut irreducible representation ρ of a compact connected Lie group G is either orbit equivalent to the isotropy representation of a symmetric space or it is one of the*

following orthogonal representations ($n \geq 2$):

G	ρ
$\mathbf{SO}(2) \times \mathbf{Spin}(9)$	$(x \oplus x^{-1}) \otimes \overset{1}{\circ} \text{---} \overset{1}{\circ} \text{---} \overset{1}{\bullet}$
$\mathbf{U}(2) \times \mathbf{Sp}(n)$	$(x \oplus x^{-1}) \otimes \overset{1}{\circ} \otimes \overset{1}{\bullet} \text{---} \cdots \text{---} \bullet \text{---} \bullet \text{---} \circ$
$\mathbf{SU}(2) \times \mathbf{Sp}(n)$	$\overset{3}{\circ} \otimes \overset{1}{\bullet} \text{---} \cdots \text{---} \bullet \text{---} \bullet \text{---} \circ$

7.2 Corollary *A taut irreducible representation of a compact connected simple Lie group is orbit equivalent to the isotropy representation of a symmetric space.*

7.3 Remark According to [46], see also [40], the representations given in the table of Theorem 7.1 are exactly the irreducible representations of compact connected Lie groups which have cohomogeneity three and are not orbit equivalent to the isotropy representation of a symmetric space.

It follows from the main results of Section 6 that Theorem 7.1 is equivalent to the following:

7.4 Theorem *A taut irreducible representation ρ of a compact connected Lie group G is one of the following:*

- (a) *The isotropy representation of an irreducible symmetric space (listed in Tables 8.11.2 and 8.11.5 of [45]).*
- (b) *A cohomogeneity one representation not listed in (a) (cf. Theorem 8.2).*
- (c) *A representation orbit equivalent to the isotropy representation of an irreducible symmetric space and not listed in (a) or (b) (cf. Proposition 6.3).*
- (d) *One of the three exceptional cases given in Theorem 7.1.*

Proof of Theorems 7.1 and 7.4. A taut irreducible representation ρ of a compact connected Lie group G is of class $\mathcal{O}^2$. If ρ is not orbit equivalent to the isotropy representation of one of the irreducible symmetric spaces, we have obtained all possibilities for it, and these are listed in Tables C.1, C.2, C.3 and C.4 in Appendix C. Now we already know from Proposition 6.2 that the representations in Table C.4 are not taut. Therefore the assertion follows from Propositions 7.5, 7.6 and 7.12 below. $\square$

We start by considering Table C.3.

7.5 Proposition *The representations listed in Table C.3 are not taut.*

Proof. Let $G = \mathbf{G}_2 \times \mathbf{Spin}(7)$. Then ρ is the tensor product of the 7-dimensional representation of $\mathbf{G}_2$ on $\mathbf{R}^7$ and the spin representation of $\mathbf{Spin}(7)$ on $\mathbf{R}^8$. Let $p = v_1 \otimes v_2 \in \mathbf{R}^7 \otimes \mathbf{R}^8$. The isotropy subalgebra at p is $\mathfrak{su}(3) + \mathfrak{g}_2$. The dimension of the orbit through p is $\dim G(p) = 13$, the dimension of the normal space at p is 43, and the slice representation at p minus the trivial component $\mathbf{R}p$ is given by the tensor product of the vector representation of $\mathfrak{su}(3)$ on $\mathbf{R}^6$ and the 7-dimensional representation of $\mathfrak{g}_2$ on $\mathbf{R}^7$. Since this representation is of complex type and Dadok invariant 3, it follows from Proposition 4.9 that it cannot be of class $\mathcal{O}^2$. In particular, it is not taut. Now Proposition 3.8 implies that ρ is not taut.

Let $G = \mathbf{Spin}(7) \times \mathbf{Spin}(9)$. Then ρ is the tensor product of the spin representation of $\mathbf{Spin}(7)$ on $\mathbf{R}^8$ and the spin representation of $\mathbf{Spin}(9)$ on $\mathbf{R}^{16}$. Let $p = v_1 \otimes v_2 \in \mathbf{R}^8 \otimes \mathbf{R}^{16}$. The isotropy subalgebra at p is $\mathfrak{g}_2 + \mathfrak{spin}(7)$. The dimension of the orbit through p is $\dim G(p) = 22$, the dimension of the normal space at p is 106, and the slice representation at p minus the trivial component $\mathbf{R}p$ is given by the real tensor product of the 7-dimensional representation of $\mathfrak{g}_2$ and the vector representation of $\mathfrak{spin}(7)$ plus the real tensor product of the 7-dimensional representation of $\mathfrak{g}_2$ and the spin representation of $\mathfrak{spin}(7)$. Since we already know that the tensor product of the 7-dimensional representation of $\mathfrak{g}_2$ and the spin representation of $\mathfrak{spin}(7)$ is not taut (see the last paragraph), it follows from Proposition 3.8 that ρ is not taut either.

Let $G = \mathbf{G}_2 \times \mathbf{Spin}(9)$. Then ρ is the tensor product of the 7-dimensional representation of $\mathbf{G}_2$ on $\mathbf{R}^7$ and the spin representation of $\mathbf{Spin}(9)$ on $\mathbf{R}^{16}$. Let $p = v_1 \otimes v_2 \in \mathbf{R}^7 \otimes \mathbf{R}^{16}$. The isotropy subalgebra at p is $\mathfrak{su}(3) + \mathfrak{spin}(7)$. The dimension of the orbit through p is $\dim G(p) = 21$, the dimension of the normal space at p is 91, and the slice representation at p minus the trivial component $\mathbf{R}p$ is given by the real tensor product of the vector representations of $\mathfrak{su}(3)$ and $\mathfrak{spin}(7)$ plus the real tensor product of the vector representation of $\mathfrak{su}(3)$ and the spin representation of $\mathfrak{spin}(7)$. Since the second summand is a representation of complex type and Dadok invariant 3, it follows from Proposition 4.9 that it cannot be of class $\mathcal{O}^2$. In particular, it is not taut. Now Proposition 3.8 implies that ρ is not taut.

Consider⁶ $G = \mathbf{Sp}(1) \cdot \mathbf{Sp}(n) \times \mathbf{G}_2$, $n \geq 2$. Let τ_n be a real form of $\circ \otimes \circ \text{---} \circ \text{---} \dots \text{---} \circ$. Then τ_n can be realized as the representation of $\mathbf{Sp}(1) \cdot \mathbf{Sp}(n)$ on $\mathbf{R}^{4n}$ given by $\tau_n(q, A)x = Axq^{-1}$, where $q \in \mathbf{Sp}(1)$, $A \in \mathbf{Sp}(n)$, $x \in \mathbf{H}^n$ and we identify $\mathbf{H}^n \cong \mathbf{R}^{4n}$. Now ρ is the real tensor product of τ_n and the 7-dimensional representation of $\mathbf{G}_2$. Let $p = v_1 \otimes v_2 \in \mathbf{R}^{4n} \otimes \mathbf{R}^7$. The isotropy subalgebra at p is isomorphic to $\mathfrak{sp}(1) + \mathfrak{sp}(n-1) + \mathfrak{su}(3)$. The dimension of the orbit through p is $\dim G(p) = 4n + 5$, the dimension of the normal space at p is $24n - 5$, and the slice representation at p minus the trivial component $\mathbf{R}p$ is given by the real tensor product of τ_{n-1} and the vector representation of $\mathfrak{su}(3)$ plus the real tensor product of the adjoint representation of $\mathfrak{sp}(1)$ and the the vector representation of $\mathfrak{su}(3)$. Since the second summand is a representation of complex type and Dadok invariant 3, it follows from Proposition 4.9 that it cannot be of class $\mathcal{O}^2$. In particular, it is not taut. Now Proposition 3.8 implies that ρ is not taut.

Consider $G = \mathbf{Sp}(1) \cdot \mathbf{Sp}(n) \times \mathbf{Spin}(7)$, $n \geq 2$. Let τ_n be as above. Now ρ is the tensor product of τ_n and the spin representation of $\mathbf{Spin}(7)$. Let $p = v_1 \otimes v_2 \in \mathbf{R}^{4n} \otimes \mathbf{R}^8$. The isotropy subalgebra at p is isomorphic to $\mathfrak{sp}(1) + \mathfrak{sp}(n-1) + \mathfrak{g}_2$. The dimension of the orbit through p is $\dim G(p) = 4n + 6$, the dimension of the normal space at p is $28n - 6$,

⁶We use the convention that $G \cdot H$ refers to a quotient of $G \times H$ by a finite central subgroup.

and the slice representation at p minus the trivial component $\mathbf{R}p$ is given by the real tensor product of τ_{n-1} and the 7-dimensional representation of $\mathfrak{g}_2$ plus the real tensor product of the adjoint representation of $\mathfrak{sp}(1)$ and the 7-dimensional representation of $\mathfrak{g}_2$. Since the first summand is not a taut representation by the above (for $n = 2$ we refer to the case of $G = \mathbf{SO}(4) \times \mathbf{G}_2$ which will be dealt with in Proposition 7.6), it follows from Proposition 3.8 that ρ is not taut either.

Let $G = \mathbf{Spin}(7) \times \mathbf{Spin}(7)$. Then ρ is the tensor product of the spin representations of each of the factors on $\mathbf{R}^8$. Let $p = v_1 \otimes v_2 \in \mathbf{R}^8 \otimes \mathbf{R}^8$. The isotropy subalgebra at p is $\mathfrak{g}_2 + \mathfrak{g}_2$. The dimension of the orbit through p is $\dim G(p) = 14$, the dimension of the normal space at p is 50, and the slice representation at p minus the trivial component $\mathbf{R}p$ is given by the real tensor product of the 7-dimensional representations of each of the factors. Since this representation is not taut (see next case), it follows from Proposition 3.8 that ρ is not taut either.

Consider $G = \mathbf{G}_2 \times \mathbf{G}_2$. Then ρ is the tensor product of the 7-dimensional representations of each of the factors on $\mathbf{R}^7$. Let $\{e_1, \dots, e_7\}$ be an orthonormal basis of $\mathbf{R}^7$ and take $p = e_1 \otimes e_1 + \dots + e_7 \otimes e_7$. Then the isotropy subalgebra at p is

$$\mathfrak{g}_p = \{(X, X) : X \in \mathfrak{g}_2\}$$

which is isomorphic to $\mathfrak{g}_2$. Now the restriction of ρ to $\mathfrak{g}_p$ is a real form of

$$\mathfrak{g}_2 : \overset{1}{\bullet} \equiv \overset{1}{\circ} \otimes (\overset{1}{\bullet} \equiv \overset{1}{\circ})^* = \overset{1}{\bullet} \equiv \overset{1}{\circ} \otimes \overset{1}{\bullet} \equiv \overset{1}{\circ}$$

acting on $\mathbf{R}^{49}$, and $\mathbf{R}^{49} = T_p G(p) \oplus N_p G(p)$ is a $\mathfrak{g}_p$ -invariant decomposition. We have the following decomposition into irreducible components:

$$\begin{aligned} \mathfrak{g}_2 : \otimes^2(\overset{1}{\bullet} \equiv \overset{1}{\circ}) &= S^2(\overset{1}{\bullet} \equiv \overset{1}{\circ}) \oplus \Lambda^2(\overset{1}{\bullet} \equiv \overset{1}{\circ}) \\ &= (\overset{2}{\bullet} \equiv \overset{2}{\circ} \oplus (\text{trivial})) \oplus (\overset{1}{\bullet} \equiv \overset{1}{\circ} \oplus \overset{1}{\bullet} \equiv \overset{1}{\circ}). \end{aligned}$$

Since $T_p G(p)$ is 14-dimensional, it follows that the 27-dimensional real representation $\overset{2}{\bullet} \equiv \overset{2}{\circ}$ must be a component of the slice representation at p . But we saw in Lemma 5.39 that it is not of class $\mathcal{O}^2$. Now Proposition 3.8 implies that ρ is not taut. $\square$

7.6 Proposition *The representations listed in Table C.2 are not taut.*

Proof. Let $G = \mathbf{SO}(m) \times \mathbf{G}_2$, $m \geq 4$. Then ρ is the tensor product of the vector representation of $\mathbf{SO}(m)$ on $\mathbf{R}^m$ and the 7-dimensional representation of $\mathbf{G}_2$ on $\mathbf{R}^7$. Let $p = v_1 \otimes v_2 \in \mathbf{R}^m \otimes \mathbf{R}^7$. The isotropy subalgebra at p is $\mathfrak{so}(m-1) + \mathfrak{su}(3)$. The dimension of the orbit through p is $\dim G(p) = m+5$, the dimension of the normal space at p is $6m-5$, and the slice representation at p minus the trivial component $\mathbf{R}p$ is given by the tensor product of the vector representations of $\mathfrak{so}(m-1)$ and $\mathfrak{su}(3)$ on $\mathbf{R}^{m-1}$ and $\mathbf{R}^6$, respectively. Since this is a representation of complex type and Dadok invariant 3, it follows from Proposition 4.9 that it cannot be of class $\mathcal{O}^2$. In particular, it is not taut. Now Proposition 3.8 implies that ρ is not taut.

The proof for $G = \mathbf{SO}(3) \times \mathbf{G}_2$ will be done in Lemma 7.8 below, since this case is more involved.

Let $G = \mathbf{SO}(m) \times \mathbf{Spin}(7)$, $m \geq 4$. Then ρ is the tensor product of the vector representation of $\mathbf{SO}(m)$ on $\mathbf{R}^m$ and the spin representation of $\mathbf{Spin}(7)$ on $\mathbf{R}^8$. Let $p = v_1 \otimes v_2 \in \mathbf{R}^m \otimes \mathbf{R}^8$. The isotropy subalgebra at p is $\mathfrak{so}(m-1) + \mathfrak{g}_2$. The dimension of the orbit through p is $\dim G(p) = m+6$, the dimension of the normal space at p is $7m-6$, and the slice representation at p minus the trivial component $\mathbf{R}p$ is given by the real tensor product of the vector representation of $\mathfrak{so}(m-1)$ and the 7-dimensional representation of $\mathfrak{g}_2$. Since this representation is not taut by the above, it follows from Proposition 3.8 that ρ is not taut either.

Let $G = \mathbf{SO}(m) \times \mathbf{Spin}(9)$, $m \geq 4$. Then ρ is the tensor product of the vector representation of $\mathbf{SO}(m)$ on $\mathbf{R}^m$ and the spin representation of $\mathbf{Spin}(9)$ on $\mathbf{R}^{16}$. Let $p = v_1 \otimes v_2 \in \mathbf{R}^m \otimes \mathbf{R}^{16}$. The isotropy subalgebra at p is $\mathfrak{so}(m-1) + \mathfrak{spin}(7)$. The dimension of the orbit through p is $\dim G(p) = m+14$, the dimension of the normal space at p is $15m-14$, and the slice representation at p minus the trivial component $\mathbf{R}p$ is given by the real tensor product of the vector representations of $\mathfrak{so}(m-1)$ and $\mathfrak{spin}(7)$ plus the real tensor product of the vector representation of $\mathfrak{so}(m-1)$ and the spin representation of $\mathfrak{spin}(7)$. If $m \geq 5$ then the second summand is not a taut representation by the above and it follows from Proposition 3.8 that ρ is not taut either. In the case $m = 4$ we give a special argument. Here we have that the connected component of the isotropy subgroup at p is $K = \mathbf{SO}(3) \times \mathbf{Spin}(7)$. Let ν_1, ν_2 be respectively the irreducible summands of dimensions 21, 24 of the slice representation at p , and denote with V_1, V_2 the representation spaces. Note that both ν_1 and ν_2 are taut representations of K , since ν_1 is the isotropy representation of a real Grassmann manifold and ν_2 is orbit equivalent to the isotropy representation of a real Grassmann manifold. Let $q_1 \in V_1$ be a regular point. Then the isotropy group K_{q_1} is isomorphic to $\mathbf{Z}_2^2 \times \mathbf{Spin}(4)$. We can choose $q_2 \in V_2$ so that the isotropy $(K_{q_1})_{q_2}$ is contained in $\{1\} \times \mathbf{Spin}(4)$. Hence the orbit $K_{q_1}(q_2)$ is disconnected. Since the orbit $K(q_1, q_2)$ is connected, Proposition 3.11 implies that the slice representation at p is not taut. Now Proposition 3.8 yields that ρ is not taut either.

We postpone the proof for the case $\mathbf{SO}(3) \times \mathbf{Spin}(9)$ to Lemma 7.7 below.

Let $G = \mathbf{SU}(n) \times \mathbf{Sp}(m)$, $n \geq 3$ and $m \geq 2$. Then ρ is the realification of the tensor product of the vector representations of $\mathbf{SU}(n)$ on $\mathbf{C}^n$ and of $\mathbf{Sp}(m)$ on $\mathbf{C}^{2m}$. Consider complex bases $\{e_1, \dots, e_n\}$ for $\mathbf{C}^n$ and $\{f_1, \dots, f_m, f_{m+1}, \dots, f_{2m}\}$ for $\mathbf{C}^{2m}$ where $f_{m+j} = \epsilon f_j$, ϵ the quaternionic structure on $\mathbf{C}^{2m}$. Let $p = e_1 \otimes f_1$. Then the isotropy $K = G_p$ is isomorphic to $T_0 \cdot \mathbf{SU}(n-1) \times \mathbf{Sp}(m-1)$, where T_0 is the circle group generated by

$$\begin{aligned} \varphi_0(t) = & (\text{diag}(e^{(n-1)it}, e^{-it}, \dots, e^{-it}), \\ & \text{diag}(e^{-(n-1)it}, 1, \dots, 1; e^{(n-1)it}, 1, \dots, 1)) \in \mathbf{SU}(n) \times \mathbf{Sp}(m), \end{aligned}$$

and the slice representation ν_p minus the trivial component corresponding to the radial direction $\mathbf{R}p$ decomposes into irreducible components as $\nu_1 \oplus \nu_2$, where ν_1 restricted to $\mathbf{SU}(n-1) \times \mathbf{Sp}(m-1)$ is the realification of the complex tensor product of the vector representations of the factors and ν_2 is the realification of the vector representation on $\mathbf{SU}(n-1)$ and it is trivial on $\mathbf{Sp}(m-1)$. Denote the representation spaces by V_1, V_2 . Then V_1 is spanned by $e_\alpha \otimes f_\beta$ and $i(e_\alpha \otimes f_\beta)$, for $2 \leq \alpha \leq n$, $2 \leq \beta \leq m$ or $m+2 \leq \beta \leq 2m$, and V_2 is spanned by $e_\alpha \otimes f_{m+1}$ and $i(e_\alpha \otimes f_{m+1})$, for $2 \leq \alpha \leq n$. Let $q_1 = e_2 \otimes f_2 \in V_1$. Then the isotropy $K_1 = K_{q_1}$ is isomorphic to $T_1 \cdot T_2 \cdot \mathbf{SU}(n-2) \times \mathbf{Sp}(m-2)$, where T_1, T_2 are circle

groups respectively generated by

$$\begin{aligned} \varphi_1(t) = & (\text{diag}(1, e^{(n-2)it}, e^{-it}, \dots, e^{-it}), \\ & \text{diag}(1, e^{-(n-2)it}, 1, \dots, 1; 1, e^{(n-2)it}, 1, \dots, 1)) \in \mathbf{SU}(n) \times \mathbf{Sp}(m), \end{aligned}$$

and

$$\begin{aligned} \varphi_2(t) = & (\text{diag}(e^{(n-1)it}, e^{-it}, e^{-it}, \dots, e^{-it}), \\ & \text{diag}(e^{-(n-1)it}, e^{it}, 1, \dots, 1; e^{(n-1)it}, e^{-it}, 1, \dots, 1)) \in \mathbf{SU}(n) \times \mathbf{Sp}(m). \end{aligned}$$

(Here we have used the hypothesis $n \geq 3$ and $m \geq 2$). Finally let $q_2 = e_2 \otimes f_{m+1} \in V_2$. Then the isotropy $K_2 = (K_{q_1})_{q_2}$ is isomorphic to $T_3 \cdot \mathbf{SU}(n-2) \times \mathbf{Sp}(m-2)$, where T_3 is the circle subgroup of $T_1 \cdot T_2$ generated by $\varphi_3(t) = \varphi_1(-t)\varphi_2(t)$. It is easy to see that K_1/K_2 is diffeomorphic to S^1 , whereas K/K_2 is simply-connected. Therefore Proposition 3.11 implies that ν_p is not taut. Now Proposition 3.8 yields that ρ is not taut either.

The case where $G = \mathbf{U}(n) \times \mathbf{Sp}(m)$, $n \geq 3$ and $m \geq 2$, is completely analogous to the previous case.

Let $G = \mathbf{SO}(m) \times \mathbf{Sp}(1) \cdot \mathbf{Sp}(n)$, $m \geq 3$ and $n \geq 2$. Then ρ is the tensor product of the vector representation of $\mathbf{SO}(m)$ on $\mathbf{R}^m$ and τ_n , where τ_n is a representation on $\mathbf{R}^{4n}$ as in Proposition 7.5. We choose an orthonormal basis $\{e_1, \dots, e_m\}$ for $\mathbf{R}^m$ and an orthonormal basis

$$\{f_1, \dots, f_n, f_1i, \dots, f_ni, f_1j, \dots, f_nj, f_1k, \dots, f_nk\}$$

for $\mathbf{R}^{4n}$. Let $p = e_1 \otimes f_1$. The connected component of the isotropy subgroup $K = G_p^0$ is isomorphic to $\mathbf{SO}(m-1) \times \mathbf{Sp}(1)' \cdot \mathbf{Sp}(n-1)$, where

$$\mathbf{Sp}(1)' = \left\{ \left(q, \begin{pmatrix} q & & & \\ & 1 & & \\ & & \ddots & \\ & & & 1 \end{pmatrix} \right) : q \in \mathbf{Sp}(1) \right\} \subset \mathbf{Sp}(1) \cdot \mathbf{Sp}(n).$$

The slice representation at p minus the trivial component $\mathbf{R}p$ decomposes as $\nu_1 \oplus \nu_2$, where ν_1 is the real tensor product of the vector representation of $\mathbf{SO}(m-1)$ and τ_{n-1} , and ν_2 is the real tensor product of the vector representation of $\mathbf{SO}(m-1)$ and the adjoint representation of $\mathbf{Sp}(1)$ (ν_2 is trivial on $\mathbf{Sp}(n-1)$). Note that ν_2 is the isotropy representation of a real Grassmann manifold. Let V_1, V_2 denote the representation spaces of ν_1, ν_2 . Then V_1 is spanned by $e_\alpha \otimes f_\beta, e_\alpha \otimes f_\beta i, e_\alpha \otimes f_\beta j, e_\alpha \otimes f_\beta k$, for $2 \leq \alpha \leq m, 2 \leq \beta \leq n$, and V_2 is spanned by $e_\alpha \otimes f_1i, e_\alpha \otimes f_1j, e_\alpha \otimes f_1k$, for $2 \leq \alpha \leq m$. First consider the case $m \geq 4$. Let $q_2 \in V_2$ be a regular point, say $q_2 = ae_2 \otimes f_1i + be_3 \otimes f_1j + ce_4 \otimes f_1k$, where a, b, c are pairwise distinct positive real numbers. Then the isotropy group K_{q_2} is isomorphic to $\mathbf{SO}(m-4) \times Q \cdot \mathbf{Sp}(n-1)$, where Q is the quaternion subgroup $\{\pm 1, \pm i, \pm j, \pm k\}$ of $\mathbf{Sp}(1)$. Consider the restriction of ν_1 to K_{q_2} . We can choose $q_1 \in V_1$ so that the orbit $K_{q_2}(q_1)$ is disconnected, say $q_1 = ae_2 \otimes f_2 + be_3 \otimes f_2i$, where a, b are distinct positive real numbers. In fact, if $(A, q, B) \in \mathbf{SO}(m-4) \times Q \cdot \mathbf{Sp}(n-1)$ is in the isotropy of q_1 , then we must have $Bf_2q^{-1} = f_2$ and $Bf_2iq^{-1} = f_2i$, which implies that $Bf_2q^{-1}i = Bf_2iq^{-1}$

and then $q^{-1}i = iq^{-1}$, so that $q = \pm 1$ or $q = \pm i$. In particular, no element of the form $(A, j, B) \in \mathbf{SO}(m-4) \times Q \cdot \mathbf{Sp}(n-1)$ is in the isotropy of q_1 , which implies that the orbit $K_{q_2}(q_1)$ is disconnected. Since the orbit $K(q_1, q_2)$ is connected, Proposition 3.11 implies that the slice representation at p is not taut. Now Proposition 3.8 yields that ρ is not taut either. Finally we analyse the case $m = 3$. Then $K = \mathbf{SO}(2) \times \mathbf{Sp}(1)' \cdot \mathbf{Sp}(n-1)$. Consider a regular point $q_2 \in V_2$, say $q_2 = ae_2 \otimes f_1i + be_3 \otimes f_1j$, where a, b are distinct positive real numbers. Then we have that the isotropy K_{q_2} is isomorphic to $\mathbf{Z}_4 \times \mathbf{Sp}(n-1)$. Here $\mathbf{Z}_4$ is generated by $(-1, k)$, where -1 is minus the identity matrix in $\mathbf{SO}(2)$ and $k \in \mathbf{Sp}(1)'$. Consider the restriction of ν_1 to K_{q_2} . We can choose $q_1 \in V_1$ as above such that its isotropy does not contain elements of the form $(-1, k, B)$, for $B \in \mathbf{Sp}(n-1)$. It follows that the orbit $K_{q_2}(q_1)$ is disconnected and the proof follows as above. $\square$

7.7 Lemma *The tensor product ρ of the vector representation of $\mathbf{SO}(3)$ on $\mathbf{R}^3$ and the spin representation of $\mathbf{Spin}(9)$ on $\mathbf{R}^{16}$ is not taut.*

Proof. Let $p = v_1 \otimes v_2 \in \mathbf{R}^3 \otimes \mathbf{R}^{16}$. Then the connected component of the isotropy group $K = G_p^0$ is isomorphic to $\mathbf{SO}(2) \times \mathbf{Spin}(7)$. The normal space at p to the orbit $G(p)$ decomposes as $\mathbf{R}p \oplus V_1 \oplus V_2$, where $V_1 = \mathbf{R}^2 \otimes \mathbf{R}^7$, $V_2 = \mathbf{R}^2 \otimes \mathbf{R}^8$ are the representation spaces for the irreducible components ν_1, ν_2 of the slice representation at p . Here ν_1 is the tensor product of the vector representations and ν_2 is the tensor product of the vector representation of $\mathbf{SO}(2)$ and the spin representation of $\mathbf{Spin}(7)$. Now let $q = v_3 \otimes v_4 \in V_2$. Then the connected component of the isotropy group K_q^0 is isomorphic to $\mathbf{G}_2$ and the slice representation at q of the representation of K on $V_1 \oplus V_2$ is a representation of $\mathbf{G}_2$ on $\mathbf{R}^{22}$ which decomposes as a trivial component $\mathbf{R}q$ plus three copies of the 7-dimensional representation of $\mathbf{G}_2$. Next we shall show that the sum $\mu = \mu_1 \oplus \mu_2 \oplus \mu_3$ of three copies of 7-dimensional representation of $H = \mathbf{G}_2$ is not taut. It then follows from Proposition 3.8 that ρ is not taut either.

In fact, let W_i be the representation space of μ_i . We can select $w_i \in W_i$ such that the isotropy groups $H_{w_1}, (H_{w_1})_{w_2}, ((H_{w_1})_{w_2})_{w_3}$ are respectively $\mathbf{SU}(3), \mathbf{SU}(2), \{1\}$. Since $H_{(w_1, w_2, w_3)} = ((H_{w_1})_{w_2})_{w_3}$, we get that the orbit $H(w_1, w_2, w_3)$ is diffeomorphic to $\mathbf{G}_2$. Now if this orbit is taut, Proposition 3.11 implies that it must have the product homology of the orbits through w_1, w_2 and w_3 , namely, S^6, S^5 and S^3 . But it is well known that $\mathbf{G}_2$ has the product homology of S^{11} and S^3 (this also follows from Satz 1 in [25]). Thus, μ is not taut. $\square$

7.8 Lemma *The tensor product ρ of the vector representation of $\mathbf{SO}(3)$ on $\mathbf{R}^3$ and the 7-dimensional representation of $\mathbf{G}_2$ is not taut.*

Proof. We shall use the reduction technique described in Subsection 3.3. Note that $G = \mathbf{SO}(3) \times \mathbf{G}_2$ is a closed subgroup of $\hat{G} = \mathbf{SO}(3) \times \mathbf{SO}(7)$ and ρ is the restriction of a representation $\hat{\rho}$ of $\hat{G}$ which is the isotropy representation of a real Grassmann manifold.

We view the Cayley numbers $\mathbf{O} = \mathbf{H} \oplus e\mathbf{H}$, so that $\{1, e, i, j, k, ei, ej, ek\}$ is a real orthonormal basis for $\mathbf{O}$. The Lie group $\mathbf{G}_2$ is the automorphism group of $\mathbf{O}$ with respect to its nonassociative algebra structure. We next compute the space of orbits of G acting by ρ on $V = \mathbf{R}^3 \otimes \mathbf{R}^7$. Take $\{e, i, j, k, ei, ej, ek\}$ as a basis for $\mathbf{R}^7$ and take an orthonormal basis $\{f_1, f_2, f_3\}$ for $\mathbf{R}^3$. Recall that the subspace spanned by $\{i, j, k, ei, ej, ek\}$ has a complex

structure given by left multiplication by e . Since all these representations are self-adjoint, it is equivalent to consider ρ acting on $\mathbf{R}^{7*} \otimes \mathbf{R}^3$ which we may identify with the space of 3×7 real matrices.

Let

$$p_1 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

Then the orbit $G(p_1)$ is 8-dimensional and its normal space at p_1 intersects all orbits. So, given an arbitrary orbit M , it contains a point of the form

$$\begin{pmatrix} a & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & * & * & * & * & * & * \\ 0 & * & * & * & * & * & * \end{pmatrix},$$

for some $a \geq 0$, for we can multiply on the left by

$$\begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

to change the sign of a , if necessary, and remain in the orbit M . Now the connected component of the isotropy subgroup at p_1 , $K_1 = \mathbf{SO}(2) \times \mathbf{SU}(3)$, acts on the normal space to $G(p_1)$ by fixing the radial direction $\mathbf{R}p_1$ and by the tensor product of the vector representations of $\mathbf{SO}(2)$ and $\mathbf{SU}(3)$ on the orthogonal complement $\mathbf{R}^{12}$. The point

$$p_2 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

is in $\mathbf{R}^{12}$ and has a 6-dimensional K_1 -orbit. So M has a representative of the form

$$\begin{pmatrix} a & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & b & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & * & * & * & * & * \end{pmatrix},$$

for some $b \geq 0$, since we can also multiply on the left by

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix},$$

if necessary, and remain in the same orbit. Finally, the connected component of the isotropy subgroup of K_1 at p_2 is $\mathbf{SU}(2)$ and it acts on the normal space to $K_1(p_2)$ with respect to $\mathbf{R}^{12}$ by fixing a 2-plane and by the vector representation on the orthogonal complement. So M has an element of the form

$$(7.9) \quad \begin{pmatrix} a & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & b & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & c & 0 & d & 0 & 0 \end{pmatrix},$$

for some $c \geq 0$. We can assume $a \geq b$ by multiplying on the left by

$$\begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix},$$

and on the right by

$$\begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix},$$

if necessary. Multiplying (7.9) on the left by

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & 0 \end{pmatrix}$$

and on the right by

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & \frac{-c}{\sqrt{c^2+d^2}} & 0 & \frac{-d}{\sqrt{c^2+d^2}} & 0 & 0 \\ 0 & \frac{c}{\sqrt{c^2+d^2}} & 0 & 0 & 0 & \frac{d}{\sqrt{c^2+d^2}} & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & \frac{d}{\sqrt{c^2+d^2}} & 0 & 0 & 0 & \frac{-c}{\sqrt{c^2+d^2}} & 0 \\ 0 & 0 & \frac{-d}{\sqrt{c^2+d^2}} & 0 & \frac{c}{\sqrt{c^2+d^2}} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

we get

$$\begin{pmatrix} a & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & \sqrt{c^2+d^2} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & \frac{bc}{\sqrt{c^2+d^2}} & 0 & \frac{bd}{\sqrt{c^2+d^2}} & 0 & 0 \end{pmatrix}.$$

We can therefore assume $b \geq \sqrt{c^2+d^2}$. It is straightforward to check that no further restrictions can be put on a, b, c, d . Hence we see that the orbit space is described by the matrices of the form (7.9) subject to the conditions $a \geq b \geq \sqrt{c^2+d^2}$ and $c \geq 0$.

Next we want to compute a principal isotropy subgroup of G . Let p be an element of the form (7.9) which is regular for both G and $\hat{G}$. The principal isotropy group $\hat{G}_p$ is isomorphic to $\mathbf{Z}_2^2 \times \mathbf{SO}(4)$, and $G_p = \hat{G}_p \cap G$. Now the $\mathbf{Z}_2^2$ -factor is contained in G_p , because

$$(\text{diag}(-1, -1, 1), \text{diag}(-1, -1, 1, -1, 1, -1, 1))$$

and

$$(\text{diag}(1, -1, -1), \text{diag}(1, -1, -1, 1, -1, -1, 1))$$

are elements of G that fix p . On the other hand, an element in the $\mathbf{SO}(4)$ -factor that fixes p and is in $G = \mathbf{SO}(3) \times \mathbf{G}_2$ is in fact in $\{1\} \times \mathbf{G}_2$ and has to fix $e, i, cj + d(ei) \in \mathbf{R}^7$. Therefore it also fixes $j = \frac{1}{c}[(cj + d(ei)) - d(ei)]$ ($c \neq 0$ because p is regular for G) and hence it is the identity. We conclude that the principal isotropy group $H = G_p$ is exactly $\mathbf{Z}_2^2$.

It is immediate to see that the fixed subspace V^H is

$$\begin{pmatrix} * & 0 & 0 & * & 0 & 0 & 0 \\ 0 & * & 0 & 0 & 0 & * & 0 \\ 0 & 0 & * & 0 & * & 0 & 0 \end{pmatrix}.$$

Let N be the normalizer of H in G and $\bar{G} = N/H$ the reduced group as in Subsection 3.3. The connected component $\bar{G}^0$ must be contained in $\{1\} \times \mathbf{G}_2$. One can use the fact that N equals the normalizer of V^H in G to verify that $\bar{G}^0$ is the 2-torus group consisting of the matrices

$$(7.10) \quad \begin{pmatrix} \cos t & 0 & 0 & -\sin t & 0 & 0 & 0 \\ 0 & \cos s & 0 & 0 & 0 & -\sin s & 0 \\ 0 & 0 & \cos(s-t) & 0 & -\sin(s-t) & 0 & 0 \\ \sin t & 0 & 0 & \cos t & 0 & 0 & 0 \\ 0 & 0 & \sin(s-t) & 0 & \cos(s-t) & 0 & 0 \\ 0 & \sin s & 0 & 0 & 0 & \cos s & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

for $s, t \in \mathbf{R}$.

Let $p \in V^H$ be a regular point and let $M = Gp$ be the corresponding principal orbit in V . We next prove that M is not taut by contradiction. In fact, assume M is taut. The orbit $\bar{G}^0 p$ is a connected component of M^H , and it is a substantial 2-torus embedded in the 6-dimensional Euclidean space V^H . Since its second osculating space is at most 5-dimensional, it cannot be taut by Theorem 2.5. It is known that a surface is taut if and only if its lines of curvature are circles. Therefore there is a parallel normal vector field $\xi(t)$ along a curve $\gamma(t)$ in $\bar{G}^0 p \subset M^H$ such that $A_{\xi(t)}^{M^H} \dot{\gamma}(t) = \lambda(t) \dot{\gamma}(t)$ and $\lambda(t)$ is not constant. (*A posteriori* it is clear that one can find $\gamma(t), \xi(t), \lambda(t)$ directly by computation using (7.10).) Since M is a principal G -orbit, the normal spaces of M and M^H in V coincide along $\gamma(t)$ implying that $\xi(t)$ is also parallel with respect to M . By Lemma 3.16 we have

$$(7.11) \quad A_{\xi(t)}^M \dot{\gamma}(t) = \lambda(t) \dot{\gamma}(t).$$

We do not claim that $\gamma(t)$ is a curvature line of M since the eigenspace corresponding to $\lambda(t)$ might not be one-dimensional. Still the argument in [36], Lemmas 1 and 2 (see also [30]) carries through and shows that (7.11) together with the tautness of M implies that $\lambda(t)$ is constant, which is a contradiction. Hence M is not taut. Thus the representation ρ is not taut either. $\square$

7.12 Proposition *The representations listed in Table C.1 are taut.*

Proof. We use the reduction technique discussed in Subsection 3.3. Consider first the case of $G = \mathbf{U}(2) \times \mathbf{Sp}(n)$, $n \geq 2$, acting on the realification of $\mathbf{C}^2 \otimes \mathbf{C}^{2n}$ by the tensor product of the vector representations. It is equivalent to consider ρ acting on the space of $2 \times 2n$ complex matrices, i. e. $\rho(A, B)X = AXB^{-1}$ for $A \in \mathbf{U}(2)$, $B \in \mathbf{Sp}(n) \subset \mathbf{SU}(2n)$, $X \in M(2 \times 2n; \mathbf{C})$. It follows rather easily that the orbit space consists of matrices of the form

$$\begin{pmatrix} a & 0 & 0 & \dots & 0 & 0 & 0 & 0 & \dots & 0 \\ 0 & b & 0 & \dots & 0 & c & 0 & 0 & \dots & 0 \end{pmatrix},$$

where $a \geq \sqrt{b^2 + c^2}$ and $b, c \geq 0$. Since these matrices are all real, it follows that the group obtained by adjoining to G the complex conjugation of matrices σ still has the same orbits. We replace G by the enlarged group and continue to denote it with the same letter.

Now the principal isotropy group H is generated by σ and the subgroup of G isomorphic to $\mathbf{U}(1) \times \mathbf{Sp}(n-2)$ consisting of elements of the form

$$\left(\begin{pmatrix} e^{i\theta} & 0 \\ 0 & e^{-i\theta} \end{pmatrix}, \begin{pmatrix} e^{i\theta} & & & 0 \\ & e^{-i\theta} & & 0 \\ & & C & D \\ 0 & & e^{-i\theta} & \\ & 0 & & e^{i\theta} \\ & & -\bar{D} & \bar{C} \end{pmatrix} \right),$$

where $\theta \in \mathbf{R}$ and $\begin{pmatrix} C & D \\ -\bar{D} & \bar{C} \end{pmatrix} \in \mathbf{Sp}(n-2)$. Now V^H is the space of matrices of the form

$$\begin{pmatrix} a & 0 & 0 & \dots & 0 & 0 & d & 0 & \dots & 0 \\ 0 & b & 0 & \dots & 0 & c & 0 & 0 & \dots & 0 \end{pmatrix},$$

whose entries are real, and $(N/H)^0$ is the circle group consisting of the coclasses defined by the matrices

$$\begin{pmatrix} \cos \theta & 0 & 0 & \sin \theta \\ 0 & \cos \theta & \sin \theta & 0 \\ & & \mathbf{1} & \mathbf{0} \\ 0 & -\sin \theta & \cos \theta & 0 \\ -\sin \theta & 0 & 0 & \cos \theta \\ & & \mathbf{0} & \mathbf{1} \end{pmatrix} \in \{1\} \times \mathbf{Sp}(n)$$

for $\theta \in \mathbf{R}$, where $\mathbf{1}$ (resp. $\mathbf{0}$) denotes the $(n-2) \times (n-2)$ identity (resp. zero) block. We see that the reduced representation $\bar{\rho} : (N/H)^0 \rightarrow \mathbf{O}(V^H)$ is just the direct sum of two copies of the vector representation of $\mathbf{SO}(2)$ on $\mathbf{R}^2$, hence taut. Finally, we conclude that ρ is taut by Proposition 3.19, where we choose L to be generated by the following elements: σ , the diagonal elements in $\mathbf{Sp}(n-2) \subset H$ with ± 1 entries, and the element

$$\left(\begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}, \begin{pmatrix} i & & & \\ & -i & & \\ & & \mathbf{1} & \\ & & & -i \\ & & & & i \\ & & & & & \mathbf{1} \end{pmatrix} \right).$$

Next consider the case of $G = \mathbf{SO}(2) \times \mathbf{Spin}(9)$ acting on $\mathbf{R}^2 \otimes \mathbf{R}^{16}$ by the tensor product of the vector and spin representations. It is equivalent to consider ρ acting on the space of 2×16 real matrices, namely, $\rho(A, B)X = AXB^{-1}$ for $A \in \mathbf{SO}(2)$, $B \in \mathbf{Spin}(9) \subset \mathbf{SO}(16)$, $X \in M(2 \times 16; \mathbf{R})$. We identify $\mathbf{R}^{16} \cong \mathbf{O} \oplus \mathbf{O}$ and use $\{1, e, i, j, k, ei, ej, ek\}$ as a basis of $\mathbf{O}$ over $\mathbf{R}$. For the computations involved in this case, Lemmas 14.61 and 14.77 from [22] are useful, which we state as follows:

7.13 Lemma *A matrix $g \in \mathbf{O}(8) \cong \mathbf{O}(\mathbf{O})$ belongs to $\mathbf{Spin}(7)$ if and only if $g(uv) = g(u)\chi(g)(v)$ for all $u, v \in \mathbf{O}$, where $\chi(g)(v) = g(g^{-1}(1)v)$ for $v \in \mathbf{O}$ defines the vector representation of $\mathbf{Spin}(7)$ on $\mathfrak{SO} \cong \mathbf{R}^7$.*

7.14 Lemma *The subgroup $\mathbf{Spin}(9) \subset \mathbf{SO}(16) \cong \mathbf{SO}(\mathbf{O} \oplus \mathbf{O})$ is generated by the 8-sphere*

$$\left\{ \begin{pmatrix} r & R_u \\ R_{\bar{u}} & -r \end{pmatrix} : r \in \mathbf{R}, u \in \mathbf{O}, r^2 + |u|^2 = 1 \right\}$$

in $\text{End}(\mathbf{O} \oplus \mathbf{O})$, where $R_u : \mathbf{O} \rightarrow \mathbf{O}$ denotes right translation by the element $u \in \mathbf{O}$.

Mainly as a consequence of

$$(7.15) \quad \text{---} \circ \text{---} \circ \text{---} \text{---} \bullet \Big|_{B_3} = (\text{trivial}) \oplus \text{---} \circ \text{---} \circ \text{---} \text{---} \bullet \oplus \text{---} \circ \text{---} \circ \text{---} \text{---} \bullet$$

one can see that the G -orbit space consists of matrices of the form

$$(7.16) \quad \begin{pmatrix} a & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & b & 0 & 0 & 0 & 0 & 0 & 0 & c & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

and, with some more effort (using Lemma 7.14), that the restrictions $a \geq \sqrt{b^2 + c^2}$ and $b, c \geq 0$ apply. Also, Lemma 7.13 shows that

$$g_1 = \text{diag}(-1, 1, 1, 1, -1, -1, -1, 1) \in \mathbf{Spin}(7),$$

which combined with (7.15) yields that

$$g_2 = \begin{pmatrix} 1 & & \\ & \chi(g_1) & \\ & & g_1 \end{pmatrix} \in \mathbf{Spin}(9),$$

where $\chi(g_1) = \text{diag}(-1, -1, -1, 1, 1, 1, -1)$. In particular, the equality $\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} p = pg_2$ where p is as in (7.16) shows that $\mathbf{O}(2) \times \mathbf{Spin}(9)$ has the same orbits as $\mathbf{SO}(2) \times \mathbf{Spin}(9)$. Hereafter we consider $G = \mathbf{O}(2) \times \mathbf{Spin}(9)$.

The principal isotropy group H is isomorphic to $\mathbf{Z}_2 \times \mathbf{Z}_2 \times \mathbf{SU}(3)$, where the first $\mathbf{Z}_2$ corresponds to the kernel of ρ and it is generated by the element $h_1 = (-1, -1) \in \mathbf{O}(2) \times \mathbf{Spin}(9)$, the second $\mathbf{Z}_2$ is generated by the element

$$h_2 = (\text{diag}(1, -1), g_2),$$

and the $\mathbf{SU}(3)$ -factor consists of matrices of the form

$$\begin{pmatrix} 1 & & & & & & & & & & & & & & & \\ & 1 & & & & & & & & & & & & & & \\ & & A & B & & & & & & & & & & & \\ & & -B & A & & & & & & & & & & & \\ & & & & 1 & & & & & & & & & & \\ & & & & & 1 & & & & & & & & & \\ & & & & & & A & B & & & & & & & \\ & & & & & & -B & A & & & & & & & \end{pmatrix} \in \{1\} \times \mathbf{Spin}(9),$$

where $A + iB \in \mathbf{SU}(3)$. Now V^H is the space of matrices of the form

$$\begin{pmatrix} a & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & d & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & b & 0 & 0 & 0 & 0 & 0 & 0 & c & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

whose entries are real, and $(N/H)^0$ is the circle group consisting of the coclasses defined by the following matrices of $\{1\} \times \mathbf{Spin}(9)$:

$$(7.17) \quad \begin{pmatrix} \cos \theta & & & & & & & \sin \theta & & & & & & & & \\ & \cos \theta & & & & & & -\sin \theta & & & & & & & & \\ & & \cos \theta \mathbf{1} & & & & & & & & & & & -\sin \theta \mathbf{1} & & \\ & & & \cos \theta \mathbf{1} & & & & & & & \sin \theta \mathbf{1} & & & & & \\ & & \sin \theta & & & & \cos \theta & & & & & & & & & \\ -\sin \theta & & & & & & & \cos \theta & & & & & & & & \\ & & & & -\sin \theta \mathbf{1} & & & & \cos \theta \mathbf{1} & & & & & & & \\ & & \sin \theta \mathbf{1} & & & & & & & \cos \theta \mathbf{1} & & & & & & \\ & & & & & & & & & & \cos \theta \mathbf{1} & & & & & \end{pmatrix}$$

for $\theta \in \mathbf{R}$, where $\mathbf{1}$ denotes the 3×3 identity block (note that the matrix (7.17) belongs to $\mathbf{Spin}(9)$ because it is the product of the two matrices in $\mathbf{Spin}(9)$ which are obtained by setting $r = \cos \theta$, $u = \sin \theta e$ and $r = 1$, $u = 0$ in Lemma 7.14). We see that the reduced representation $\bar{\rho}$ is just the direct sum of two copies of the vector representation of $\mathbf{SO}(2)$ on $\mathbf{R}^2$, hence taut. We conclude that ρ is taut by Proposition 3.19, where we choose L to be generated by h_1 , h_2 and the elements in the $\mathbf{SU}(3)$ -factor of H with A diagonal with ± 1 entries and $B = 0$.

Finally we consider the case of $G = \mathbf{SU}(2) \times \mathbf{Sp}(n)$, $n \geq 2$, acting on a real form V of $\mathbf{C}^4 \otimes \mathbf{C}^{2n}$. It is equivalent to consider ρ acting on a subspace of $4 \times 2n$ complex matrices, namely, $\rho(A, B)Z = \eta(A)ZB^{-1}$ for $A \in \mathbf{SU}(2)$, $B \in \mathbf{Sp}(n) \subset \mathbf{SU}(2n)$, $Z \in V = \left\{ \begin{pmatrix} X & Y \\ -\bar{Y} & \bar{X} \end{pmatrix} \in M(4 \times 2n; \mathbf{C}) \right\}$, where $\eta : \mathbf{SU}(2) \rightarrow \mathbf{Sp}(2) \subset \mathbf{SU}(4)$ is the 4-dimensional complex representation of $\mathbf{SU}(2)$ given by

$$\eta \begin{pmatrix} \alpha & \beta \\ -\bar{\beta} & \bar{\alpha} \end{pmatrix} = \begin{pmatrix} \alpha^3 & \sqrt{3}\alpha\beta^2 & \beta^3 & \sqrt{3}\alpha^2\bar{\beta} \\ \sqrt{3}\alpha\bar{\beta}^2 & \alpha\bar{\alpha}^2 - 2\bar{\alpha}\beta\bar{\beta} & \sqrt{3}\bar{\alpha}^2\beta & \beta\bar{\beta}^2 - 2\alpha\bar{\alpha}\bar{\beta} \\ -\bar{\beta}^3 & -\sqrt{3}\bar{\alpha}^2\bar{\beta} & \bar{\alpha}^3 & \sqrt{3}\bar{\alpha}\bar{\beta}^2 \\ -\sqrt{3}\alpha^2\bar{\beta} & 2\alpha\bar{\alpha}\beta - \beta^2\bar{\beta} & \sqrt{3}\bar{\alpha}\beta^2 & \alpha^2\bar{\alpha} - 2\alpha\beta\bar{\beta} \end{pmatrix}.$$

Let

$$p_1 = \begin{pmatrix} 1 & 0 & 0 & \dots & 0 & 0 & 0 & 0 & \dots & 0 \\ 0 & 0 & 0 & \dots & 0 & 0 & 0 & 0 & \dots & 0 \\ 0 & 0 & 0 & \dots & 0 & 1 & 0 & 0 & \dots & 0 \\ 0 & 0 & 0 & \dots & 0 & 0 & 0 & 0 & \dots & 0 \end{pmatrix} \in V.$$

Then the orbit $G(p_1)$ is $(4n+1)$ -dimensional and its normal space intersects all orbits at p_1 . So, given an arbitrary orbit M , it contains a point of the form

$$(7.18) \quad \begin{pmatrix} a & 0 & 0 & \dots & 0 & 0 & 0 & 0 & \dots & 0 \\ x_1 & x_2 & x_3 & \dots & x_n & 0 & y_2 & y_3 & \dots & y_n \\ 0 & 0 & 0 & \dots & 0 & a & 0 & 0 & \dots & 0 \\ 0 & -\bar{y}_2 & -\bar{y}_3 & \dots & -\bar{y}_n & \bar{x}_1 & \bar{x}_2 & \bar{x}_3 & \dots & \bar{x}_n \end{pmatrix}$$

where $x_i, y_i \in \mathbf{C}$ and $a \geq 0$, for we can multiply on the left by the minus identity matrix to change the sign of a , if necessary, and remain in the orbit M . The isotropy subgroup $K_1 = G_{p_1}$ is isomorphic to $S^1 \times \mathbf{Sp}(n-1)$, where the circle factor consists of the elements

$$\left(\begin{pmatrix} e^{i\theta} & \\ & e^{-i\theta} \end{pmatrix}, \begin{pmatrix} e^{3i\theta} & & \\ & \mathbf{1} & \\ & & e^{-3i\theta} \\ & & & \mathbf{1} \end{pmatrix} \right)$$

for $\theta \in \mathbf{R}$, where $\mathbf{1}$ denotes the $(n-1) \times (n-1)$ identity block. By letting K_1 act on the matrix (7.18) we now see that the G -orbit space consists of real matrices of the form

$$\begin{pmatrix} a & 0 & 0 & \dots & 0 & 0 & 0 & 0 & \dots & 0 \\ c & b & 0 & \dots & 0 & 0 & 0 & 0 & \dots & 0 \\ 0 & 0 & 0 & \dots & 0 & a & 0 & 0 & \dots & 0 \\ 0 & 0 & 0 & \dots & 0 & c & b & 0 & \dots & 0 \end{pmatrix}.$$

(One can also check that some restrictions on a, b and c apply, but this is unimportant to us.)

The principal isotropy group

$$H = \left\{ \left(q, \begin{pmatrix} \eta(q) & \\ & C \end{pmatrix} \right) : \begin{array}{l} C \in \mathbf{Sp}(n-2), \quad q = \begin{pmatrix} \alpha & \beta \\ -\bar{\beta} & \bar{\alpha} \end{pmatrix}, \\ (\alpha, \beta) = \pm(1, 0), \pm(i, 0), \pm(0, 1), \pm(0, i) \end{array} \right\}$$

is isomorphic to $Q \times \mathbf{Sp}(n-2)$, where Q is quaternion group $\{\pm 1, \pm i, \pm j, \pm k\}$. Now V^H is the space of matrices of the form

$$\begin{pmatrix} a & d & 0 & \dots & 0 & 0 & 0 & 0 & \dots & 0 \\ c & b & 0 & \dots & 0 & 0 & 0 & 0 & \dots & 0 \\ 0 & 0 & 0 & \dots & 0 & a & d & 0 & \dots & 0 \\ 0 & 0 & 0 & \dots & 0 & c & b & 0 & \dots & 0 \end{pmatrix},$$

where the entries are real, and $(N/H)^0$ is the circle group consisting of the coclasses defined by the matrices

$$\begin{pmatrix} \cos \theta & -\sin \theta & & & \\ \sin \theta & \cos \theta & & & \\ & & \mathbf{1} & & \\ & & & \cos \theta & -\sin \theta \\ & & & \sin \theta & \cos \theta \\ & & & & & \mathbf{1} \end{pmatrix} \in \{1\} \times \mathbf{Sp}(n)$$

We see that the reduced representation $\bar{\rho}$ is just the direct sum of two copies of the vector representation of $\mathbf{SO}(2)$ on $\mathbf{R}^2$, hence taut. We conclude that ρ is taut by Proposition 3.19, where we choose L to be generated by Q and the diagonal elements in $\mathbf{Sp}(n-2) \subset H$ with ± 1 entries. $\square$

8 Other applications

In this last section we collect some further applications of our methods.

8.1 Theorem (Dadok) *A polar representation of a compact connected Lie group is orbit equivalent to the isotropy representation of a symmetric space.*

Proof. Conlon's theorem (Theorem 2.3) says that a polar representation is variationally complete. Now use Theorem 6.4. $\square$

The next theorem, which is a special case of a result of Montgomery and Samelson on transitive homeomorphic actions on spheres ([31]), see also [2, 3] by Borel, was used in Sections 5 and 6.

8.2 Theorem *The transitive effective linear actions of compact connected Lie groups G on the spheres S^d are given by the following tables (in each case, the action is the obvious one):*

G	$\mathbf{SO}(n)$	$\mathbf{G}_2$	$\mathbf{Spin}(7)$	$\mathbf{Spin}(9)$	$\mathbf{Sp}(1) \cdot \mathbf{Sp}(n)$
S^d	S^{n-1}	S^6	S^7	S^{15}	S^{4n-1}

G	$\mathbf{SU}(n)$	$S^1 \cdot \mathbf{SU}(n)$	$\mathbf{Sp}(n)$	$S^1 \cdot \mathbf{Sp}(n)$
S^d	S^{2n-1}		S^{4n-1}	

Proof. We may assume that the action is given by a real irreducible representation ρ of G . It is enough to classify ρ up to image equivalence and then factor the kernel out. If ρ is of quaternionic or complex type, then Lemma 5.12 implies that $k(\lambda) = 1$ and we get the standard action of $\mathbf{SO}(2)$ on S^1 and the actions in the second table. If ρ is of real type, then Lemmas 5.11, 5.14 and 5.15 together give the remaining actions in the first table. $\square$

Representations $\rho : G \rightarrow \mathbf{O}(V)$ with cohomogeneity equal to two were classified up to orbit equivalence in [26] by Hsiang and Lawson⁷. It is not difficult to prove that a

⁷The classification of cohomogeneity two and three representations in [26] contain gaps which were to our knowledge first corrected in the papers [44, 46].

representation with cohomogeneity two is polar. Hence it is also taut and if it is irreducible it is therefore in class $\mathcal{O}^2$.

We next show that it is quite easy to see directly that an irreducible representation with cohomogeneity two is in class $\mathcal{O}^2$. We formulate this in the next proposition.

8.3 Proposition *An irreducible representation $\rho : G \rightarrow \mathbf{O}(V)$ with cohomogeneity two is in class $\mathcal{O}^2$.*

Proof. We can clearly restrict our attention to orbits contained in the unit sphere S of V . Let p in S be a point whose orbit has codimension two. The second osculating space $\mathcal{O}_p^2(Gp)$ is the direct sum of $T_p(Gp)$ and the linear span of $\{\alpha(X, Y) \mid X, Y \in T_p(Gp)\}$ where α is the second fundamental form of the orbit Gp as a submanifold of V . It is a general property of submanifolds contained in the sphere S that $\langle \alpha(X, X), \xi \rangle = 1$ for all unit vectors $X \in T_p(Gp)$, where ξ is the inward pointing unit normal vector of S in p . We therefore need to prove that the linear span of $\{\alpha(X, Y) \mid X, Y \in T_p(Gp)\}$ is not one dimensional. Let η be a normal vector of Gp at p that is tangent to the sphere and let A_η^S be the Weingarten map of Gp at p as a submanifold of S . The orbit Gp cannot be umbilical in S since then it would be a round sphere contradicting the assumption that ρ is irreducible. Hence A_η^S has at least two different eigenvalues λ_1 and λ_2 with corresponding eigenvectors X_1 and X_2 that we assume to have length one. Then

$$\begin{aligned} \alpha(X_i, X_i) &= \langle \alpha(X_i, X_i), \xi \rangle \xi + \langle \alpha(X_i, X_i), \eta \rangle \eta \\ &= \xi + \lambda_i \eta \end{aligned}$$

for $i = 1, 2$. It follows that $\alpha(X_1, X_1)$ and $\alpha(X_2, X_2)$ are not linearly dependent and hence that $V = \mathcal{O}_p^2(Gp)$.

Now assume that the orbit through p in S is singular. If we restrict the action of ρ to S , then we have a cohomogeneity one action. It is well known that the slice representations of cohomogeneity one actions are irreducible and transitive on spheres in the normal space. If we now think of the orbit Gp as a submanifold of V , then we see that the slice representation of G_p on the normal space $N_p(Gp)$ decomposes into the trivial representation on $\mathbf{R}\xi$ and an irreducible representation on $N_p^S(Gp)$ where ξ is a normal vector of the sphere S as above and $N_p^S(Gp)$ is the normal space of Gp in S . The linear span of $\{\alpha(X, Y) \mid X, Y \in T_p(Gp)\}$ is clearly invariant under the isotropy representation of G_p . It follows that it contains $N_p^S(Gp)$ since it must have a component in $N_p^S(Gp)$. (If it had no component in $N_p^S(Gp)$, then Gp would be totally geodesic in S , hence a round sphere, contradicting the assumption that ρ is irreducible.) The linear span of $\{\alpha(X, Y) \mid X, Y \in T_p(Gp)\}$ cannot coincide with $N_p^S(Gp)$, since $\alpha(X, X) \notin N_p^S(Gp)$ for a nonvanishing X in $T_p(Gp)$. It follows that $V = \mathcal{O}_p^2(Gp)$. $\square$

8.4 Theorem (Hsiang-Lawson) *The cohomogeneity two linear actions of compact connected Lie groups on the Euclidean spaces are given, up to orbit equivalence, by the isotropy representations of the rank two symmetric spaces.*

Proof. A cohomogeneity two reducible representation is the direct sum of two cohomogeneity one representations and therefore orbit equivalent to the isotropy representation of the product of two spheres.

A cohomogeneity two irreducible representation of a compact connected Lie group is of class $\mathcal{O}^2$ (Proposition 8.3). Therefore the result follows from the fact that no representation listed in Tables C.1, C.2, C.3 and C.4 is of cohomogeneity two, which is easy to check. $\square$


A Description of the maximal subsets of strongly orthogonal roots of the complex simple Lie algebras

In this appendix, for each one of the complex simple Lie algebras, we give a description of the maximal subset of strongly orthogonal roots $\mathcal{B} = \{\beta_1, \dots, \beta_s\}$ which was introduced in Proposition 4.1. In each case, Δ^+ is the positive root system, $\mathcal{S} = \{\alpha_1, \dots, \alpha_n\}$ is the simple root system, and $\lambda_1, \dots, \lambda_r$ are the fundamental highest weights ($\theta_1, \theta_2, \dots$ refer to specific linear forms on the Cartan subalgebra; see e. g. [6]).


$\mathbf{A}_n : \overset{\alpha_1}{\circ} - \overset{\alpha_2}{\circ} - \dots - \overset{\alpha_n}{\circ}$		
$\mathcal{S} = \{\alpha_1 = \theta_1 - \theta_2, \dots, \alpha_n = \theta_n - \theta_{n+1}\}$		
$\Delta^+ = \{\theta_i - \theta_j : 1 \leq i < j \leq n+1\}$		
$\lambda_1 = \theta_1$ $\lambda_2 = \theta_1 + \theta_2$ $\vdots$ $\lambda_n = \theta_1 + \theta_2 + \dots + \theta_n$	$\beta_1 = \theta_1 - \theta_{n+1}$ $\beta_2 = \theta_2 - \theta_n$ $\vdots$ $\beta_n = \theta_{\frac{n}{2}} - \theta_{\frac{n}{2}+2}$ (n even)	$\beta_1 = \theta_1 - \theta_{n+1}$ $\beta_2 = \theta_2 - \theta_n$ $\vdots$ $\beta_n = \theta_{\frac{n+1}{2}} - \theta_{\frac{n+1}{2}+1}$ (n odd)


$\mathbf{B}_n : \overset{\alpha_1}{\circ} - \overset{\alpha_2}{\circ} - \dots - \overset{\alpha_{n-1}}{\circ} \equiv \overset{\alpha_n}{\bullet}$		
$\mathcal{S} = \{\alpha_1 = \theta_1 - \theta_2, \dots, \alpha_{n-1} = \theta_{n-1} - \theta_n, \alpha_n = \theta_n\}$		
$\Delta^+ = \{\theta_i \pm \theta_j : 1 \leq i < j \leq n\} \cup \{\theta_i : 1 \leq n\}$		
$\lambda_1 = \theta_1$ $\lambda_2 = \theta_1 + \theta_2$ $\vdots$ $\lambda_{n-2} = \theta_1 + \theta_2 + \dots + \theta_{n-2}$ $\lambda_{n-1} = \theta_1 + \theta_2 + \dots + \theta_{n-2} + \theta_{n-1}$ $\lambda_n = \frac{1}{2}(\theta_1 + \theta_2 + \dots + \theta_{n-1} + \theta_n)$	$\beta_1 = \theta_1 + \theta_2$ $\beta_2 = \theta_1 - \theta_2$ $\vdots$ $\beta_{n-2} = \theta_{n-2} + \theta_{n-1}$ $\beta_{n-1} = \theta_{n-2} - \theta_{n-1}$ $\beta_n = \theta_n$ (n odd)	$\beta_1 = \theta_1 + \theta_2$ $\beta_2 = \theta_1 - \theta_2$ $\vdots$ $\beta_{n-1} = \theta_{n-1} + \theta_n$ $\beta_n = \theta_{n-1} - \theta_n$ (n even)


$\mathbf{C}_n : \overset{\alpha_1}{\bullet} - \overset{\alpha_2}{\bullet} - \dots - \overset{\alpha_{n-1}}{\bullet} \equiv \overset{\alpha_n}{\circ}$	
$\mathcal{S} = \{\alpha_1 = \theta_1 - \theta_2, \dots, \alpha_{n-1} = \theta_{n-1} - \theta_n, \alpha_n = 2\theta_n\}$	
$\Delta^+ = \{\theta_i \pm \theta_j : 1 \leq i < j \leq n\} \cup \{2\theta_i : 1 \leq n\}$	
$\lambda_1 = \theta_1$ $\lambda_2 = \theta_1 + \theta_2$ $\vdots$ $\lambda_n = \theta_1 + \theta_2 + \dots + \theta_n$	$\beta_1 = 2\theta_1$ $\beta_2 = 2\theta_2$ $\vdots$ $\beta_n = 2\theta_n$


$\mathbf{D}_n :$ 		
$\mathcal{S} = \{\alpha_1 = \theta_1 - \theta_2, \dots, \alpha_{n-2} = \theta_{n-2} - \theta_{n-1}, \alpha_{n-1} = \theta_{n-1} - \theta_n, \alpha_n = \theta_{n-1} + \theta_n\}$		
$\Delta^+ = \{\theta_i \pm \theta_j : 1 \leq i < j \leq n\}$		
$\lambda_1 = \theta_1$ $\lambda_2 = \theta_1 + \theta_2$ $\vdots$ $\lambda_{n-2} = \theta_1 + \theta_2 + \dots + \theta_{n-2}$ $\lambda_{n-1} = \frac{1}{2}(\theta_1 + \theta_2 + \dots + \theta_{n-1} - \theta_n)$ $\lambda_n = \frac{1}{2}(\theta_1 + \theta_2 + \dots + \theta_{n-1} + \theta_n)$	$\beta_1 = \theta_1 + \theta_2$ $\beta_2 = \theta_1 - \theta_2$ $\vdots$ $\beta_{n-2} = \theta_{n-2} + \theta_{n-1}$ $\beta_{n-1} = \theta_{n-2} - \theta_{n-1}$ (n odd)	$\beta_1 = \theta_1 + \theta_2$ $\beta_2 = \theta_1 - \theta_2$ $\vdots$ $\beta_{n-1} = \theta_{n-1} + \theta_n$ $\beta_n = \theta_{n-1} - \theta_n$ (n even)

$\mathbf{G}_2 :$ 	
$\Delta^+ = \{\alpha_1, \alpha_2, \alpha_1 + \alpha_2, 2\alpha_1 + \alpha_2, 3\alpha_1 + \alpha_2, 3\alpha_1 + 2\alpha_2\}$	
$\lambda_1 = 2\alpha_1 + \alpha_2$ $\lambda_2 = 3\alpha_1 + 2\alpha_2$	$\beta_1 = 3\alpha_1 + 2\alpha_2$ $\beta_2 = \alpha_1$

$\mathbf{F}_4 :$ 	
$\mathcal{S} = \{\alpha_1 = \theta_2 - \theta_3, \alpha_2 = \theta_3 - \theta_4, \alpha_3 = \theta_4, \alpha_4 = \frac{1}{2}(\theta_1 - \theta_2 - \theta_3 - \theta_4)\}$	
$\Delta^+ = \{\theta_i : 1 \leq i \leq 4\} \cup \{\theta_i \pm \theta_j : 1 \leq i < j \leq 4\} \cup \{\frac{1}{2}(\theta_1 \pm \theta_2 \pm \theta_3 \pm \theta_4)\}$	
$\lambda_1 = \theta_1 + \theta_2$ $\lambda_2 = 2\theta_1 + \theta_2 + \theta_3$ $\lambda_3 = \frac{1}{2}(3\theta_1 + \theta_2 + \theta_3 + \theta_4)$ $\lambda_4 = \theta_1$	$\beta_1 = \theta_1 + \theta_2$ $\beta_2 = \theta_1 - \theta_2$ $\beta_3 = \theta_3 + \theta_4$ $\beta_4 = \theta_3 - \theta_4$

$\mathbf{E}_6 :$ 	
$\mathcal{S} = \{\alpha_1 = \frac{1}{2}(\theta_1 - \theta_2 - \theta_3 - \theta_4 - \theta_5 - \theta_6 - \theta_7 + \theta_8),$ $\alpha_2 = \theta_1 + \theta_2, \alpha_3 = \theta_2 - \theta_1, \alpha_4 = \theta_3 - \theta_2, \alpha_5 = \theta_4 - \theta_3, \alpha_6 = \theta_5 - \theta_4\}$	
$\Delta^+ = \{\pm\theta_i + \theta_j : 1 \leq i < j \leq 5\} \cup \{\frac{1}{2}(\theta_8 - \theta_7 - \theta_6 + \sum_{i=1}^5 \epsilon_i \theta_i) : \prod_{i=1}^5 \epsilon_i = +1\}$	
$\lambda_1 = \frac{2}{3}(\theta_8 - \theta_7 - \theta_6)$ $\lambda_2 = \frac{1}{2}(\theta_1 + \theta_2 + \theta_3 + \theta_4 + \theta_5 - \theta_6 - \theta_7 + \theta_8)$ $\lambda_3 = \frac{5}{6}(\theta_8 - \theta_7 - \theta_6) + \frac{1}{2}(-\theta_1 + \theta_2 + \theta_3 + \theta_4 + \theta_5)$ $\lambda_4 = \theta_3 + \theta_4 + \theta_5 - \theta_6 - \theta_7 + \theta_8$ $\lambda_5 = \frac{2}{3}(\theta_8 - \theta_7 - \theta_6) + \theta_4 + \theta_5$ $\lambda_6 = \frac{1}{3}(\theta_8 - \theta_7 - \theta_6) + \theta_5$	$\beta_1 = \frac{1}{2}(\theta_1 + \theta_2 + \theta_3 + \theta_4 + \theta_5 - \theta_6 - \theta_7 + \theta_8)$ $\beta_2 = \frac{1}{2}(-\theta_1 - \theta_2 - \theta_3 - \theta_4 + \theta_5 - \theta_6 - \theta_7 + \theta_8)$ $\beta_3 = \theta_4 - \theta_1$ $\beta_4 = \theta_3 - \theta_2$

$\mathbf{E}_7 :$ 	
$\mathcal{S} = \{\alpha_1 = \frac{1}{2}(\theta_1 - \theta_2 - \theta_3 - \theta_4 - \theta_5 - \theta_6 - \theta_7 + \theta_8),$ $\alpha_2 = \theta_1 + \theta_2, \alpha_3 = \theta_2 - \theta_1, \alpha_4 = \theta_3 - \theta_2, \alpha_5 = \theta_4 - \theta_3, \alpha_6 = \theta_5 - \theta_4, \alpha_7 = \theta_6 - \theta_5\}$	
$\Delta^+ = \{\pm\theta_i + \theta_j : 1 \leq i < j \leq 6\} \cup \{\theta_8 - \theta_7\} \cup \{\frac{1}{2}(\theta_8 - \theta_7 + \sum_{i=1}^6 \epsilon_i \theta_i) : \prod_{i=1}^6 \epsilon_i = -1\}$	
$\lambda_1 = \theta_8 - \theta_7$ $\lambda_2 = \frac{1}{2}(\theta_1 + \theta_2 + \theta_3 + \theta_4 + \theta_5 + \theta_6 - 2\theta_7 + 2\theta_8)$ $\lambda_3 = \frac{1}{2}(-\theta_1 + \theta_2 + \theta_3 + \theta_4 + \theta_5 + \theta_6 - 3\theta_7 + 3\theta_8)$ $\lambda_4 = \theta_3 + \theta_4 + \theta_5 + \theta_6 + 2(\theta_8 - \theta_7)$ $\lambda_5 = \frac{1}{2}(2\theta_4 + 2\theta_5 + 2\theta_6 + 3(\theta_8 - \theta_7))$ $\lambda_6 = \theta_5 + \theta_6 - \theta_7 + \theta_8$ $\lambda_7 = \theta_6 + \frac{1}{2}(\theta_8 - \theta_7)$	$\beta_1 = \theta_8 - \theta_7$ $\beta_2 = \theta_6 + \theta_5$ $\beta_3 = \theta_6 - \theta_5$ $\beta_4 = \theta_4 + \theta_3$ $\beta_5 = \theta_4 - \theta_3$ $\beta_6 = \theta_2 + \theta_1$ $\beta_7 = \theta_2 - \theta_1$

$\mathbf{E}_8 :$ 	
$\mathcal{S} = \{\alpha_1 = \frac{1}{2}(\theta_1 - \theta_2 - \theta_3 - \theta_4 - \theta_5 - \theta_6 - \theta_7 + \theta_8), \alpha_2 = \theta_1 + \theta_2,$ $\alpha_3 = \theta_2 - \theta_1, \alpha_4 = \theta_3 - \theta_2, \alpha_5 = \theta_4 - \theta_3, \alpha_6 = \theta_5 - \theta_4, \alpha_7 = \theta_6 - \theta_5, \alpha_8 = \theta_7 - \theta_6\}$	
$\Delta^+ = \{\pm\theta_i + \theta_j : 1 \leq i < j \leq 8\} \cup \{\frac{1}{2}(\theta_8 + \sum_{i=1}^7 \epsilon_i \theta_i) : \prod_{i=1}^7 \epsilon_i = +1\}$	
$\lambda_1 = 2\theta_8$ $\lambda_2 = \frac{1}{2}(\theta_1 + \theta_2 + \theta_3 + \theta_4 + \theta_5 + \theta_6 + \theta_7 + 5\theta_8)$ $\lambda_3 = \frac{1}{2}(-\theta_1 + \theta_2 + \theta_3 + \theta_4 + \theta_5 + \theta_6 + \theta_7 + 7\theta_8)$ $\lambda_4 = \theta_3 + \theta_4 + \theta_5 + \theta_6 + \theta_7 + 5\theta_8$ $\lambda_5 = \theta_4 + \theta_5 + \theta_6 + \theta_7 + 4\theta_8$ $\lambda_6 = \theta_5 + \theta_6 + \theta_7 + 3\theta_8$ $\lambda_7 = \theta_6 + \theta_7 + 2\theta_8$ $\lambda_8 = \theta_7 + \theta_8$	$\beta_1 = \theta_7 + \theta_8$ $\beta_2 = \theta_8 - \theta_7$ $\beta_3 = \theta_6 + \theta_5$ $\beta_4 = \theta_6 - \theta_5$ $\beta_5 = \theta_4 + \theta_3$ $\beta_6 = \theta_4 - \theta_3$ $\beta_7 = \theta_2 + \theta_1$ $\beta_8 = \theta_2 - \theta_1$

B Table for Dadok's invariant $k(\lambda)$

We next tabulate the value of the Dadok invariant $k(\lambda)$ for the the complex irreducible representations π_λ of the complex simple Lie algebras, where λ is a fundamental highest weight (since $k(\lambda)$ is linear on λ , this already determines the value of $k(\lambda)$ for all λ). The number in parenthesis next to the vertex of the Dynkin diagram which specifies α_i is $k(\lambda_i)$. The last column tells, for each complex simple Lie algebra, whether all representations are self-dual, or else, the dual of π_{λ_i} is given by π_{λ_j} , where the vertices that specify α_i and α_j are images one of the other under the symmetry of the Dynkin diagram.

G	$k(\lambda)$	$Conditions$
A_{2n-1}		duality given by symmetry of diagram
A_{2n}		duality given by symmetry of diagram
B_{2n-1}		all self-dual
B_{2n}		all self-dual
C_n		all self-dual
D_{2n}		all self-dual
D_{2n+1}		duality given by symmetry of diagram
G_2		all self-dual
F_4		all self-dual
E_6		duality given by symmetry of diagram
E_7		all self-dual
E_8		all self-dual

Table B.1: The invariant of Dadok.

C Table of candidates to class $\mathcal{O}^2$

In the tables below are compiled all the real irreducible representations of compact connected Lie groups that are not orbit equivalent to the isotropy representation of a symmetric space but that have a chance of being a representation of class $\mathcal{O}^2$. These are just the representations mentioned in Propositions 5.1, 5.2, 5.17, 5.18, 5.19, 5.20 and 5.21, except those excluded by the discussion in Section 6; note however that the presentation here slightly differs from the presentation in the above mentioned propositions for the sake of convenience.

G	ρ	<i>Conditions</i>
$\mathbf{SO}(2) \times \mathbf{Spin}(9)$	$(x \oplus x^{-1}) \otimes \overset{1}{\circ} \text{---} \overset{1}{\circ} \text{---} \overset{1}{\circ} \text{---} \bullet$	—
$\mathbf{U}(2) \times \mathbf{Sp}(n)$	$(x \oplus x^{-1}) \otimes \overset{1}{\circ} \otimes \overset{1}{\bullet} \text{---} \dots \text{---} \bullet \text{---} \bullet$	$n \geq 2$
$\mathbf{SU}(2) \times \mathbf{Sp}(n)$	$\overset{3}{\circ} \otimes \overset{1}{\bullet} \text{---} \dots \text{---} \bullet \text{---} \bullet$	$n \geq 2$

Table C.1: Taut representations.

G	ρ	Conditions
$\mathrm{SU}(n) \times \mathrm{Sp}(m)$		$n \geq 3, m \geq 2$
$\mathrm{U}(n) \times \mathrm{Sp}(m)$		$n \geq 3, m \geq 2$
$\mathrm{SO}(m) \times \mathrm{Sp}(1) \cdot \mathrm{Sp}(n)$ ($n \geq 2$)		$m = 2p > 4$ $m = 2p + 1 > 3$ $m = 4$ $m = 3$
$\mathrm{SO}(m) \times \mathrm{G}_2$		$m = 2p > 4$ $m = 2p + 1 > 3$ $m = 4$ $m = 3$
$\mathrm{SO}(m) \times \mathrm{Spin}(7)$		$m = 2p > 4$ $m = 2p + 1 \geq 5$ $m = 4$
$\mathrm{SO}(m) \times \mathrm{Spin}(9)$		$m = 2p > 4$ $m = 2p + 1 > 3$ $m = 4$ $m = 3$

Table C.2: Nontaut representations (first group).

G	ρ	Conditions
$\mathbf{G}_2 \times \mathbf{G}_2$		—
$\mathbf{G}_2 \times \mathbf{Spin}(7)$		—
$\mathbf{G}_2 \times \mathbf{Spin}(9)$		—
$\mathbf{Spin}(7) \times \mathbf{Spin}(7)$		—
$\mathbf{Spin}(7) \times \mathbf{Spin}(9)$		—
$\mathbf{Sp}(1) \cdot \mathbf{Sp}(n) \times \mathbf{G}_2$		$n \geq 2$
$\mathbf{Sp}(1) \cdot \mathbf{Sp}(n) \times \mathbf{Spin}(7)$		$n \geq 2$

Table C.3: Nontaut representations (second group).

G	ρ	Conditions
$\mathbf{Sp}(n) \times \mathbf{SU}(6)$		$n \geq 2$
$\mathbf{Sp}(n) \times \mathbf{Spin}(11)$		—
$\mathbf{Sp}(n) \times \mathbf{Spin}(12)$		$n \geq 2$
$\mathbf{Sp}(1) \times \mathbf{Spin}(13)$		—
$\mathbf{Sp}(1) \times \mathbf{Sp}(2)$		—
$\mathbf{Sp}(n) \times \mathbf{Sp}(3)$		$n \geq 2$
$\mathbf{Sp}(n) \times \mathbf{E}_7$		$n \geq 2$
$\mathbf{Spin}(7)$		—
$\mathbf{Spin}(9)$		—
$\mathbf{Spin}(15)$		—
$\mathbf{Spin}(17)$		—

Table C.4: Nontaut representations (third group).

References

- [1] I. Bergmann, *Polar actions*, Ph.D. thesis, Univ. Augsburg, 1999.
- [2] A. Borel, *Some remarks about Lie groups transitive on spheres and tori*, Bull. Amer. Math. Soc. **55** (1949), 580–587.
- [3] ———, *Le plan projectif des octaves et les sphères comme espaces homogènes*, C. R. Acad. Sc. Paris **230** (1950), 1378–1380.
- [4] R. Bott, *An application of the Morse theory to the topology of Lie groups*, Bull. Soc. Math. France **84** (1956), 251–281.
- [5] R. Bott and H. Samelson, *Applications of the theory of Morse to symmetric spaces*, Amer. J. Math. **80** (1958), 964–1029, Correction in Amer. J. Math. **83** (1961), 207–208.
- [6] N. Bourbaki, *Groupes et algèbres de Lie*, Éléments de Mathématique, Hermann, Paris, 1968.
- [7] G. E. Bredon, *Introduction to compact transformation groups*, Pure and Applied Mathematics, vol. 46, Academic Press, 1972.
- [8] T. Cecil and P. Ryan, *Tight and taut immersions of manifolds*, Research Notes in Mathematics, no. 107, Pitman, 1985.
- [9] L. Conlon, *Variational completeness and K -transversal domains*, J. Differential Geom. **5** (1971), 135–147.
- [10] S. Console and A. Fino, *Symmetric weights and s -representations*, Kodai Math. J. **23** (2000), 266–280.
- [11] S. Console and G. Thorbergsson, *Geometric characterizations of orthogonal representations*, Geometry and Topology of Submanifolds, VIII (F. Dillen et al, ed.), World Scientific, 1996, pp. 74–84.
- [12] J. Dadok, *Polar actions induced by actions of compact Lie groups*, Trans. Amer. Math. Soc. **288** (1985), 125–137.
- [13] A. J. Di Scala and C. Olmos, *Variationally complete representations are polar*, preprint.
- [14] J. J. Duistermaat, *Convexity and tightness for restrictions of Hamiltonian functions to fixed point sets of an antisymplectic involution*, Trans. Amer. Math. Soc. **275** (1983), 417–429.
- [15] J. Eschenburg and E. Heintze, *On the classification of polar representations*, Math. Z. **232** (1999), 391–398.
- [16] ———, *Polar representations and symmetric spaces*, J. Reine. Angew. Math. **507** (1999), 93–106.

- [17] E.E. Floyd, *On periodic maps and the Euler characteristics of associated spaces*, Trans. Amer. Math. Soc. **72** (1952), 138–147.
- [18] W. Fulton and J. Harris, *Representation theory: A first course*, Graduate Texts in Mathematics, no. 129, Springer-Verlag, 1991.
- [19] B. Galemann, *Tautness and linear representations of the classical compact groups*, Ph.D. thesis, Univ. of Notre Dame, 1993.
- [20] C. Gorodski and G. Thorbergsson, *Cycles of bott-samelson type for taut representations*, preprint.
- [21] K. Grove and C. Searle, *Global G -manifold reductions and resolutions*, Ann. Global Anal. and Geom. **18** (2000), 437–446, Special issue in memory of Alfred Gray (1939–1998).
- [22] F. R. Harvey, *Spinors and calibrations*, Perspectives in Mathematics, Academic Press, 1990.
- [23] E. Heintze, R. S. Palais, C.-L. Terng, and G. Thorbergsson, *Hyperpolar actions and k -flat homogeneous spaces*, J. Reine Angew. Math. **454** (1994), 163–179.
- [24] S. Helgason, *Differential geometry, Lie groups, and symmetric spaces*, Pure and Applied Mathematics, no. 80, Academic Press, 1978.
- [25] H. Hopf, *Über die Topologie von Gruppen-Mannigfaltigkeiten und ihrer Verallgemeinerungen*, Ann. of Math. **42** (1941), 22–52.
- [26] W.-Y. Hsiang and H. B. Lawson Jr., *Minimal submanifolds of low cohomogeneity*, J. Differential Geom. **5** (1971), 1–38.
- [27] N. H. Kuiper, *Sur les immersions à courbure totale minimale*, Séminaire de Topologie et Géométrie Différentielle C. Ereshmann, Paris, vol. II, 1961, Recueil d'exposés faits en 1958-1959-1960.
- [28] D. Luna, *Adhérences d'orbite et invariants*, Invent. Math. **29** (1975), 231–238.
- [29] D. Luna and R. W. Richardson, *A generalization of the Chevalley restriction theorem*, Duke Math. J. **46** (1979), 487–496.
- [30] R. Miyaoka, *Taut embeddings and Dupin hypersurfaces*, Differential Geometry of Submanifolds (Kyoto, 1984) (K. Kenmotsu, ed.), Lecture Notes in Math., no. 1090, Springer, Berlin, 1984, pp. 15–23.
- [31] D. Montgomery and H. Samelson, *Transformation groups of spheres*, Ann. of Math. **44** (1943), 454–470.
- [32] C. Olmos, *Homogeneous submanifolds of higher rank and parallel mean curvature*, J. Differential Geom. **39** (1994), 605–627.

- [33] A. L. Onishchik and E. B. Vinberg, *Lie groups and algebraic groups*, Springer-Verlag, 1990.
- [34] T. Ozawa, *On the critical sets of distance functions to a taut submanifold*, Math. Ann. **276** (1986), 91–96.
- [35] R. S. Palais and C.-L. Terng, *Critical point theory and submanifold geometry*, Lect. Notes in Math., no. 1353, Springer-Verlag, 1988.
- [36] U. Pinkall, *Curvature properties of taut submanifolds*, Geom. Dedicata **20** (1986), 79–83.
- [37] G. W. Schwartz, *Lifting smooth homotopies of orbit spaces*, I.H.E.S. Publ. in Math. **51** (1980), 37–135.
- [38] T. Skjelbred and E. Straume, *A note on the reduction principle for compact transformation groups*, preprint, 1995.
- [39] E. Straume, *On the invariant theory and geometry of compact linear groups of cohomogeneity ≤ 3* , Diff. Geom. and its Appl. **4** (1994), 1–23.
- [40] ———, *Compact connected lie transformation groups on spheres with low cohomogeneity, I*, Memoirs, no. 569, Amer. Math. Soc., 1996.
- [41] H. Tasaki and O. Yasukura, *R-spaces associated with a Hermitian symmetric pair*, Tsukuba J. Math. **10** (1986), 165–170.
- [42] C.-L. Terng, *Variational completeness and infinite dimensional geometry*, Geometry and Topology of Submanifolds, III (L. Vestraelen and A. West, eds.), World Scientific, 1991, pp. 279–293.
- [43] C.-L. Terng and G. Thorbergsson, *Taut immersions into complete Riemannian manifolds*, Tight and Taut Submanifolds (T. E. Ryan and S.-S. Chern, eds.), Math. Sci. Res. Inst. Publ. 32, Cambridge University Press, 1997, pp. 181–228.
- [44] F. Uchida, *An orthogonal transformation group of $(8k - 1)$ -sphere*, J. Differential Geom. **15** (1980), 569–574.
- [45] J.A. Wolf, *Spaces of constant curvature*, 5th ed., Publish or Perish, Houston, 1984.
- [46] O. Yasukura, *A classification of orthogonal transformation groups of low cohomogeneity*, Tsukuba J. Math. **10** (1986), 299–326.